

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
4 October 2001 (04.10.2001)

PCT

(10) International Publication Number
WO 01/72774 A2

- (51) International Patent Classification⁷: **C07K 14/00**
- (21) International Application Number: PCT/GB01/01297
- (22) International Filing Date: 23 March 2001 (23.03.2001)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
0007268.6 24 March 2000 (24.03.2000) GB
- (71) Applicant (for all designated States except US): **CYCLACEL LIMITED** [GB/GB]; Dundee Technopole, James Lindsay Place, Dundee DD1 5JJ (GB).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **DEAK, Peter** [HU/GB]; 27 George Nuttall Close, Cambridge CB4 1YE (GB). **GLOVER, David, Moore** [GB/GB]; Vincent Cottage, 20 Fox Street, Great Gransdon, Sandy, Bedfordshire SG19 3AA (GB). **MIDGLEY, Carol** [GB/GB]; Daisy Cottage, 9 Mount Pleasant, Aspley Guise, Milton Keynes MK17 8JZ (GB).
- (74) Agents: **KHOO, Chong-Yee** et al.; D Young & Co., 21 New Fetter Lane, London EC4A 1DA (GB).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:
— without international search report and to be republished upon receipt of that report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 01/72774 A2

(54) Title: CELL CYCLE PROGRESSION PROTEINS

(57) Abstract: Polynucleotides encoding a number of *Drosophila* gene products are provided. Polynucleotide probes derived from these nucleotide sequences, polypeptides encoded by the polynucleotides and antibodies that bind to the polypeptides are also provided.

CELL CYCLE PROGRESSION PROTEINS

The present invention relates to a number of genes implicated in the processes of cell cycle progression, including mitosis and meiosis.

We have now identified a large number of genes in *Drosophila*, mutations in
5 which disrupt cell cycle progression, for example the processes of mitosis and/or meiosis. We have determined the phenotypes of these mutations and recovered nucleotide sequences associated with the corresponding genes. Many of these nucleotide sequences correspond to protein open reading frames (ORFs) present in the *Drosophila* genome.

Accordingly the present invention provides in one aspect a polynucleotide selected
10 from: (a) polynucleotides comprising any one of the nucleotide sequences set out in Examples 1 to 70 or the complement thereof; (b) polynucleotides comprising a nucleotide sequence capable of hybridising to the nucleotide sequences set out in Examples 1 to 70, or a fragment thereof; (c) polynucleotides comprising a nucleotide sequence capable of hybridising to the complement of the nucleotide sequences set out in Examples 1 to 70 or
15 a fragment thereof; (d) polynucleotides comprising a polynucleotide sequence which is degenerate as a result of the genetic code to the polynucleotides defined in (a), (b) or (c).

There is provided, according to another aspect of the present invention, a polynucleotide selected from: (a) polynucleotides comprising any one of the nucleotide sequences set out in Examples 1 to 14 or the complement thereof; (b) polynucleotides
20 comprising a nucleotide sequence capable of hybridising to the nucleotide sequences set out in Examples 1 to 14, or a fragment thereof; (c) polynucleotides comprising a nucleotide sequence capable of hybridising to the complement of the nucleotide sequences set out in Examples 1 to 14 or a fragment thereof; (d) polynucleotides comprising a polynucleotide sequence which is degenerate as a result of the genetic code to the polynucleotides defined
25 in (a), (b) or (c).

We provide, according to yet a further aspect of the present invention, a polynucleotide selected from: (a) polynucleotides comprising any one of the nucleotide

sequences set out in Examples 15 to 19 or the complement thereof; (b) polynucleotides comprising a nucleotide sequence capable of hybridising to the nucleotide sequences set out in Examples 15 to 19, or a fragment thereof; (c) polynucleotides comprising a nucleotide sequence capable of hybridising to the complement of the nucleotide sequences set out in
5 Examples 15 to 19 or a fragment thereof; (d) polynucleotides comprising a polynucleotide sequence which is degenerate as a result of the genetic code to the polynucleotides defined in (a), (b) or (c).

As a further aspect of the present invention, there is provided a polynucleotide selected from: (a) polynucleotides comprising any one of the nucleotide sequences set out
10 in Examples 20 to 30 or the complement thereof; (b) polynucleotides comprising a nucleotide sequence capable of hybridising to the nucleotide sequences set out in Examples 20 to 30, or a fragment thereof; (c) polynucleotides comprising a nucleotide sequence capable of hybridising to the complement of the nucleotide sequences set out in Examples 20 to 30 or a fragment thereof; (d) polynucleotides comprising a polynucleotide sequence
15 which is degenerate as a result of the genetic code to the polynucleotides defined in (a), (b) or (c).

We provide, according to a yet further aspect of the present invention, a polynucleotide selected from: (a) polynucleotides comprising any one of the nucleotide sequences set out in Examples 31 to 53 or the complement thereof; (b) polynucleotides
20 comprising a nucleotide sequence capable of hybridising to the nucleotide sequences set out in Examples 31 to 53, or a fragment thereof; (c) polynucleotides comprising a nucleotide sequence capable of hybridising to the complement of the nucleotide sequences set out in 31 to 53 or a fragment thereof; (d) polynucleotides comprising a polynucleotide sequence which is degenerate as a result of the genetic code to the polynucleotides defined in (a),
25 (b) or (c).

The present invention, in a further aspect, provides a polynucleotide selected from: (a) polynucleotides comprising any one of the nucleotide sequences set out in 54 to 70 or the complement thereof; (b) polynucleotides comprising a nucleotide sequence capable of hybridising to the nucleotide sequences set out in 54 to 70, or a fragment thereof; (c)

polynucleotides comprising a nucleotide sequence capable of hybridising to the complement of the nucleotide sequences set out in 54 to 70 or a fragment thereof; (d) polynucleotides comprising a polynucleotide sequence which is degenerate as a result of the genetic code to the polynucleotides defined in (a), (b) or (c).

- 5 A polynucleotide probe which comprises a fragment of at least 15 nucleotides of a polynucleotide according to any of the above aspects of the invention.

 The present invention also provides a polypeptide which comprises any one of the amino acid sequences set out in Examples 1 to 70 or in any of Examples 1 to 14, Examples 15 to 19, Examples 20 to 30, Examples 31 to 53 and Examples 54 to 70, or a
10 homologue, variant, derivative or fragment thereof.

 Preferably the polypeptide is encoded by a cDNA sequence obtainable from a eukaryotic cDNA library, preferably a metazoan cDNA library (such as insect or mammalian) said DNA sequence comprising a DNA sequence being selectively detectable with a *Drosophila* nucleotide sequence as shown in any one of Examples 1 to 70.

- 15 The term "selectively detectable" means that the cDNA used as a probe is used under conditions where a target cDNA of the invention is found to hybridize to the probe at a level significantly above background. The background hybridization may occur because of other cDNAs present in the cDNA library. In this event background implies a level of signal generated by interaction between the probe and a non-specific cDNA
20 member of the library which is less than 10 fold, preferably less than 100 fold as intense as the specific interaction observed with the target cDNA. The intensity of interaction may be measured, for example, by radiolabelling the probe, e.g. with ³²P. Suitable conditions may be found by reference to the Examples, as well as in the detailed description below.

 A polynucleotide encoding a polypeptide of the invention is also provided.

- 25 The present invention further provides a vector comprising a polynucleotide of the invention, for example an expression vector comprising a polynucleotide of the invention

operably linked to a regulatory sequence capable of directing expression of said polynucleotide in a host cell.

Also provided is an antibody capable of binding a polypeptide of the invention.

5 In a further aspect the present invention provides a method for detecting the presence or absence of a polynucleotide of the invention in a biological sample which method comprises: (a) bringing the biological sample containing DNA or RNA into contact with a probe comprising a nucleotide of the invention under hybridising conditions; and (b) detecting any duplex formed between the probe and nucleic acid in the sample.

10 In another aspect the invention provides a method for detecting a polypeptide of the invention present in a biological sample which comprises: (a) providing an antibody of the invention; (b) incubating a biological sample with said antibody under conditions which allow for the formation of an antibody-antigen complex; and (c) determining whether antibody-antigen complex comprising said antibody is formed.

15 Knowledge of the genes involved in cell cycle progression allows the development of therapeutic agents for the treatment of medical conditions associated with aberrant cell cycle progression. Accordingly, the present invention provides a polynucleotide of the invention for use in therapy. The present invention also provides a polypeptide of the invention for use in therapy. The present invention further provides an antibody of the
20 invention for use in therapy.

In a specific embodiment, the present invention provides a method of treating a tumour or a patient suffering from a proliferative disease, comprising administering to a patient in need of treatment an effective amount of a polynucleotide, polypeptide and/or antibody of the invention.

25 The present invention also provides the use of a polypeptide of the invention in a method of identifying a substance capable of affecting the function of the corresponding

gene. For example, in one embodiment the present invention provides the use of a polypeptide of the invention in an assay for identifying a substance capable of inhibiting cell cycle progression. The substance may inhibit any of the steps or stages in the cell cycle, for example, formation of the nuclear envelope, exit from the quiescent phase of the cell cycle (G0), G1 progression, chromosome decondensation, nuclear envelope breakdown, START, initiation of DNA replication, progression of DNA replication, termination of DNA replication, centrosome duplication, G2 progression, activation of mitotic or meiotic functions, chromosome condensation, centrosome separation, microtubule nucleation, spindle formation and function, interactions with microtubule motor proteins, chromatid separation and segregation, inactivation of mitotic functions, formation of contractile ring, and cytokinesis functions. For example, possible functions of genes of the invention for which it may be desired to identify substances which affect such functions include chromatin binding, formation of replication complexes, replication licensing, phosphorylation or other secondary modification activity, proteolytic degradation, microtubule binding, actin binding, septin binding, microtubule organising centre nucleation activity and binding to components of cell cycle signalling pathways.

In a further aspect the present invention provides a method for identifying a substance capable of binding to a polypeptide of the invention, which method comprises incubating the polypeptide with a candidate substance under suitable conditions and determining whether the substance binds to the polypeptide.

In an additional aspect, the invention provides kits comprising polynucleotides, polypeptides or antibodies of the invention and methods of using such kits in diagnosing the presence of absence of polynucleotides and polypeptides of the invention including deleterious mutant forms.

Also provided is a substance identified by the above methods of the invention. Such substances may be used in a method of therapy, such as in a method of affecting cell cycle progression, for example mitosis and/or meiosis.

The invention also provides a process comprising the steps of: (a) performing one of the above methods; and (b) preparing a quantity of those one or more substances identified as being capable of binding to a polypeptide of the invention.

Also provided is a process comprising the steps of: (a) performing one of the above
5 methods; and (b) preparing a pharmaceutical composition comprising one or more substances identified as being capable of binding to a polypeptide of the invention.

We further provide a method for identifying a substance capable of modulating the function of a polypeptide of the invention or a polypeptide encoded by a polynucleotide of the invention, the method comprising the steps of: incubating the polypeptide with a
10 candidate substance and determining whether activity of the polypeptide is thereby modulated.

A substance identified by a method or assay according to any of the above methods or processes is also provided, as is the use of such a substance in a method of inhibiting the function of a polypeptide. Use of such a substance in a method of regulating a cell
15 division cycle function is also provided.

DETAILED DESCRIPTION OF THE INVENTION

The practice of the present invention will employ, unless otherwise indicated, conventional techniques of chemistry, molecular biology, microbiology, recombinant DNA and immunology, which are within the capabilities of a person of ordinary skill in
20 the art. Such techniques are explained in the literature. See, for example, J. Sambrook, E. F. Fritsch, and T. Maniatis, 1989, *Molecular Cloning: A Laboratory Manual*, Second Edition, Books 1-3, Cold Spring Harbor Laboratory Press; Ausubel, F. M. et al. (1995 and periodic supplements; *Current Protocols in Molecular Biology*, ch. 9, 13, and 16, John Wiley & Sons, New York, N.Y.); B. Roe, J. Crabtree, and A. Kahn, 1996, *DNA Isolation and Sequencing: Essential Techniques*, John Wiley & Sons; J. M. Polak and James O'D.
25 McGee, 1990, *In Situ Hybridization: Principles and Practice*, Oxford University Press; M. J. Gait (Editor), 1984, *Oligonucleotide Synthesis: A Practical Approach*, Irl Press; and, D.

M. J. Lilley and J. E. Dahlberg, 1992, *Methods of Enzymology: DNA Structure Part A: Synthesis and Physical Analysis of DNA* Methods in Enzymology, Academic Press. Each of these general texts is herein incorporated by reference.

Preferably, the polypeptides and polynucleotides of the invention are such that they
 5 give rise to or are associated with defined phenotypes when mutated.

For example, mutations in the polypeptides and polynucleotides of the invention may be associated with a failure to complete cytokinesis; such polypeptides and polynucleotides are conveniently categorised as "Category 1". Phenotypes associated with Category 1 polypeptides and polynucleotides include any one or more of the following,
 10 singly or in combination: Mitotic defects in brain: cytokinesis defect (polyploidy); Male semi-sterile, Meiotic defects in testis: cytokinesis defects, segregation defects.(Seg-01/62); Meiotic defects in testis: cytokinesis defects, abnormal spindles.(Ab-02/12); Mitotic defects in brain: cytokinesis defect (no overcondensation of diploids, high polyploidy); Meiotic defects in testis: cytokinesis defects. Dark bands in eyes, dominant; Meiotic
 15 defects in testis: cytokinesis defects; Meiotic defects in testis:segregation defect, cytokinesis defect(Ck-09/32); Mitotic defects in brain: cytokinesis defect (no overcondensation of diploids, very high polyploidy); Mitotic defects in brain: cytokinesis defect(very high polyploidy); Mitotic defects in brain: cytokinesis defect. Meiotic defects in testis: cytokinesis defects (Mitotic higher level of condensation, polyploidy, Meiotic:
 20 Ck05/07); Mitotic defects in brain, Cytokinesis defect (no overcondensation of diploids, high polyploidy); Mitotic defects in brain: cytokinesis defect (very high polyploidy, chromosomes entangled?); Mitotic defects in brain: cytokinesis defect (very high polyploidy; Meiotic defects in testis: cytokinesis defects (Ck-04/06) `; Female sterile (anaphase bridges, lagging chromosomes); Mitotic defects in brain: cytokinesis defect.
 25 Meiotic defects in testis: cytokinesis defects:(mitotic: high polyploidy, no diploids, higher mitotic index, meiotic: Ck-01/05); Meiotic defects in testis: cytokinesis defects; Meiotic defects in testis: cytokinesis defects(Ck-06/09); Meiotic defects in testis: segregation defects, cytokinesis defect(Ck-07/35); Meiotic defects in testis: cytokinesis defects.

Alternatively, mutations in the polypeptides and polynucleotides of the invention may be associated with a failure to enter M-phase; such polypeptides and polynucleotides are conveniently categorised as "Category 2". Phenotypes associated with Category 2 polypeptides and polynucleotides include any one or more of the following, singly or in
 5 combination: Meiotic defects in testis: no division(no meiosis); Mitotic defects in brain: no mitosis; Meiotic defects in testis: segregation defects, meiotic failure(Mf-07/75); Meiotic defects in testis: segregation defects, meiotic failure(Mf-05/31); Meiotic defects in testis: cytokinesis defects, meiotic failure(Mf-02/15).

Mutations in the polypeptides and polynucleotides of the invention may be
 10 associated with a metaphase arrest phenotype ("Category 3"). Phenotypes associated with Category 3 polypeptides and polynucleotides include any one or more of the following, singly or in combination: Mitotic defects in brain: prometaphase arrest (overcondensation, polyploidy, scattered chromosomes with bipolar spindle); Male sterile, Female sterile, Mitotic defects in brain: prometaphase arrest (Overcondensation, polyploidy, fewer
 15 anaphases, high mitotic index, scattered chromosomes with bipolar spindle); Mitotic defects in brain: (weak overcondensation, metaphase with bipolar spindle); Mitotic defects in brain: prometaphase arrest; Mitotic defects in brain: metaphase arrest; Mitotic defects in brain: metaphase arrest. (overcondensation, polyploidy, aneuploidy, few anaphases, high mitotic index, metaphase with bent bipolar spindle); Mitotic defects in brain: metaphase
 20 arrest.(overcondensation, polyploidy, few anaphases, high mitotic index, metaphase with bent bipolar spindle); Mitotic defects in brain: Metaphase arrest (overcondensation, polyploidy, aneuploidy, no anaphases, high mitotic index, metaphase with bipolar spindle); Mitotic defects in brain: metaphase arrest (overcondensation, metaphase with bipolar spindle; Meiotic defects in testis: segregation defects, multipolar spindles (Mul-
 25 02/29); Meiotic defects in testis: cytokinesis defects, abnormal spindles.(Ab-01/03); Mitotic defects in brain: metaphase arrest; Mitotic defects in brain: metaphase arrest (overcondensation, polyploidy, metaphase with bipolar spindle); Mitotic defects in brain: metaphase arrest. Meiotic defects in testis: segregation defects. Abnormal spindles (mitotic: High mitotic index, meiotic: Ab-08/24); Mitotic defects in brain: metaphase
 30 arrest(overcondensation, few anaphases, some polyploids); Mitotic defects in brain: prometaphase arrest (overcondensation, fewer anaphases, metaphase with bipolar spindle);

Mitotic defects in brain: metaphase arrest(condensation, no polyploidy, no anaphases, metaphase with bipolar spindle).

Mutations in Category 4 polypeptides and polynucleotides of the invention may be associated with an anaphase defect phenotype; phenotypes associated with Category 4

5 polypeptides and polynucleotides include any one or more of the following, singly or in combination: Mitotic defects in brain: anaphase defects (overcondensation, high polyploidy, some lagging chromosomes); Meiotic defects in testis: segregation defects; Male and female sterile, small wings, meiotic defects in testis: segregation defects, elongation defect; Mitotic defects in brain: anaphase defects(overcondensation, anaphase

10 bridge, metaphase with swollen chromosomes and bipolar spindle); Mitotic defects in brain: Anaphase defects. (overcondensation, aneuploidy, some lagging chromosomes and breaks); Meiotic defects in testis: segregation defects; Meiotic defects in testis: segregation defects, multi-stage defects (PI-02/17); Meiotic defects in testis: segregation defects, multi-stage defects (PI-02/18); Meiotic defects in testis: cytokinesis defects,

15 segregation defects (seg-01/01); Mitotic defects in brain: cytokinesis defect. Meiotic defects in testis: cytokinesis defect. Multi-stage defects Polyploidy, no overcondensation PI-01/10; Meiotic defects in testis: segregation defects, abnormal spindles. (Ab-03/30); Mitotic defects in brain: anaphase defects (weak, higher condensation, some polyploidy, fewer anaphases, polyploids with monopolar spindles); Mitotic defects in brain: anaphase

20 defects (overcondensation, polyploidy (with overcondensation), few anaphases, metaphase with bipolar spindle); Meiotic defects in testis: cytokinesis defects; Meiotic defects in testis: segregation defects,multipolar spindles(Mul-02/22); Meiotic defects in testis: segregation defects, abnormal spindles (Ab-04/26); Meiotic defects in testis: cytokinesis defects,abnormal spindles (Ab-16/13); Mitotic defects in brain: anaphase defects. Meiotic

25 defects in testis: segregation defects, abnormal spindles (mitotic : Overcondensation, lagging chromosomes/less aligned metaphase with bipolar spindles, Meiotic: Ab-06/20); Meiotic defects in testis: segregation defects; Meiotic defects in testis: no division (no meiosis); Meiotic defects in testis: segregation defects, abnormal spindles (Ab-12/48); Meiotic defects in testis: segregation defects, multipolar spindles(mitotic: High polyploids,

30 no diploids, higher mitotic index Meiotic: Mul-02/59); Meiotic defects in testis: segregation defect; Meiotic defects in testis: segregation defects,abnormal spindles

(meiotic: Ab-08/42); Female sterile. Meiotic defects in testis: cytokinesis defects, segregation defects (Mitotic: Less condensed chromosomes, nuclear bridges, Meiotic: Seg-01/02; Mitotic defects in brain: anaphase defects; Meiotic defects in testis: cytokinesis defects, abnormal spindles (Ab-01/04); Meiotic defects in testis: segregation defects (overcondensation, fewer anaphases); Mitotic defects in brain: (some overcondensation, anaphase bridge, metaphase with swollen chromosome and bipolar spindle).

A fifth category ("Category 5") of polypeptides and polynucleotides of the invention are associated with the presence of small imaginal discs (block to proliferation). Phenotypes associated with Category 5 polypeptides and polynucleotides include any one or more of the following, singly or in combination: 2nd chromosome, small imaginal discs.

The polypeptides and polynucleotides of the invention may also be categorised according to their function, or their putative function.

For example, the polypeptides described here preferably comprise, and the polynucleotides described here are ones which preferably encode polypeptides comprising, any one or more of the following: a CBP activator protein; a CCR4-associated regulator of polymerase II transcription; a CTP synthase (CTPS); a Cyclin specific ubiquitin conjugating enzyme; a DNA packaging protein; a DNA repair protein; a DNA-binding protein involved in chromosomal organisation; a DNase IV; a EIF4G2 translation initiation factor; a eukaryotic translation initiation factor 6; a Ecdysone-induced protein 78C; a Egf2 translation factor; a G protein-coupled receptor kinase 7; a GTPase exchange factor; a phosphatidylinositol transfer protein beta isoform; a His-rich protein; a Lk6 kinase; a MAP kinase; a MAP kinase interacting kinase 1; a N-arginine dibasic convertase; a Phosphatidylinositol transfer protein; a RIP protein kinase; a RNA binding motif, single stranded interacting protein; a RNA binding protein; a RYK receptor tyrosine kinase; a Ribosomal protein L1; a selenide, water dikinase 1; a selenium donor protein 1; a selenophosphate synthetase 1; a Sqv-7-like protein; a sugar modification protein; a protein involved in cytokinesis and signalling; a TEK tyrosine kinase; a Translation elongation

factor; a UDP-galactose transporter; a v-erba related protein; a WD40 protein; a brahma protein; a calcium binding protein; a cell adhesion protein; a chaperone; a chromodomain helicase DNA binding protein; a chromodomain-helicase-DNA-binding protein; a coiled coil protein with ubiquitin like domain; a component of the 19S proteasome regulatory
5 particle; a couch potato RNA binding protein; a cytidine 5-prime triphosphate synthetase; a cytoskeletal structural protein; a death domain containing protein; a developmentally expressed in axons of the CNS; a diacylglycerol-activated/phospholipid dependent protein kinase C inhibitor; a diazepam binding inhibitor; a diphosphate kinase; a dodecasatellite DNA binding protein; a doughnut protein tyrosine kinase; an elongation factor 2; a
10 endoplasmic reticulum ATPase; a eukaryotic translation initiation factor 4E binding protein 2; a factor involved in axon guidance; a fatty-acid-Coenzyme A ligase; a flap structure-specific endonuclease 1; a protein involved in the formation of the contractile ring and the initiation of cytokinesis; a glucose-6-phosphate transporter; a glycoprotein glucosyltransferase; a growth factor; a transmembrane receptor protein tyrosine kinase
15 involved in cell growth and maintenance; a guanyl-nucleotide exchange factor involved in signal transduction; a heat shock protein; a helicase; a high density lipoprotein binding protein; a histone acetyl transferase transcriptional activator; a histone acetyltransferase; a histone acetyltransferase GCN5; a protein involved in development of the abdomen (embryos); a protein involved in the development of the imaginal discs (larvae or pupae);
20 a kinesin like protein 67a; a ligand-dependent nuclear receptor; a ligand-dependent nuclear receptor; a lola-like specific RNA polymerase II transcription factor; a matrix associated protein; a membrane glycoprotein; a mitotic heterochromatin fragment clone CH(2)6; a motor protein; a motor protein involved in cytoskeleton organization; a mushroom body RNA binding protein; a myosin like proteins; a nemo-like kinase; a non-ATPase protein; a
25 nuclear receptor NR1E1; a nucleic acid binding protein; a nucleoside diphosphate kinase (NBR-A); a oly(rC)-binding protein 2 (hnRNP-E1); a peroxisome biogenesis factor 1; a phospholipid transporter involved in lipid metabolism; a phosphatase or enhancer of Pi uptake protein; a protease; a proteasome regulatory particle; a protein involved in cytoskeleton organization and/or biogenesis; a protein kinase associated with
30 microtubules; a protein kinase mitogen-activated 7; a protein serine/threonine kinase involved in cell cycle, possibly targeted to cytoskeleton; a protein serine/threonine kinase involved in eye morphogenesis; a protein which associates with cdc25 phosphatase; a

protein which induces apoptosis; a ribonuclease P; a ribonuclease P protein subunit p29; a ser/thr phosphatase; a signal transduction protein; a signal transport protein; a sin3-associated polypeptide; a single stranded DNA/RNA binding protein; a sodium-dependent dicarboxylate transporters; a ssDNA/RNA binding proteins; a striatin, calmodulin-binding protein (STRN); a structural protein of ribosome involved in protein biosynthesis; a subtelomeric heterochromatin repeats; a sugar acetylase; a sugar modification protein; a suppressor of ras; a tRNA processing enzyme Ribonuclease P protein subunit; a thyroid hormone responsive gene; a tie receptor protein tyrosine kinase; a transacylase; a transcription factor; a transcription factor involved in chromatin remodelling; a transcriptional regulation of c-myc expression; a transcriptional regulator; a transcriptional regulators/telomeric silencing; a translation initiation factor; a tumor metastasis inhibitor; a tyrosine 3-monooxygenase/tryptophan 5-monooxygenase activation protein; a ubiquitin carrier protein; a ubiquitin-conjugating enzyme; a ugtUDP-glucose-glycoprotein glucosyltransferase; a zinc finger protein; an RNA polymerase II transcription factor; an acetylcholinesterase (YT blood group) precursor; an actin binding protein; an actin dependent regulator of chromatin; an acyl-CoA-binding protein; an alanine:glyoxylate aminotransferase; an alpha esterase; an ankyrin protein; an imitation-SWI protein; and an integrin beta 4 binding protein.

POLYPEPTIDES

- 20 It will be understood that polypeptides of the invention are not limited to polypeptides having the amino acid sequence set out in Examples 1 to 70 or fragments thereof but also include homologous sequences obtained from any source, for example related viral/bacterial proteins, cellular homologues and synthetic peptides, as well as variants or derivatives thereof.
- 25 Thus polypeptides of the invention also include those encoding homologues from other species including animals such as mammals (e.g. mice, rats or rabbits), especially primates, more especially humans. More specifically, homologues included within the scope of the invention include human homologues.

Thus, the present invention covers variants, homologues or derivatives of the amino acid sequence set out in Examples 1 to 70, as well as variants, homologues or derivatives of the nucleotide sequence coding for the amino acid sequences of the present invention.

5 In the context of the present invention, a homologous sequence is taken to include an amino acid sequence which is at least 15, 20, 25, 30, 40, 50, 60, 70, 80 or 90% identical, preferably at least 95 or 98% identical at the amino acid level over at least 50 or 100, preferably 200, 300, 400 or 500 amino acids with any one of the polypeptide sequences shown in the Examples. In particular, homology should typically be considered
10 with respect to those regions of the sequence known to be essential for protein function rather than non-essential neighbouring sequences. This is especially important when considering homologous sequences from distantly related organisms.

 Although homology can also be considered in terms of similarity (i.e. amino acid residues having similar chemical properties/functions), in the context of the present
15 invention it is preferred to express homology in terms of sequence identity.

 Homology comparisons can be conducted by eye, or more usually, with the aid of readily available sequence comparison programs. These publicly and commercially available computer programs can calculate % homology between two or more sequences.

 % homology may be calculated over contiguous sequences, i.e. one sequence is
20 aligned with the other sequence and each amino acid in one sequence directly compared with the corresponding amino acid in the other sequence, one residue at a time. This is called an "ungapped" alignment. Typically, such ungapped alignments are performed only over a relatively short number of residues (for example less than 50 contiguous amino acids).

25 Although this is a very simple and consistent method, it fails to take into consideration that, for example, in an otherwise identical pair of sequences, one insertion or deletion will cause the following amino acid residues to be put out of alignment, thus

potentially resulting in a large reduction in % homology when a global alignment is performed. Consequently, most sequence comparison methods are designed to produce optimal alignments that take into consideration possible insertions and deletions without penalising unduly the overall homology score. This is achieved by inserting “gaps” in the sequence alignment to try to maximise local homology.

However, these more complex methods assign “gap penalties” to each gap that occurs in the alignment so that, for the same number of identical amino acids, a sequence alignment with as few gaps as possible - reflecting higher relatedness between the two compared sequences - will achieve a higher score than one with many gaps. “Affine gap costs” are typically used that charge a relatively high cost for the existence of a gap and a smaller penalty for each subsequent residue in the gap. This is the most commonly used gap scoring system. High gap penalties will of course produce optimised alignments with fewer gaps. Most alignment programs allow the gap penalties to be modified. However, it is preferred to use the default values when using such software for sequence comparisons. For example when using the GCG Wisconsin Bestfit package (see below) the default gap penalty for amino acid sequences is -12 for a gap and -4 for each extension.

Calculation of maximum % homology therefore firstly requires the production of an optimal alignment, taking into consideration gap penalties. A suitable computer program for carrying out such an alignment is the GCG Wisconsin Bestfit package (University of Wisconsin, U.S.A; Devereux *et al.*, 1984, Nucleic Acids Research 12:387). Examples of other software than can perform sequence comparisons include, but are not limited to, the BLAST package (see Ausubel *et al.*, 1999 *ibid* – Chapter 18), FASTA (Atschul *et al.*, 1990, J. Mol. Biol., 403-410) and the GENWORKS suite of comparison tools. Both BLAST and FASTA are available for offline and online searching (see Ausubel *et al.*, 1999 *ibid*, pages 7-58 to 7-60). However it is preferred to use the GCG Bestfit program.

Although the final % homology can be measured in terms of identity, the alignment process itself is typically not based on an all-or-nothing pair comparison. Instead, a scaled similarity score matrix is generally used that assigns scores to each

pairwise comparison based on chemical similarity or evolutionary distance. An example of such a matrix commonly used is the BLOSUM62 matrix - the default matrix for the BLAST suite of programs. GCG Wisconsin programs generally use either the public default values or a custom symbol comparison table if supplied (see user manual for
5 further details). It is preferred to use the public default values for the GCG package, or in the case of other software, the default matrix, such as BLOSUM62.

Once the software has produced an optimal alignment, it is possible to calculate % homology, preferably % sequence identity. The software typically does this as part of the sequence comparison and generates a numerical result.

10 The terms "variant" or "derivative" in relation to the amino acid sequences of the present invention includes any substitution of, variation of, modification of, replacement of, deletion of or addition of one (or more) amino acids from or to the sequence providing the resultant amino acid sequence retains substantially the same activity as the unmodified sequence, preferably having at least the same activity as the polypeptides presented in the
15 sequence listings in the Examples.

Polypeptides having the amino acid sequence shown in the Examples, or fragments or homologues thereof may be modified for use in the present invention. Typically, modifications are made that maintain the biological activity of the sequence. Amino acid substitutions may be made, for example from 1, 2 or 3 to 10, 20 or 30 substitutions
20 provided that the modified sequence retains the biological activity of the unmodified sequence. Alternatively, modifications may be made to deliberately inactivate one or more functional domains of the polypeptides of the invention. Amino acid substitutions may include the use of non-naturally occurring analogues, for example to increase blood plasma half-life of a therapeutically administered polypeptide.

25 Conservative substitutions may be made, for example according to the Table below. Amino acids in the same block in the second column and preferably in the same line in the third column may be substituted for each other:

ALIPHATIC	Non-polar	G A P
		I L V
	Polar - uncharged	C S T M
		N Q
	Polar - charged	D E
		K R
AROMATIC		H F W Y

Polypeptides of the invention also include fragments of the full length sequences mentioned above. Preferably said fragments comprise at least one epitope. Methods of identifying epitopes are well known in the art. Fragments will typically comprise at least 6 amino acids, more preferably at least 10, 20, 30, 50 or 100 amino acids.

5 Proteins of the invention are typically made by recombinant means, for example as described below. However they may also be made by synthetic means using techniques well known to skilled persons such as solid phase synthesis. Proteins of the invention may also be produced as fusion proteins, for example to aid in extraction and purification. Examples of fusion protein partners include glutathione-S-transferase (GST), 6xHis,
 10 GAL4 (DNA binding and/or transcriptional activation domains) and β -galactosidase. It may also be convenient to include a proteolytic cleavage site between the fusion protein partner and the protein sequence of interest to allow removal of fusion protein sequences. Preferably the fusion protein will not hinder the function of the protein of interest sequence. Proteins of the invention may also be obtained by purification of cell extracts
 15 from animal cells.

Proteins of the invention may be in a substantially isolated form. It will be understood that the protein may be mixed with carriers or diluents which will not interfere with the intended purpose of the protein and still be regarded as substantially isolated. A protein of the invention may also be in a substantially purified form, in which case it will
 20 generally comprise the protein in a preparation in which more than 90%, e.g. 95%, 98% or 99% of the protein in the preparation is a protein of the invention.

A polypeptide of the invention may be labeled with a revealing label. The revealing label may be any suitable label which allows the polypeptide to be detected. Suitable labels include radioisotopes, e.g. ¹²⁵I, enzymes, antibodies, polynucleotides and linkers such as biotin. Labeled polypeptides of the invention may be used in diagnostic procedures such as

5 immunoassays to determine the amount of a polypeptide of the invention in a sample. Polypeptides or labeled polypeptides of the invention may also be used in serological or cell-mediated immune assays for the detection of immune reactivity to said polypeptides in animals and humans using standard protocols.

A polypeptide or labeled polypeptide of the invention or fragment thereof may also

10 be fixed to a solid phase, for example the surface of an immunoassay well or dipstick. Such labeled and/or immobilised polypeptides may be packaged into kits in a suitable container along with suitable reagents, controls, instructions and the like. Such polypeptides and kits may be used in methods of detection of antibodies to the polypeptides or their allelic or species variants by immunoassay.

15 Immunoassay methods are well known in the art and will generally comprise: (a) providing a polypeptide comprising an epitope bindable by an antibody against said protein; (b) incubating a biological sample with said polypeptide under conditions which allow for the formation of an antibody-antigen complex; and (c) determining whether antibody-antigen complex comprising said polypeptide is formed.

20 Polypeptides of the invention may be used in *in vitro* or *in vivo* cell culture systems to study the role of their corresponding genes and homologues thereof in cell function, including their function in disease. For example, truncated or modified polypeptides may be introduced into a cell to disrupt the normal functions which occur in the cell. The polypeptides of the invention may be introduced into the cell by *in situ* expression of the

25 polypeptide from a recombinant expression vector (see below). The expression vector optionally carries an inducible promoter to control the expression of the polypeptide.

The use of appropriate host cells, such as insect cells or mammalian cells, is expected to provide for such post-translational modifications (e.g. myristolation,

glycosylation, truncation, lapidation and tyrosine, serine or threonine phosphorylation) as may be needed to confer optimal biological activity on recombinant expression products of the invention. Such cell culture systems in which polypeptides of the invention are expressed may be used in assay systems to identify candidate substances which interfere
5 with or enhance the functions of the polypeptides of the invention in the cell.

POLYNUCLEOTIDES

Polynucleotides of the invention include polynucleotides that comprise any one or more of the nucleic acid sequences set out in Examples 1 to 70 and fragments thereof. Polynucleotides of the invention also include polynucleotides encoding the polypeptides
10 of the invention. It will be understood by a skilled person that numerous different polynucleotides can encode the same polypeptide as a result of the degeneracy of the genetic code. In addition, it is to be understood that skilled persons may, using routine techniques, make nucleotide substitutions that do not affect the polypeptide sequence encoded by the polynucleotides of the invention to reflect the codon usage of any
15 particular host organism in which the polypeptides of the invention are to be expressed.

Polynucleotides of the invention may comprise DNA or RNA. They may be single-stranded or double-stranded. They may also be polynucleotides which include within them synthetic or modified nucleotides. A number of different types of modification to oligonucleotides are known in the art. These include methylphosphonate
20 and phosphorothioate backbones, addition of acridine or polylysine chains at the 3' and/or 5' ends of the molecule. For the purposes of the present invention, it is to be understood that the polynucleotides described herein may be modified by any method available in the art. Such modifications may be carried out in order to enhance the *in vivo* activity or life span of polynucleotides of the invention.

25 The terms "variant", "homologue" or "derivative" in relation to the nucleotide sequence of the present invention include any substitution of, variation of, modification of, replacement of, deletion of or addition of one (or more) nucleic acid from or to the

sequence. Preferably said variant, homologues or derivatives code for a polypeptide having biological activity.

As indicated above, with respect to sequence homology, preferably there is at least 50 or 75%, more preferably at least 85%, more preferably at least 90% homology to the sequences shown in the sequence listing herein. More preferably there is at least 95%, more preferably at least 98%, homology. Nucleotide homology comparisons may be conducted as described above. A preferred sequence comparison program is the GCG Wisconsin Bestfit program described above. The default scoring matrix has a match value of 10 for each identical nucleotide and -9 for each mismatch. The default gap creation penalty is -50 and the default gap extension penalty is -3 for each nucleotide.

The present invention also encompasses nucleotide sequences that are capable of hybridising selectively to the sequences presented herein, or any variant, fragment or derivative thereof, or to the complement of any of the above. Nucleotide sequences are preferably at least 15 nucleotides in length, more preferably at least 20, 30, 40 or 50 nucleotides in length.

The term "hybridization" as used herein shall include "the process by which a strand of nucleic acid joins with a complementary strand through base pairing" as well as the process of amplification as carried out in polymerase chain reaction technologies.

Polynucleotides of the invention capable of selectively hybridising to the nucleotide sequences presented herein, or to their complement, will be generally at least 70%, preferably at least 80 or 90% and more preferably at least 95% or 98% homologous to the corresponding nucleotide sequences presented herein over a region of at least 20, preferably at least 25 or 30, for instance at least 40, 60 or 100 or more contiguous nucleotides.

The term "selectively hybridizable" means that the polynucleotide used as a probe is used under conditions where a target polynucleotide of the invention is found to hybridize to the probe at a level significantly above background. The background

hybridization may occur because of other polynucleotides present, for example, in the cDNA or genomic DNA library being screening. In this event, background implies a level of signal generated by interaction between the probe and a non-specific DNA member of the library which is less than 10 fold, preferably less than 100 fold as intense as the
5 specific interaction observed with the target DNA. The intensity of interaction may be measured, for example, by radiolabelling the probe, e.g. with ^{32}P .

Hybridization conditions are based on the melting temperature (T_m) of the nucleic acid binding complex, as taught in Berger and Kimmel (1987, Guide to Molecular Cloning Techniques, Methods in Enzymology, Vol 152, Academic Press, San Diego CA), and
10 confer a defined "stringency" as explained below.

Maximum stringency typically occurs at about $T_m - 5^\circ\text{C}$ (5°C below the T_m of the probe); high stringency at about 5°C to 10°C below T_m ; intermediate stringency at about 10°C to 20°C below T_m ; and low stringency at about 20°C to 25°C below T_m . As will be understood by those of skill in the art, a maximum stringency hybridization can be used to
15 identify or detect identical polynucleotide sequences while an intermediate (or low) stringency hybridization can be used to identify or detect similar or related polynucleotide sequences.

In a preferred aspect, the present invention covers nucleotide sequences that can hybridise to the nucleotide sequence of the present invention under stringent conditions
20 (e.g. 65°C and $0.1\times\text{SSC}$ { $1\times\text{SSC} = 0.15\text{ M NaCl}$, $0.015\text{ M Na}_3\text{ Citrate pH } 7.0$ }).

Where the polynucleotide of the invention is double-stranded, both strands of the duplex, either individually or in combination, are encompassed by the present invention. Where the polynucleotide is single-stranded, it is to be understood that the complementary sequence of that polynucleotide is also included within the scope of the present invention.

25 Polynucleotides which are not 100% homologous to the sequences of the present invention but fall within the scope of the invention can be obtained in a number of ways. Other variants of the sequences described herein may be obtained for example by probing

DNA libraries made from a range of individuals, for example individuals from different populations. In addition, other viral/bacterial, or cellular homologues particularly cellular homologues found in mammalian cells (e.g. rat, mouse, bovine and primate cells), may be obtained and such homologues and fragments thereof in general will be capable of

5 selectively hybridising to the sequences shown in the Examples. Such sequences may be obtained by probing cDNA libraries made from or genomic DNA libraries from other animal species, and probing such libraries with probes comprising all or part of any one of the sequences shown in the Examples under conditions of medium to high stringency. The nucleotide sequences of the human homologues described in the Examples, may

10 preferably be used to identify other primate/mammalian homologues since nucleotide homology between human sequences and mammalian sequences is likely to be higher than is the case for the *Drosophila* sequences identified herein.

Similar considerations apply to obtaining species homologues and allelic variants of the polypeptide or nucleotide sequences of the invention.

15 Variants and strain/species homologues may also be obtained using degenerate PCR which will use primers designed to target sequences within the variants and homologues encoding conserved amino acid sequences within the sequences of the present invention. Conserved sequences can be predicted, for example, by aligning the amino acid sequences from several variants/homologues. Sequence alignments can be performed

20 using computer software known in the art. For example the GCG Wisconsin PileUp program is widely used.

The primers used in degenerate PCR will contain one or more degenerate positions and will be used at stringency conditions lower than those used for cloning sequences with single sequence primers against known sequences. It will be appreciated by the skilled

25 person that overall nucleotide homology between sequences from distantly related organisms is likely to be very low and thus in these situations degenerate PCR may be the method of choice rather than screening libraries with labeled fragments the sequences disclosed in the Examples.

In addition, homologous sequences may be identified by searching nucleotide and/or protein databases using search algorithms such as the BLAST suite of programs. This approach is described in the Examples.

Alternatively, such polynucleotides may be obtained by site directed mutagenesis of characterised sequences, such as the sequences disclosed in the Examples. This may be useful where for example silent codon changes are required to sequences to optimise codon preferences for a particular host cell in which the polynucleotide sequences are being expressed. Other sequence changes may be desired in order to introduce restriction enzyme recognition sites, or to alter the property or function of the polypeptides encoded by the polynucleotides. For example, further changes may be desirable to represent particular coding changes found in the sequences disclosed in the Examples which give rise to mutant genes which have lost their regulatory function. Probes based on such changes can be used as diagnostic probes to detect such mutants.

Polynucleotides of the invention may be used to produce a primer, e.g. a PCR primer, a primer for an alternative amplification reaction, a probe e.g. labeled with a revealing label by conventional means using radioactive or non-radioactive labels, or the polynucleotides may be cloned into vectors. Such primers, probes and other fragments will be at least 8, 9, 10, or 15, preferably at least 20, for example at least 25, 30 or 40 nucleotides in length, and are also encompassed by the term polynucleotides of the invention as used herein.

Polynucleotides such as a DNA polynucleotides and probes according to the invention may be produced recombinantly, synthetically, or by any means available to those of skill in the art. They may also be cloned by standard techniques.

In general, primers will be produced by synthetic means, involving a step wise manufacture of the desired nucleic acid sequence one nucleotide at a time. Techniques for accomplishing this using automated techniques are readily available in the art.

Longer polynucleotides will generally be produced using recombinant means, for example using a PCR (polymerase chain reaction) cloning techniques. This will involve making a pair of primers (e.g. of about 15 to 30 nucleotides) flanking a region of the lipid targeting sequence which it is desired to clone, bringing the primers into contact with mRNA or cDNA obtained from an animal or human cell, performing a polymerase chain reaction under conditions which bring about amplification of the desired region, isolating the amplified fragment (e.g. by purifying the reaction mixture on an agarose gel) and recovering the amplified DNA. The primers may be designed to contain suitable restriction enzyme recognition sites so that the amplified DNA can be cloned into a suitable cloning vector

Polynucleotides or primers of the invention may carry a revealing label. Suitable labels include radioisotopes such as ^{32}P or ^{35}S , enzyme labels, or other protein labels such as biotin. Such labels may be added to polynucleotides or primers of the invention and may be detected using by techniques known *per se*.

Polynucleotides or primers of the invention or fragments thereof labeled or unlabeled may be used by a person skilled in the art in nucleic acid-based tests for detecting or sequencing polynucleotides of the invention in the human or animal body.

Such tests for detecting generally comprise bringing a biological sample containing DNA or RNA into contact with a probe comprising a polynucleotide or primer of the invention under hybridising conditions and detecting any duplex formed between the probe and nucleic acid in the sample. Such detection may be achieved using techniques such as PCR or by immobilising the probe on a solid support, removing nucleic acid in the sample which is not hybridised to the probe, and then detecting nucleic acid which has hybridised to the probe. Alternatively, the sample nucleic acid may be immobilised on a solid support, and the amount of probe bound to such a support can be detected. Suitable assay methods of this and other formats can be found in for example WO89/03891 and WO90/13667.

Tests for sequencing nucleotides of the invention include bringing a biological sample containing target DNA or RNA into contact with a probe comprising a polynucleotide or primer of the invention under hybridising conditions and determining the sequence by, for example the Sanger dideoxy chain termination method (see
5 Sambrook *et al.*).

Such a method generally comprises elongating, in the presence of suitable reagents, the primer by synthesis of a strand complementary to the target DNA or RNA and selectively terminating the elongation reaction at one or more of an A, C, G or T/U residue; allowing strand elongation and termination reaction to occur; separating out
10 according to size the elongated products to determine the sequence of the nucleotides at which selective termination has occurred. Suitable reagents include a DNA polymerase enzyme, the deoxynucleotides dATP, dCTP, dGTP and dTTP, a buffer and ATP. Dideoxynucleotides are used for selective termination.

Tests for detecting or sequencing nucleotides of the invention in a biological
15 sample may be used to determine particular sequences within cells in individuals who have, or are suspected to have, an altered gene sequence, for example within cancer cells including leukaemia cells and solid tumours such as breast, ovary, lung, colon, pancreas, testes, liver, brain, muscle and bone tumours. Cells from patients suffering from a proliferative disease may also be tested in the same way.

20 In addition, the identification of the genes described in the Examples will allow the role of these genes in hereditary diseases to be investigated. In general, this will involve establishing the status of the gene (e.g. using PCR sequence analysis), in cells derived from animals or humans with, for example, neurological disorders or neoplasms.

The probes of the invention may conveniently be packaged in the form of a test kit
25 in a suitable container. In such kits the probe may be bound to a solid support where the assay format for which the kit is designed requires such binding. The kit may also contain suitable reagents for treating the sample to be probed, hybridising the probe to nucleic acid in the sample, control reagents, instructions, and the like.

NUCLEIC ACID VECTORS

Polynucleotides of the invention can be incorporated into a recombinant replicable vector. The vector may be used to replicate the nucleic acid in a compatible host cell. Thus in a further embodiment, the invention provides a method of making polynucleotides of the invention by introducing a polynucleotide of the invention into a replicable vector, introducing the vector into a compatible host cell, and growing the host cell under conditions which bring about replication of the vector. The vector may be recovered from the host cell. Suitable host cells include bacteria such as *E. coli*, yeast, mammalian cell lines and other eukaryotic cell lines, for example insect Sf9 cells.

Preferably, a polynucleotide of the invention in a vector is operably linked to a control sequence that is capable of providing for the expression of the coding sequence by the host cell, i.e. the vector is an expression vector. The term "operably linked" means that the components described are in a relationship permitting them to function in their intended manner. A regulatory sequence "operably linked" to a coding sequence is ligated in such a way that expression of the coding sequence is achieved under condition compatible with the control sequences.

The control sequences may be modified, for example by the addition of further transcriptional regulatory elements to make the level of transcription directed by the control sequences more responsive to transcriptional modulators.

Vectors of the invention may be transformed or transfected into a suitable host cell as described below to provide for expression of a protein of the invention. This process may comprise culturing a host cell transformed with an expression vector as described above under conditions to provide for expression by the vector of a coding sequence encoding the protein, and optionally recovering the expressed protein. Vectors will be chosen that are compatible with the host cell used.

The vectors may be for example, plasmid or virus vectors provided with an origin of replication, optionally a promoter for the expression of the said polynucleotide and

optionally a regulator of the promoter. The vectors may contain one or more selectable marker genes, for example an ampicillin resistance gene in the case of a bacterial plasmid or a neomycin resistance gene for a mammalian vector. Vectors may be used, for example, to transfect or transform a host cell.

- 5 Control sequences operably linked to sequences encoding the polypeptide of the invention include promoters/enhancers and other expression regulation signals. These control sequences may be selected to be compatible with the host cell for which the expression vector is designed to be used in. The term promoter is well-known in the art and encompasses nucleic acid regions ranging in size and complexity from minimal
10 promoters to promoters including upstream elements and enhancers.

- The promoter is typically selected from promoters which are functional in mammalian cells, although prokaryotic promoters and promoters functional in other eukaryotic cells, such as insect cells, may be used. The promoter is typically derived from promoter sequences of viral or eukaryotic genes. For example, it may be a promoter
15 derived from the genome of a cell in which expression is to occur. With respect to eukaryotic promoters, they may be promoters that function in a ubiquitous manner (such as promoters of α -actin, β -actin, tubulin) or, alternatively, a tissue-specific manner (such as promoters of the genes for pyruvate kinase). They may also be promoters that respond to specific stimuli, for example promoters that bind steroid hormone receptors. Viral
20 promoters may also be used, for example the Moloney murine leukaemia virus long terminal repeat (MMLV LTR) promoter, the rous sarcoma virus (RSV) LTR promoter or the human cytomegalovirus (CMV) IE promoter.

- It may also be advantageous for the promoters to be inducible so that the levels of expression of the heterologous gene can be regulated during the life-time of the cell.
25 Inducible means that the levels of expression obtained using the promoter can be regulated.

In addition, any of these promoters may be modified by the addition of further regulatory sequences, for example enhancer sequences. Chimeric promoters may also be

used comprising sequence elements from two or more different promoters described above.

Polynucleotides according to the invention may also be inserted into the vectors described above in an antisense orientation to provide for the production of antisense
5 RNA. Antisense RNA or other antisense polynucleotides may also be produced by synthetic means. Such antisense polynucleotides may be used in a method of controlling the levels of RNAs transcribed from genes comprising any one of the polynucleotides of the invention.

HOST CELLS

10 Vectors and polynucleotides of the invention may be introduced into host cells for the purpose of replicating the vectors/polynucleotides and/or expressing the polypeptides of the invention encoded by the polynucleotides of the invention. Although the polypeptides of the invention may be produced using prokaryotic cells as host cells, it is preferred to use eukaryotic cells, for example yeast, insect or mammalian cells, in
15 particular mammalian cells.

Vectors/polynucleotides of the invention may be introduced into suitable host cells using a variety of techniques known in the art, such as transfection, transformation and electroporation. Where vectors/polynucleotides of the invention are to be administered to animals, several techniques are known in the art, for example infection with recombinant
20 viral vectors such as retroviruses, herpes simplex viruses and adenoviruses, direct injection of nucleic acids and biolistic transformation.

PROTEIN EXPRESSION AND PURIFICATION

Host cells comprising polynucleotides of the invention may be used to express polypeptides of the invention. Host cells may be cultured under suitable conditions which
25 allow expression of the proteins of the invention. Expression of the polypeptides of the invention may be constitutive such that they are continually produced, or inducible, requiring a stimulus to initiate expression. In the case of inducible expression, protein

production can be initiated when required by, for example, addition of an inducer substance to the culture medium, for example dexamethasone or IPTG.

Polypeptides of the invention can be extracted from host cells by a variety of techniques known in the art, including enzymatic, chemical and/or osmotic lysis and
5 physical disruption.

Polypeptides of the invention may also be produced recombinantly in an *in vitro* cell-free system, such as the TnTTM (Promega) rabbit reticulocyte system.

ANTIBODIES

The invention also provides monoclonal or polyclonal antibodies to polypeptides of
10 the invention or fragments thereof. Thus, the present invention further provides a process for the production of monoclonal or polyclonal antibodies to polypeptides of the invention.

If polyclonal antibodies are desired, a selected mammal (e.g., mouse, rabbit, goat, horse, etc.) is immunised with an immunogenic polypeptide bearing an epitope(s) from a polypeptide of the invention. Serum from the immunised animal is collected and treated
15 according to known procedures. If serum containing polyclonal antibodies to an epitope from a polypeptide of the invention contains antibodies to other antigens, the polyclonal antibodies can be purified by immunoaffinity chromatography. Techniques for producing and processing polyclonal antisera are known in the art. In order that such antibodies may be made, the invention also provides polypeptides of the invention or fragments thereof
20 haptenised to another polypeptide for use as immunogens in animals or humans.

Monoclonal antibodies directed against epitopes in the polypeptides of the invention can also be readily produced by one skilled in the art. The general methodology for making monoclonal antibodies by hybridomas is well known. Immortal antibody-producing cell lines can be created by cell fusion, and also by other techniques such as
25 direct transformation of B lymphocytes with oncogenic DNA, or transfection with Epstein-Barr virus. Panels of monoclonal antibodies produced against epitopes in the

polypeptides of the invention can be screened for various properties; i.e., for isotype and epitope affinity.

An alternative technique involves screening phage display libraries where, for example the phage express scFv fragments on the surface of their coat with a large variety
5 of complementarity determining regions (CDRs). This technique is well known in the art.

Antibodies, both monoclonal and polyclonal, which are directed against epitopes from polypeptides of the invention are particularly useful in diagnosis, and those which are neutralising are useful in passive immunotherapy. Monoclonal antibodies, in particular, may be used to raise anti-idiotypic antibodies. Anti-idiotypic antibodies are
10 immunoglobulins which carry an "internal image" of the antigen of the agent against which protection is desired.

Techniques for raising anti-idiotypic antibodies are known in the art. These anti-idiotypic antibodies may also be useful in therapy.

For the purposes of this invention, the term "antibody", unless specified to the
15 contrary, includes fragments of whole antibodies which retain their binding activity for a target antigen. Such fragments include Fv, F(ab') and F(ab')₂ fragments, as well as single chain antibodies (scFv). Furthermore, the antibodies and fragments thereof may be humanised antibodies, for example as described in EP-A-239400.

Antibodies may be used in method of detecting polypeptides of the invention
20 present in biological samples by a method which comprises: (a) providing an antibody of the invention; (b) incubating a biological sample with said antibody under conditions which allow for the formation of an antibody-antigen complex; and (c) determining whether antibody-antigen complex comprising said antibody is formed.

Suitable samples include extracts tissues such as brain, breast, ovary, lung, colon,
25 pancreas, testes, liver, muscle and bone tissues or from neoplastic growths derived from such tissues.

Antibodies of the invention may be bound to a solid support and/or packaged into kits in a suitable container along with suitable reagents, controls, instructions and the like.

ASSAYS

The present invention provides assays that are suitable for identifying substances
5 which bind to polypeptides of the invention and which affect, for example, formation of the nuclear envelope, exit from the quiescent phase of the cell cycle (G0), G1 progression, chromosome decondensation, nuclear envelope breakdown, START, initiation of DNA replication, progression of DNA replication, termination of DNA replication, centrosome duplication, G2 progression, activation of mitotic or meiotic functions, chromosome
10 condensation, centrosome separation, microtubule nucleation, spindle formation and function, interactions with microtubule motor proteins, chromatid separation and segregation, inactivation of mitotic functions, formation of contractile ring, cytokinesis functions, chromatin binding, formation of replication complexes, replication licensing, phosphorylation or other secondary modification activity, proteolytic degradation,
15 microtubule binding, actin binding, septin binding, microtubule organising centre nucleation activity and binding to components of cell cycle signalling pathways.

In addition, assays suitable for identifying substances that interfere with binding of polypeptides of the invention, where appropriate, to components of cell division cycle machinery. This includes not only components such as microtubules but also signalling
20 components and regulatory components as indicated above. Such assays are typically *in vitro*. Assays are also provided that test the effects of candidate substances identified in preliminary *in vitro* assays on intact cells in whole cell assays. The assays described below, or any suitable assay as known in the art, may be used to identify these substances.

According to one aspect of the invention, therefore, we provide one or more
25 substances identified by any of the assays described below, viz, mitosis assays, meiotic assays, polypeptide binding assays, microtubule binding/polymerisation assays, microtubule purification and binding assays, microtubule organising centre (MTOC) nucleation activity assays, motor protein assay, assay for spindle assembly and function,

assays for dna replication, chromosome condensation assays, kinase assays, kinase inhibitor assays, and whole cell assays, each as described in further detail below.

CANDIDATE SUBSTANCES

A substance that inhibits cell cycle progression as a result of an interaction with a polypeptide of the invention may do so in several ways. For example, if the substance inhibits cell division, mitosis and/or meiosis, it may directly disrupt the binding of a polypeptide of the invention to a component of the spindle apparatus by, for example, binding to the polypeptide and masking or altering the site of interaction with the other component. A substance which inhibits DNA replication may do so by inhibiting the phosphorylation or de-phosphorylation of proteins involved in replication. For example, it is known that the kinase inhibitor 6-DMAP (6-dimethylaminopurine) prevents the initiation of replication (Blow, JJ, 1993, *J Cell Biol*122,993-1002). Candidate substances of this type may conveniently be preliminarily screened by *in vitro* binding assays as, for example, described below and then tested, for example in a whole cell assay as described below. Examples of candidate substances include antibodies which recognise a polypeptide of the invention.

A substance which can bind directly to a polypeptide of the invention may also inhibit its function in cell cycle progression by altering its subcellular localisation and hence its ability to interact with its normal substrate. The substance may alter the subcellular localisation of the polypeptide by directly binding to it, or by indirectly disrupting the interaction of the polypeptide with another component. For example, it is known that interaction between the p68 and p180 subunits of DNA polymerase alpha-primase enzyme is necessary in order for p180 to translocate into the nucleus (Mizuno et al (1998) *Mol Cell Biol*18,3552-62), and accordingly, a substance which disrupts the interaction between p68 and p180 will affect nuclear translocation and hence activity of the primase. A substance which affects mitosis may do so by preventing the polypeptide and components of the mitotic apparatus from coming into contact within the cell.

These substances may be tested using, for example the whole cells assays described below. Non-functional homologues of a polypeptide of the invention may also be tested

for inhibition of cell cycle progression since they may compete with the wild type protein for binding to components of the cell division cycle machinery whilst being incapable of the normal functions of the protein or block the function of the protein bound to the cell division cycle machinery. Such non-functional homologues may include naturally occurring mutants and modified sequences or fragments thereof.

Alternatively, instead of preventing the association of the components directly, the substance may suppress the biologically available amount of a polypeptide of the invention. This may be by inhibiting expression of the component, for example at the level of transcription, transcript stability, translation or post-translational stability. An example of such a substance would be antisense RNA or double-stranded interfering RNA sequences which suppresses the amount of mRNA biosynthesis.

Suitable candidate substances include peptides, especially of from about 5 to 30 or 10 to 25 amino acids in size, based on the sequence of the polypeptides described in the Examples, or variants of such peptides in which one or more residues have been substituted. Peptides from panels of peptides comprising random sequences or sequences which have been varied consistently to provide a maximally diverse panel of peptides may be used.

Suitable candidate substances also include antibody products (for example, monoclonal and polyclonal antibodies, single chain antibodies, chimeric antibodies and CDR-grafted antibodies) which are specific for a polypeptide of the invention. Furthermore, combinatorial libraries, peptide and peptide mimetics, defined chemical entities, oligonucleotides, and natural product libraries may be screened for activity as inhibitors of binding of a polypeptide of the invention to the cell division cycle machinery, for example mitotic/meiotic apparatus (such as microtubules). The candidate substances may be used in an initial screen in batches of, for example 10 substances per reaction, and the substances of those batches which show inhibition tested individually. Candidate substances which show activity in *in vitro* screens such as those described below can then be tested in whole cell systems, such as mammalian cells which will be exposed to the inhibitor and tested for inhibition of any of the stages of the cell cycle.

Polypeptide Binding Assays

One type of assay for identifying substances that bind to a polypeptide of the invention involves contacting a polypeptide of the invention, which is immobilised on a solid support, with a non-immobilised candidate substance determining whether and/or to what extent the polypeptide of the invention and candidate substance bind to each other. Alternatively, the candidate substance may be immobilised and the polypeptide of the invention non-immobilised.

In a preferred assay method, the polypeptide of the invention is immobilised on beads such as agarose beads. Typically this is achieved by expressing the component as a GST-fusion protein in bacteria, yeast or higher eukaryotic cell lines and purifying the GST-fusion protein from crude cell extracts using glutathione-agarose beads (Smith and Johnson, 1988). As a control, binding of the candidate substance, which is not a GST-fusion protein, to the immobilised polypeptide of the invention is determined in the absence of the polypeptide of the invention. The binding of the candidate substance to the immobilised polypeptide of the invention is then determined. This type of assay is known in the art as a GST pulldown assay. Again, the candidate substance may be immobilised and the polypeptide of the invention non-immobilised.

It is also possible to perform this type of assay using different affinity purification systems for immobilising one of the components, for example Ni-NTA agarose and histidine-tagged components.

Binding of the polypeptide of the invention to the candidate substance may be determined by a variety of methods well-known in the art. For example, the non-immobilised component may be labeled (with for example, a radioactive label, an epitope tag or an enzyme-antibody conjugate). Alternatively, binding may be determined by immunological detection techniques. For example, the reaction mixture can be Western blotted and the blot probed with an antibody that detects the non-immobilised component. ELISA techniques may also be used.

Candidate substances are typically added to a final concentration of from 1 to 1000 nmol/ml, more preferably from 1 to 100 nmol/ml. In the case of antibodies, the final concentration used is typically from 100 to 500 µg/ml, more preferably from 200 to 300 µg/ml.

5 ***Microtubule Binding/Polymerisation Assays***

In the case of polypeptides of the invention that bind to microtubules, another type of *in vitro* assay involves determining whether a candidate substance modulates binding of a polypeptide of the invention to microtubules. Such an assay typically comprises contacting a polypeptide of the invention with microtubules in the presence or absence of
10 the candidate substance and determining if the candidate substance has an effect on the binding of the polypeptide of the invention to the microtubules. This assay can also be used in the absence of candidate substances to confirm that a polypeptide of the invention does indeed bind to microtubules. Microtubules may be prepared and assays conducted as follows:

15 ***Microtubule Purification and Binding Assays***

Microtubules are purified from 0-3h-old *Drosophila* embryos essentially as described previously (Saunders, *et al.*, 1997). About 3 ml of embryos are homogenized with a Dounce homogenizer in 2 volumes of ice-cold lysis buffer (0.1 M Pipes/NaOH, pH6.6, 5 mM EGTA, 1 mM MgSO₄, 0.9 M glycerol, 1 mM DTT, 1 mM PMSF, 1 µg/ml
20 aprotinin, 1 µg/ml leupeptin and 1 µg/ml pepstatin). The microtubules are depolymerized by incubation on ice for 15 min, and the extract is then centrifuged at 16,000 g for 30 min at 4°C. The supernatant is recentrifuged at 135,000 g for 90 min at 4°C. Microtubules in this later supernatant are polymerized by addition of GTP to 1 mM and taxol to 20 µM and incubation at room temperature for 30 min. A 3 ml aliquot of the extract is layered on top
25 of 3 ml 15% sucrose cushion prepared in lysis buffer. After centrifuging at 54,000g for 30 min at 20°C using a swing out rotor, the microtubule pellet is resuspended in lysis buffer.

Microtubule overlay assays are performed as previously described (Saunders *et al.*, 1997). 500 ng per lane of recombinant Asp, recombinant polypeptide, and bovine serum albumin (BSA, Sigma) are fractionated by 10% SDS-PAGE and blotted onto PVDF

membranes (Millipore). The membranes are preincubated in TBST (50mM Tris pH 7.5, 150 mM NaCl, 0.05% Tween 20) containing 5% low fat powdered milk (LFPM) for 1 h and then washed 3 times for 15 min in lysis buffer. The filters are then incubated for 30 minutes in lysis buffer containing either 1 mM GDP, 1 mM GTP, or 1 mM GTP- γ -S.

5 MAP-free bovine brain tubulin (Molecular Probes) is polymerised at a concentration of 2 μ g/ml in lysis buffer by addition of GTP to a final concentration of 1 mM and incubated at 37°C for 30 min. The nucleotide solutions are removed and the buffer containing polymerised microtubules added to the membranes for incubation for 1h at 37°C with addition of taxol at a final concentration of 10 μ M for the final 30 min. The blots are then

10 washed 3 times with TBST and the bound tubulin detected using standard Western blot procedures using anti- β -tubulin antibodies (Boehringer Mannheim) at 2.5 μ g/ml and the Super Signal detection system (Pierce).

It may be desirable in one embodiment of this type of assay to deplete the polypeptide of the invention from cell extracts used to produce polymerise microtubules.

15 This may, for example, be achieved by the use of suitable antibodies.

A simple extension to this type of assay would be to test the effects of purified polypeptide of the invention upon the ability of tubulin to polymerise *in vitro* (for example, as used by Andersen and Karsenti, 1997) in the presence or absence of a candidate substance (typically added at the concentrations described above). *Xenopus* cell-

20 free extracts may conveniently be used, for example as a source of tubulin.

Microtubule Organising Centre (MTOC) Nucleation Activity Assays

Candidate substances, for example those identified using the binding assays described above, may be screening using a microtubule organising centre nucleation activity assay to determine if they are capable of disrupting MTOCs as measured by, for example, aster formation. This assay in its simplest form comprises adding the candidate

25 substance to a cellular extract which in the absence of the candidate substance has microtubule organising centre nucleation activity resulting in formation of asters.

In a preferred embodiment, the assay system comprises (i) a polypeptide of the invention and (ii) components required for microtubule organising centre nucleation activity except for functional polypeptide of the invention, which is typically removed by immunodepletion (or by the use of extracts from mutant cells). The components themselves are typically in two parts such that microtubule nucleation does not occur until the two parts are mixed. The polypeptide of the invention may be present in one of the two parts initially or added subsequently prior to mixing of the two parts.

Subsequently, the polypeptide of the invention and candidate substance are added to the component mix and microtubule nucleation from centrosomes measured, for example by immunostaining for the polypeptide of the invention and visualising aster formation by immuno-fluorescence microscopy. The polypeptide of the invention may be preincubated with the candidate substance before addition to the component mix. Alternatively, both the polypeptide of the invention and the candidate substance may be added directly to the component mix, simultaneously or sequentially in either order.

The components required for microtubule organising centre formation typically include salt-stripped centrosomes prepared as described in Moritz *et al.*, 1998. Stripping centrosome preparations with 2 M KI removes the centrosome proteins CP60, CP190, CNN and γ -tubulin. Of these, neither CP60 nor CP190 appear to be required for microtubule nucleation. The other minimal components are typically provided as a depleted cellular extract, or conveniently, as a cellular extract from cells with a non-functional variant of a polypeptide of the invention. Typically, labeled tubulin (usually β -tubulin) is also added to assist in visualising aster formation.

Alternatively, partially purified centrosomes that have not been salt-stripped may be used as part of the components. In this case, only tubulin, preferably labeled tubulin is required to complete the component mix.

Candidate substances are typically added to a final concentration of from 1 to 1000 nmol/ml, more preferably from 1 to 100 nmol/ml. In the case of antibodies, the final

concentration used is typically from 100 to 500 µg/ml, more preferably from 200 to 300 µg/ml.

The degree of inhibition of aster formation by the candidate substance may be determined by measuring the number of normal asters per unit area for control untreated cell preparation and measuring the number of normal asters per unit area for cells treated with the candidate substance and comparing the result. Typically, a candidate substance is considered to be capable of disrupting MTOC integrity if the treated cell preparations have less than 50%, preferably less than 40, 30, 20 or 10% of the number of asters found in untreated cells preparations. It may also be desirable to stain cells for γ -tubulin to determine the maximum number of possible MTOCs present to allow normalisation between samples.

Motor Protein Assay

Polypeptides of the invention may interact with motor proteins such as the Eg5-like motor protein *in vitro*. The effects of candidate substances on such a process may be determined using assays wherein the motor protein is immobilised on coverslips. Rhodamine labeled microtubules are then added and their translocation can be followed by fluorescent microscopy. The effect of candidate substances may thus be determined by comparing the extent and/or rate of translocation in the presence and absence of the candidate substance. Generally, candidate substances known to bind to a polypeptide of the invention, would be tested in this assay. Alternatively, a high throughput assay may be used to identify modulators of motor proteins and the resulting identified substances tested for affects on a polypeptide of the invention as described above.

Typically this assay uses microtubules stabilised by taxol (e.g. Howard and Hyman 1993; Chandra and Endow, 1993 – both chapters in “Motility Assays for Motor Proteins” Ed Jon Scholey, pub Academic Press). If however, a polypeptide of the invention were to promote stable polymerisation of microtubules (see above) then these microtubules could be used directly in motility assays.

Simple protein-protein binding assays as described above, using a motor protein and a polypeptide of the invention may also be used to confirm that the polypeptide of the invention binds to the motor protein, typically prior to testing the effect of candidate substances on that interaction.

5 *Assay for Spindle Assembly and Function*

A further assay to investigate the function of polypeptide of the invention and the effect of candidate substances on those functions is an assay which measures spindle assembly and function. Typically, such assays are performed using *Xenopus* cell free systems, where two types of spindle assembly are possible. In the "half spindle" assembly
10 pathway, a cytoplasmic extract of CSF arrested oocytes is mixed with sperm chromatin. The half spindles that form subsequently fuse together. A more physiological method is to induce CSF arrested extracts to enter interphase by addition of calcium, whereupon the DNA replicates and kinetochores form. Addition of fresh CSF arrested extract then induces mitosis with centrosome duplication and spindle formation (for discussion of
15 these systems see Tournebize and Heald, 1996).

Again, generally, candidate substances known to bind to a polypeptide of the invention, or non-functional polypeptide variants of the invention, would be tested in this assay. Alternatively, a high throughput assay may be used to identify modulators of spindle formation and function and the resulting identified substances tested for affects
20 binding of the polypeptide of the invention as described above.

Assays for DNA Replication

Another assay to investigate the function of polypeptide of the invention and the effect of candidate substances on those functions is as assay for replication of DNA. A number of cell free systems have been developed to assay DNA replication. These can be
25 used to assay the ability of a substance to prevent or inhibit DNA replication, by conducting the assay in the presence of the substance. Suitable cell-free assay systems include, for example the SV-40 assay (Li and Kelly, 1984, *Proc. Natl. Acad. Sci USA* 81, 6973-6977; Waga and Stillman, 1994, *Nature* 369, 207-212.). A *Drosophila* cell free replication system, for example as described by Crevel and Cotteril (1991), *EMBO J.* 10,

4361-4369, may also be used. A preferred assay is a cell free assay derived from *Xenopus* egg low speed supernatant extracts described in Blow and Laskey (1986, *Cell* 47,577-587) and Sheehan et al. (1988, *J. Cell Biol.* 106, 1-12), which measures the incorporation of nucleotides into a substrate consisting of *Xenopus* sperm DNA or HeLa nuclei. The nucleotides may be radiolabelled and incorporation assayed by scintillation counting. Alternatively and preferably, bromo-deoxy-uridine (BrdU) is used as a nucleotide substitute and replication activity measured by density substitution. The latter assay is able to distinguish genuine replication initiation events from incorporation as a result of DNA repair. The human cell-free replication assay reported by Krude, et al (1997), *Cell* 88, 109-19 may also be used to assay the effects of substances on the polypeptides of the invention.

Other In Vitro Assays

Other assays for identifying substances that bind to a polypeptide of the invention are also provided. For example, substances which affect chromosome condensation may be assayed using the *in vitro* cell free system derived from *Xenopus* eggs, as known in the art.

Substances which affect kinase activity or proteolysis activity are of interest. It is known, for example, that temporal control of ubiquitin-proteasome mediated protein degradation is critical for normal G1 and S phase progression (reviewed in Krek 1998, *Curr Opin Genet Dev* 8, 36-42). A number of E3 ubiquitin protein ligases, designated SCFs (Skp1-cullin-F-box protein ligase complexes), confer substrate specificity on ubiquitination reactions, while protein kinases phosphorylate substrates destined for destruction and convert them into preferred targets for ubiquitin modification catalyzed by SCFs. Furthermore, ubiquitin-mediated proteolysis due to the anaphase-promoting complex/cyclosome (APC/C) is essential for separation of sister chromatids during mitosis, and exit from mitosis (Listovsky et al., 2000, *Exp Cell Res* 255, 184-191).

Substances which inhibit or affect kinase activity may be identified by means of a kinase assay as known in the art, for example, by measuring incorporation of ³²P into a suitable peptide or other substrate in the presence of the candidate substance. Similarly,

substances which inhibit or affect proteolytic activity may be assayed by detecting increased or decreased cleavage of suitable polypeptide substrates.

Assays for these and other protein or polypeptide activities are known to those skilled in the art, and may suitably be used to identify substances which bind to a
5 polypeptide of the invention and affect its activity.

Whole Cell Assays

Candidate substances may also be tested on whole cells for their effect on cell cycle progression, including mitosis and/or meiosis. Preferably the candidate substances have been identified by the above-described *in vitro* methods. Alternatively, rapid
10 throughput screens for substances capable of inhibiting cell division, typically mitosis, may be used as a preliminary screen and then used in the *in vitro* assay described above to confirm that the affect is on a particular polypeptide of the invention.

The candidate substance, i.e. the test compound, may be administered to the cell in several ways. For example, it may be added directly to the cell culture medium or injected
15 into the cell. Alternatively, in the case of polypeptide candidate substances, the cell may be transfected with a nucleic acid construct which directs expression of the polypeptide in the cell. Preferably, the expression of the polypeptide is under the control of a regulatable promoter.

Typically, an assay to determine the effect of a candidate substance identified by
20 the method of the invention on a particular stage of the cell division cycle comprises administering the candidate substance to a cell and determining whether the substance inhibits that stage of the cell division cycle. Techniques for measuring progress through the cell cycle in a cell population are well known in the art. The extent of progress through the cell cycle in treated cells is compared with the extent of progress through the cell cycle
25 in an untreated control cell population to determine the degree of inhibition, if any. For example, an inhibitor of mitosis or meiosis may be assayed by measuring the proportion of cells in a population which are unable to undergo mitosis/meiosis and comparing this to the proportion of cells in an untreated population.

The concentration of candidate substances used will typically be such that the final concentration in the cells is similar to that described above for the *in vitro* assays.

A candidate substance is typically considered to be an inhibitor of a particular stage in the cell division cycle (for example, mitosis) if the proportion of cells undergoing that particular stage (i.e., mitosis) is reduced to below 50%, preferably below 40, 30, 20 or 10% of that observed in untreated control cell populations.

THERAPEUTIC USES

Many tumours are associated with defects in cell cycle progression, for example loss of normal cell cycle control. Tumour cells may therefore exhibit rapid and often aberrant mitosis. One therapeutic approach to treating cancer may therefore be to inhibit mitosis in rapidly dividing cells. Such an approach may also be used for therapy of any proliferative disease in general. Thus, since the polypeptides of the invention appear to be required for normal cell cycle progression, they represent targets for inhibition of their functions, particularly in tumour cells and other proliferative cells.

The term proliferative disorder is used herein in a broad sense to include any disorder that requires control of the cell cycle, for example, cardiovascular disorders such as restenosis and cardiomyopathy, auto-immune disorders such as glomerulonephritis and rheumatoid arthritis, dermatological disorders such as psoriasis, anti-inflammatory, anti-fungal, antiparasitic disorders such as malaria, emphysema and alopecia.

One possible approach is to express anti-sense constructs directed against polynucleotides of the invention, preferably selectively in tumour cells, to inhibit gene function and prevent the tumour cell from progressing through the cell cycle. Anti-sense constructs may also be used to inhibit gene function to prevent cell cycle progression in a proliferative cell. Another approach is to use non-functional variants of polypeptides of the invention that compete with the endogenous gene product for cellular components of cell cycle machinery, resulting in inhibition of function. Alternatively, compounds identified by the assays described above as binding to a polypeptide of the invention may

be administered to tumour or proliferative cells to prevent the function of that polypeptide. This may be performed, for example, by means of gene therapy or by direct administration of the compounds. Suitable antibodies of the invention may also be used as therapeutic agents.

5 Alternatively, double-stranded (ds) RNA is a powerful way of interfering with gene expression in a range of organisms that has recently been shown to be successful in mammals (Wianny and Zernicka-Goetz, 2000, Nat Cell Biol 2000, 2, 70-75). Double
10 stranded RNA corresponding to the sequence of a polynucleotide according to the invention can be introduced into or expressed in oocytes and cells of a candidate organism to interfere with cell division cycle progression.

 In addition, a number of the mutations described herein exhibit aberrant meiotic phenotypes. Aberrant meiosis is an important factor in infertility since mutations that affect only meiosis and not mitosis will lead to a viable organism but one that is unable to produce viable gametes and hence reproduce. Consequently, the elucidation of genes
15 involved in meiosis is an important step in diagnosing and preventing/treating fertility problems. Thus the polypeptides of the invention identified in mutant *Drosophila* having meiotic defects (as is clearly indicated in the Examples) may be used in methods of identifying substances that affect meiosis. In addition, these polypeptides, and
20 corresponding polynucleotides, may be used to study meiosis and identify possible mutations that are indicative of infertility. This will be of use in diagnosing infertility problems.

ADMINISTRATION

 Substances identified or identifiable by the assay methods of the invention may preferably be combined with various components to produce compositions of the
25 invention. Preferably the compositions are combined with a pharmaceutically acceptable carrier or diluent to produce a pharmaceutical composition (which may be for human or animal use). Suitable carriers and diluents include isotonic saline solutions, for example phosphate-buffered saline. The composition of the invention may be administered by

direct injection. The composition may be formulated for parenteral, intramuscular, intravenous, subcutaneous, intraocular or transdermal administration. Typically, each protein may be administered at a dose of from 0.01 to 30 mg/kg body weight, preferably from 0.1 to 10 mg/kg, more preferably from 0.1 to 1 mg/kg body weight.

- 5 Polynucleotides/vectors encoding polypeptide components (or antisense constructs) for use in inhibiting cell cycle progression, for example, inhibiting mitosis or meiosis, may be administered directly as a naked nucleic acid construct. They may further comprise flanking sequences homologous to the host cell genome. When the polynucleotides/vectors are administered as a naked nucleic acid, the amount of nucleic acid administered may typically be in the range of from 1 µg to 10 mg, preferably from 100 µg to 1 mg. It is particularly preferred to use polynucleotides/ vectors that target specifically tumour or proliferative cells, for example by virtue of suitable regulatory constructs or by the use of targeted viral vectors.

- 15 Uptake of naked nucleic acid constructs by mammalian cells is enhanced by several known transfection techniques for example those including the use of transfection agents. Example of these agents include cationic agents (for example calcium phosphate and DEAE-dextran) and lipofectants (for example lipofectamTM and transfectamTM). Typically, nucleic acid constructs are mixed with the transfection agent to produce a composition.

- 20 Preferably the polynucleotide, polypeptide, compound or vector described here may be conjugated, joined, linked, fused, or otherwise associated with a membrane translocation sequence.

- 25 Preferably, the polynucleotide, polypeptide, compound or vector, etc described here may be delivered into cells by being conjugated with, joined to, linked to, fused to, or otherwise associated with a protein capable of crossing the plasma membrane and/or the nuclear membrane (i.e., a membrane translocation sequence). Preferably, the substance of interest is fused or conjugated to a domain or sequence from such a protein responsible for the translocational activity. Translocation domains and sequences for example include

domains and sequences from the HIV-1-trans-activating protein (Tat), *Drosophila* Antennapedia homeodomain protein and the herpes simplex-1 virus VP22 protein. In a highly preferred embodiment, the substance of interest is conjugated with penetratin protein or a fragment of this. Penetratin comprises the sequence

5 RQIKIWFQNRRMKWKK and is described in Derossi, *et al.*, (1994), *J. Biol. Chem.* 269, 10444-50; use of penetratin-drug conjugates for intracellular delivery is described in WO/00/01417. Truncated and modified forms of penetratin may also be used, as described in WO/00/29427.

Preferably the polynucleotide, polypeptide, compound or vector according to the

10 invention is combined with a pharmaceutically acceptable carrier or diluent to produce a pharmaceutical composition. Suitable carriers and diluents include isotonic saline solutions, for example phosphate-buffered saline. The composition may be formulated for parenteral, intramuscular, intravenous, subcutaneous, intraocular or transdermal administration.

15 The routes of administration and dosages described are intended only as a guide since a skilled practitioner will be able to determine readily the optimum route of administration and dosage for any particular patient and condition.

The invention will now be further described by way of Examples, which are meant to serve to assist one of ordinary skill in the art in carrying out the invention and are not

20 intended in any way to limit the scope of the invention.

EXAMPLES

Generation and Identification of Lethal, Semi-Lethal and Sterile Third Chromosome Mutants Having Defects in Mitosis and/Or Meiosis, and Second Chromosome Mutants Having Defects in Imaginal Disc Development By P-Element

5 Insertion Mutagenesis

P-element mutagenesis

Transposable elements are widely used for mutagenesis in *Drosophila melanogaster* as they couple the advantages of providing effective genetic lesions with ease of detecting disrupted genes for the purpose of molecular cloning. To achieve near
10 saturation of the genome with mutations resulting from mobilisation of the P-lacW transposon (a P-element marked with a mini-white gene, bearing the *E.coli lacZ* gene as an enhancer trap, and an *E.coli* replicon and ampicillin resistance gene to facilitate ‘plasmid rescue’ of sequences at the site of the P-insertion), *Drosophila* females that are homozygous for *P-lacW* (inserted on the X chromosome) are crossed with males carrying
15 the transposase source P(Δ 2-3) (Deak et al., 1997). Random transpositions of the mutator element are then ‘captured’ in lines lacking transposase activity. Stable, or balanced, stocks bearing single lethal *P-lacW* insertions are made.

More than 41,000 lines are derived, of which approximately one-half are on the third chromosome. Originally some 3100 lethal or strong semi-lethal lines (in homozygous
20 conditions) are identified. During preliminary characterisation unstable lines and clusters of the same mutation event are eliminated leaving 2460 lines to be characterised.

Screening for Mitotic and Meiotic Defects

About half of the mutants in the collection are embryonic lethals. We have carried out cytological screens of the 1155 lines that comprise late larval lethals, pupal lethals,
25 pharate and adult semi-lethals for defective mitosis in the developing larval CNS. This has identified 69 mutations falling into 43 complementation groups that affect all stages of the mitotic cycle. The cytological screens involve examining orcein-stained squashed preparations of the larval CNS to detect abnormal mitotic cells. In lines where defects are

identified, the larval CNS is subjected to immunostaining to identify centromeres, spindle microtubules and DNA for further examination. This leads to clarification of the mitotic defect.

As a set of common functions are essential to both mitosis and meiosis, we then
5 identify mutations resulting in sterility and failed progression through male meiosis. This involves examining squashed preparations larval, pupal or adult testes by phase contrast microscopy. We examine “onion stage” spermatids in the 519 pupal and pharate lethal lines and 463 adult “semi-lethal” and viable lines for variations in size and number of nuclei which provides an indication of whether there have been defects in either
10 chromosome segregation or cytokinesis, respectively. A total of 54 lines of the 519 pupal and pharate lethal lines and 22 of the adult lines show such defects. However, another 67 lines show male sterility without having onion-stage defects. 12 lines showing onion stage defects have been scored as having mitotic defects in the independent cytological screen of squashed preparations of the larval CNS. Twelve further lines with onion stage defects
15 show female sterility and of these, 10 show maternal effect mitotic defects in syncytial embryos. Thus greater than one half of the meiotic mutants scored appear to represent cell division functions specific to male meiosis or have targeted male germ-line specific enhancer elements, thus revealing their meiotic function but in this test not their mitotic function.

20 Further characterisation of testis preparations of each line by phase-contrast microscopy with and without staining with Hoechst to reveal DNA defined 6 broad categories of meiotic mutants:

8 mutants from the collection show defects in meiotic entry or at early stages in the first meiotic division (MF1-8).

25 18 mutants (15 complementation groups) show abnormal meiotic spindles (AB1-16). Mutants in this group almost invariably show an associated weak defect in cytokinesis, and 7 show a strong defect in spermatid differentiation. 3 of these mutants

also show mitotic defects in larval brains or in embryos derived from homozygous mutant mothers.

18 mutants (16 complementation groups) also show abnormal meiotic spindles that are strongly multipolar (MUL1-15). Three of these also show maternal effect mitotic abnormalities of multipolar spindles in syncytial embryos.

4 mutants (3 complementation groups) show strong defects at all stages of spermatogenesis from the pre-meiotic stages to spermatid elongation stages (PL1-3). In this respect they resemble the *polo*¹ mutation.

4 mutants show segregation defects as indicated by spermatid nuclei of heterogeneous sizes (SEG1-4). The spindles appear normal but all have what are either chromosome bridges or lagging chromosomes. One of these also shows a maternal effect.

9 mutants (7 complementation groups) show predominant cytokinesis defects. Two complementation groups also have cytokinesis defects in mitotic cells in the larval brain.

In the Examples below, the designations MF, AB, MUL, PL, SEG or CK are included in the category description where available. Further phenotype information for each mutant described in the results section is provided in the "Phenotype" field. There is considerable overlap between these categories, and it will be of much interest to distinguish between mutants in which the primary defect results in secondary consequences, and mutants that affect more than one aspect of spermatogenesis, as for example appears to be the case with *polo* mutants (Sunkel and Glover, 1988; Carmena et al, 1998).

In the Examples, lines exhibiting mitotic and meiotic phenotypes are categorised generally into four categories:

Category 1 : Failure to complete cytokinesis

Category 2 : Failure to enter M-phase

Category 3: Metaphase arrest

Category 4: Anaphase defect

Category 5: Small Imaginal Discs (Block to Proliferation; see below)

5 Category 1 phenotypes are exhibited by mutations in Examples 1 to 14; while
Category 2 phenotypes are exhibited by mutations in Examples 15 to 19. Category 3
phenotypes are exhibited by mutations in Examples 20 to 30, Category 4 phenotypes are
exhibited by mutations in Examples 31 to 53. Mutations in Examples 54 to 74 exhibit a
Category 5 phenotype.

10 *Generation and identification of second chromosome mutants having small or
no imaginal discs.*

In the case of the second chromosome the flies used were from a second
chromosome P-element collection established in Szeged, Hungary (Torok et al., 1993).
The process of P-element insertion mutagenesis is essentially as described above. 15475
15 insertions were recovered, of which 2711 were lethal or semi-lethal. After elimination of
clusters of identical mutants, 2399 lines representing 1748 independent lethal insertions
were recovered. Lines were chosen from the second chromosome collection on the basis
of having small or no imaginal discs, to indicate a disruption in cell cycle progression that
leads to underdevelopment of the discs. All the second chromosome mutants referred to in
20 the results section are noted under the "Phenotype" field as "second chromosome, small
imaginal discs" and comprise Category 5.

Cytological Mapping of the P-Element Insertion Sites

The site of insertion of the P-element in each mutant line was determined by *in situ*
hybridisation of P-element DNA to salivary gland polytene chromosomes as described in
25 Saunders et al., 1989. Wandering third stage larvae were dissected and fixed as described
and incubated with biotin-labeled DNA made from the *P-lacW* plasmid. After signal

detection chromosomes were stained with Giemsa and examined by microscopy and signals indicating the presence of P elements were assigned to polytene chromosome bands referring to the Bridges map (Lefevre, 1976). In the majority of cases a single P element was detected, only 10% of lines having multiple (two or three) insertions. The site of insertion is given as the "Map Position" field in the results section (for example 77B)

Plasmid Rescue of P-Elements from Mutant Drosophila Lines

Genomic DNA was isolated from adult flies by the method of Jowett et al., 1986, and plasmid rescue from the genomic DNA was performed according to Pirrotta et al., 1986. This allows the recovery of genomic DNA adjacent to the P-element which facilitates the identification of the site of P-element insertion and of genes which may be disrupted by the insertion. Essentially, genomic DNA derived from about 200 flies was digested with a restriction enzyme known to have a site within the P-element (EcoRI or SacII for cloning sequences to the left of the element, or XbaI, BglII, PstI or BamHI for sequences to the right of the element). The digested DNA was ligated overnight, and plasmids recovered by electroporation of the ligated DNA into *E.coli* XL1-blue competent cells. Appropriate primers from within the P-lacW sequence were used to determine the sequence of the genomic DNA flanking the element (on average, 400 bp of sequence were obtained). The rescue sequences are provided in the results section under the heading "Rescue sequence". Where more than one sequence was recovered, the orientation of each sequence is also given.

Sequence Analysis of P Element Insertion Lines

Sequences flanking the insertion site of the P-element were derived by P element rescue as described above. In some cases sequence was obtained from only one side of the insertion, while in other cases sequences were obtained from both sides of the insertion.

As a first step, each P element rescue sequence was used to search a total database of *Drosophila melanogaster* sequences (database of the Berkley *Drosophila* Genome project) using the BLASTN program (which compares a nucleic acid sequence with a nucleic acid database, (Altschul and Lipman 1990)) with default parameters.

The search may identify a number of different types of match including *Drosophila* ESTs, known *Drosophila* genes and cloned genomic regions.

The ability to identify genes already known to be essential for cell cycle progression using this approach was confirmed, in this example, by the rescue sequence
5 obtained from line 1324/8 which mapped to the 77B locus which was used to search the database. A BLASTN search identified a number of matching *Drosophila* ESTs, a match with the known cell cycle regulatory gene *polo* and a cloned genomic region designated CSC: AC018188. These matches are recorded in the results sections under the field headings "*Drosophila* ESTs", "*Drosophila* gene hit" and "Genomic hit, Accession No.",
10 respectively. Any entries under "*Drosophila* gene hit" are further annotated with "(BLASTN with Rescue sequence)" to show that the match was obtained using the rescue sequence rather than a *Drosophila* EST or genomic clone ORF (see below). Accession numbers of ESTs, genes and genomic clones are provided where known. Genomic clones designations often include the Genbank designation as part of a longer designation.
15 However the Genbank designation is always the code beginning with "AC" and followed by six digits.

Where an EST was identified, this was subsequently used to search using the BLASTX program (default parameters) against databases of sequences from *Drosophila* and Homo sapiens (databases of the National centre for Biotechnology Information
20 (NCBI), National Library of Medicine, National Institute of Health, USA). In the case of line 1104/16, the search identified a known human gene, phosphatidylinositol transfer protein (accession no. P48739) implying a novel function for this protein in cytokinesis. Human Homologues identified as a result of a BLASTX search using a *Drosophila* EST are shown in the results section under the heading "Human homologues" and annotated
25 with "(BLASTX with EST)". *Drosophila* genes identified as a result of a BLASTX search using a *Drosophila* EST are shown in the results section under the heading "*Drosophila* gene hit" and annotated with "(BLASTX with EST)".

Where no *Drosophila* gene was identified using the initial BLASTN search but a matching genomic clone was found (a Bac or P1 clone often in excess of 100 kilobases), a

20 kilobase segment of this genomic sequence (10 kilobases either side flanking the site of the P-element insertion) was subjected to a number of analyses.

If the rescue sequence matched sequences that lie within a known gene present within the genomic clone then these are presented under the heading "*Drosophila* gene hit
5 (BLASTN with Rescue sequence". The known gene sequence was then used in a BLASTX search of a human database (NCBI – see above) to identify any human homologues. These are shown in the "Human homologue" field and annotated with "(BLASTX with *Drosophila* gene)".

If the rescue sequence does not match any sequences that lie with a known gene
10 within the genomic clone then the occurrence of ORFs within the 20 kilobase genomic segment was predicted using the Genscan programme (Burge and Karlin, 1997). Where the P-element was observed to be inserted into the coding region or within the 5' untranslated region (which we defined as within 2 kilobases of the predicted start of the coding region) we assume the P element to be capable of disrupting the expression of the
15 predicted gene. Each predicted open reading frame (or predicted coding sequence) was then used to search *Drosophila* and human databases using the TBLASTN program (compares a protein query sequence against a nucleotide sequence database dynamically translated in all reading frames) and/or the TBLASTX program (compares a nucleotide query sequence dynamically translated in all reading frames against a nucleotide sequence
20 database dynamically translated in all reading frames) to determine whether the predicted open reading frame corresponded to a known gene. Typically, TBLASTX is only used when no matches are found using TBLASTN.

Where the TBLASTN search found a known *Drosophila* gene, then this is indicated in the results in the "*Drosophila* gene hit" field, annotated with "(TBLASTN
25 with predicted ORF)". The *Drosophila* gene sequence was then typically used to search a human database (NCBI – see above) to identify any human homologues using BLASTX. These are shown in the "Human homologue" field and annotated with "(BLASTX with *Drosophila* gene)".

Where the TBLASTN and/or TBLASTX search found a known human gene, then this is indicated in the results in the "Human homologue" field, annotated with "(TBLASTN (or TBLASTX) with predicted ORF)".

5 If the TBLASTN and/or TBLASTX search found no *Drosophila* or human genes, then it was assumed that the original ORF corresponds to a novel gene. If the TBLASTN search found no *Drosophila* genes but identified a human homologue, then it was assumed that the original ORF corresponds to a novel *Drosophila* homologue of a known human gene.

10 ***Additional Sequence Analysis using the Annotated D. melanogaster Sequence (GadFly).***

Rescue sequences were also used to search the fully annotated version of the *Drosophila* genome (GadFly; Adams, et al., 2000, Science 287, 2185-2195), using GlyBLAST at the Berkeley *Drosophila* Genome Projects web site to identify the genome segment (usually approximately 200-250 kb) containing the P-element insertion site. The
15 graphic representation of the genomic fragment available at GadFly allows the identification of all real and theoretical genes which flank the site of insertion. Candidate genes where the P-element is either inserted within the gene or close to the 5' end of the gene were identified. In GadFly, the *Drosophila* genes are given the designation CG (Complete gene) and usually details of human homologues are also given. In most cases,
20 this data confirms the data derived from the sequence analysis procedure described above, and in some cases new data is obtained. Where available both sets of data are included in the individual Examples described below. To identify further candidate human homologues, BLASTP (amino acid query sequence against amino acid database) searches with *Drosophila* sequences are used against the human genome project database and also
25 the Ensembl dataset. The Ensembl dataset comprises GeneWise gene predictions using a protein template where possible or Genscan followed by BLAST confirmation via protein, cDNA or EST hits. These are matched using WUBLASTP with default parameters (Altschul et al., 1990, *J Mol Biol* 215, 403-10). The results are filtered to contain only potential homologues. Only matches with the identity of more than 50% and length of
30 more than 50 amino acids are included.

Confirmation of Cell Cycle Involvement of Candidate Genes Using Double Stranded RNA Interference (RNAi)

P-elements usually insert into the region 5' to a *Drosophila* gene. This means that there is sometimes more than one candidate gene affected, as the P-element can insert into the 5' regions of two diverging genes (one on each DNA strand). In order to confirm which of the candidate genes is responsible for the cell cycle phenotype observed in the fly line, we use the technique of double stranded RNA interference to specifically knock out gene expression in *Drosophila* cells in tissue culture (Clemens, et al., 2000, *Proc. Natl. Acad. Sci. USA*, 6499-6503). The overall strategy is to prepare double stranded RNA (dsRNA) specific to each gene of interest and to transfect this into Schneider's *Drosophila* line 2 to inhibit the expression of the particular gene. The dsRNA is prepared from a double stranded, gene specific PCR product with a T7 RNA polymerase binding site at each end. The PCR primers consist of 25-30 bases of gene specific sequence fused to a T7 polymerase binding site (TAATACGACTCACTATAGGGACA), and are designed to amplify a DNA fragment of around 500bp. Although this is the optimal size, the sequences in fact range from 450 bp to 650 bp. Where possible, PCR amplification is performed using genomic DNA purified from Schneider's *Drosophila* line 2 as a template. This is only feasible where the gene has an exon of 450 bp or more. In instances where the gene possesses only short exons of less than 450 bp, primers are designed in different exons and PCR amplification is performed using cDNA derived from Schneider's *Drosophila* line 2 as a template.

A sample of PCR product is analysed by horizontal gel electrophoresis and the DNA purified using a Qiagen QiaQuick PCR purification kit. 1µg of DNA is used as the template in the preparation of gene specific single stranded RNA using the Ambion T7 Megascript kit. Single stranded RNA is produced from both strands of the template and is purified and immediately annealed by heating to 90 degrees C for 15 mins followed by gradual cooling to room temperature overnight. A sample of the dsRNA is analysed by horizontal gel electrophoresis.

3µg of dsRNA is transfected into Schneider's *Drosophila* line 2 using the transfection agent, Transfect (Gibco) and the cells incubated for 72 hours prior to fixation.

The DNA content of the cells is analysed by staining with propidium iodide and standard FACS analysis for DNA content. The cells in G1 and G2/S phases of the cell cycle are visualised as two separate population peaks in normal cycling S2 cells. In each experiment, Red Fluorescent Protein dsRNA is used as a negative control. In some cases

5 the phenotype is confirmed by fixing cells on poly-lysine covered slides which are then stained for DNA using DAPI and for tubulin using an anti-tubulin antibody YL1/2 and appropriate fluorescent secondary antibody to visualise aberrant mitoses.

It should be noted that RNAi could not confirm phenotype in all cases. This is to be expected as the method relies on the ability of dsRNA to prevent new protein

10 expression. Consequently, it is necessary that S2 cells express the specific cDNA of the gene in question, and also that the protein is turned over rapidly. It would therefore probably be difficult to sufficiently reduce levels of very stable proteins using this approach.

The layout of a typical entry in the results section is shown below. Not all fields

15 present in the actual results section contain information for each individual *Drosophila* line described.

TYPICAL RESULTS LAYOUT

20	Line ID	- <i>Drosophila</i> line designation
	Category	- Description of phenotype
	Reversion	- R = revertant, NR = non revertant, ? = not determined
	Map Position	- according to the Bridges map (Lefevre, 1976).

25	Rescue ID
	Rescue Sequence
	[nucleotide sequence]

Genomic hit, Accession No.

30	Associated ORF
	GENSCAN_predicted_peptide [results of Genscan - amino acid sequence]
	GENSCAN_predicted_CDS [results of Genscan nucleotide sequence]

35	<i>Drosophila</i> Gene Hit
	(BLASTN with rescue sequence)

(TBLASTN (or TBLASTX) with predicted ORF)
(BLASTX with EST)

Human Homologue

- 5 (BLASTX with *Drosophila* gene)
(TBLASTN (or TBLASTX) with predicted ORF)
(BLASTX with EST)
Drosophila EST

- 10 Annotated *Drosophila* genome genomic segment
Annotated *Drosophila* genome Complete gene candidate
Human homologue of Complete gene candidate

- Putative function Derived from homologies or *Drosophila* experimental data
15 Confirmation by RNAi Description of Facs analysis DNA content profile

A specific example is as follows:

- 20 **Line ID** 1324/8
Category Mitotic defects in brain: metaphase arrest
 (overcondensation, some circular chromosomes, no anaphases,
 very high mitotic index, metaphase (or less aligned) with bipolar
25 spindle, no CP190 staining)
Reversion R
Map Position 77B

Rescue ID B1E
30 **Rescue Sequence**
GTTTTGCCCATCGATTGCACGAAAACCAAGCACAAAGCGGAGAACGCGCCGA
AACCGTTCGATTTTTTAAATGCCAAAATGAATTGGACGTGAAGCGTCAGCTGA
ATTGGTGTGCCCGTTTCGGTGGCTATCGCACACTTTCTGGTATTTATCGCGGTA
TTTTGTTGAGTGTTGAACAACAAATTCTATGGCCGTTACCCTTTTGAATTTACT
35 TACTGGCGTTTACTCTGTTTGAATTGAGCGCAATATTTTTTCTATTGCTCTGC
GCAACACTGTGTTTTAACCGCTATTTATTTGAAAATCTACAAAACTAACCGTT
TACATTTTTGAAATTTCCAAAAGGGTTTTCCATAAATTGAGTTTTACTAAAACC
AGTCCAACGGTCCAACTTTATATTGTTAGAAGCCCCTTTTCTAATTTGAATTG
GCTTGCAAACGTTTTCTGAATTTAAAAATACTGCCACCCTTGTTAATTGCAGG
40 TTTTCCGAATCCCTGATTTGTTGTTTTAAAAAGAAAATTTATTAGAAACAGCTA
TCTCAACC

Genomic hit, Accession No. CSC:AC018188

Drosophila Gene Hit Polo (X63361)

- 45 **Human Homologue** BLASTX PLK-1 (P53350)
Drosophila EST several including LD11851 (AA392613) which match polo

Annotated *Drosophila* genome genomic segment AE003514

- Annotated *Drosophila* genome Complete gene candidate** CG12306
Human homolog of Complete gene candidate 1e-169 1709658 P53350
PLK1_HUMAN
SERINE/THREONINE-
5 PROTEIN KINASE PLK
(PLK-1)
- Putative function** Serine/threonine kinase known to be required for mitosis
- 10 **Confirmation by RNAi** Reduced G1 and G2/M peaks indicating fewer cycling cells, microscopy analysis of DNA and tubulin staining identified monopolar spindles characteristic of polo mutation in *Drosophila*.

CATEGORY 1: FAILURE TO COMPLETE CYTOKINESIS

Example 1 (Category 1)

5	Line ID 1031/14 Category Mitotic defects in brain: cytokinesis defect (polyploidy) Reversion R Map Position 74B
10	Rescue ID 2A3B Rescue Sequence 1 CCCCCGAACATATGTTCAAGTGTGGCCGCAGCAGAGTTGTCAAAACACGCTCCC CAATGAAATAACCTAAATGTGCCATCACTGTTACTTAACAGTTTCTGTTACTTT TCTAGCGGCATGTCAAAAAAACAAAAATATAGAAAATGCTAAATATATATTG 15 GACTAATGTGTTTAAATGTAACCTTACACTAGTAACAGATCCCCATTAATAAAA GCCAAACTCTAAAATTCTGCCACAAGTACTATTTCTCACGTAACACCTTACTA ACGGATTTTACATGATATCTACGACAAGAACTGTTTGCTGATATAAAATTGC TATCACCGCTTTCCGTAAACACTTTTACACTGATGGATTACAAGTTCAATTAAT ACATCAACTTACCTTAACAATTTTAAGACAACCTAACACTCCCACAATTTAATT 20 CAACCTACACCGCTTGATAATCAGCTGTTCTGTACAAAAACAATAACACTGT TAACAACAGCGCACAGTGGATAATACAGTCCTAAAGGCAATATACCCATTG GCATTTTT
25	Rescue ID 2A3S Rescue Sequence 2 TTCCGGGGAGAATGGCTGCGATTTTCGCGTCGGTAAAAATAGCAAATACTCGTTA ATGTGCTGTGGGAACGCTTCCTCCCCGGCCCCAAAGTGGCCCCGAAGAAAGTGA GCAAATGTGCGCGCCGCAAGATAGTCGCCGCCGAACAAACGATAGTGACGAAA GTGATTTAATTCAACTACCAGCACTCCCGCAAATACGATGAGTATGTCGCGCGG 30 CGGCAACACAACCTCTGGACTTGCAGCCGCTCCTGGCGGAGAGCGATGTCGGAA ACAGGGAGCTGGAGGAGAAGATGGGCGGATCGGCGGATCGGTCATCGCTGCTC GATGGATCCGGTTCGAAGGAGCTGAGTCACCGGGAACGCGAGGACTCGGCGTT GTTCGTCAAGAAGATCGGGAGCGCCTTGTTCTATGGCTTGTCTCCTTCATGATT ACGGTGGTAAACAAGACGGTGCTTACCTCCTACCACTTCCCCTCGTTCCTGTTCC 35 TCAGCCTCGGGCAACTTACTGCTAGCATTGTGGTCCTGGGCATGGGCAAAGCGC CTGAAAATGGTGAACCTTTTCCCTTTTGCAGAGGAATACCTTCGCCAAGATCTTT CCGCTGCCACTGATATTTCTGGGAAACATGATGTTTGACTGGGTGGCACAAAA ACCTTGAGTCTGCCCATGTTTCGCAGCCCTACGAC
40	Genomic hit, Accession No. AC019515

Associated ORF

Genscan ORF1 predicted sequences:>15:31:57|GENSCAN_predicted_peptide_4|373_aa

MSMSRGGNTTLDLQPLLAESDVGNRELEEKMGGSADRSSLDDGSGSKELSHRER
 EDSALFVKKIGSALFYGLSSFMITVVNKT VLT SYHFPSFLFLSLGQLTASIVVLGMG
 KRLKLVNFPPLQRNTFAKIFPLPLIFLGNMMFGLGGTKTLSLPMFAALRRFSILMT
 MLELKILGLRPSNAVQVSVYAMIGGALLAASDDL SFNMRGYIYVMITNAL TASN
 5 GVVYKKKLDTSEIGKYGLMYNSL F MFLPALALNYVTGNLDQALNFEQWNSV
 FVVQFLLSCVMGFILSYSTILCTQFNSALTTIVGCLKNICVTYLG MFIGGDYVFSW
 LNCIGINISVLASLLYTYVTFRRKRAPDKQDHL PSTRGENV

>15:31:57|GENSCAN_predicted_CDS_4|1122_bp
 10 atgagtatgtcgcgcgggcggaacacaactctggacttgagccgctcctggcggagagcgatgtcggaacagggagctgga
 ggagaagatggcgggatcgcggaatcggtcatcgctgctgatccggctcgaaggagctgagtcaccgggaacgcgag
 gactcggcggtgttcgtcaagaagatcgggagcgctgttctatggctgtcctcctcatgattacgggtgtaacaagacgggtgc
 ttacctctaccacttccctcgttctgttctcagcctcgggcaactfactgctagcattgtgttcctgggcatgggcaagcgcct
 gaaattggtgaacttccccctctgcagaggaataccttcgcaagatcttccgctgccactgataattctgggaaacatgatgtttg
 15 gactgggtggcacaaaaaccttgagtctgcccattgtcgcagccctacgacgcttctctatcctgatgacctgtgctggagctca
 agatcctgggactgcgacctcgaatgcggttcaggtcagcgtatacgaatgatcggtggagcgctgctggccgcctctgatga
 tctgtcctcaacatgaggggctacatctatgtgatgactaactgccttgaccgctcgaatggcgtatgtgaagaaaaaactc
 gacacctcggagatcggaagtacggcctaatagtactacaactcgtgttatgtttctgcctgccctggccctcaactatgttacag
 ggaatctagatcaggcgctgaacttgaacaatggaatgactcagtggttggtgcagttcctgctcagttgcgttatgggttcac
 20 ctatcgtacagcaccatcctgtgcacgaattcaactcggcgctgaccaccaccattgtgggatgcctgaaaaacatctgcgtaac
 atatctgggcatgttcattggaggcgactacgtcttctcgtgctcaactgtattgggatcaacatcagcgtgctggttagtctgctc
 acacgtacgtcacttttcggcggaagcggtcctccgataagcaggaccactgccagcaccgcggcgagaatgtctag

25 **Human Homologue** (TBLASTN with ORF1): KIAA0260 gene (D87449) and putative
 Sqv-7-like protein (AJ005866)
Drosophila EST CK00510 (AA140776)

Annotated Drosophila genome genomic segment AE003524
Annotated Drosophila genome Complete gene candidate CG3874 – novel glucose-6-
 30 phosphate transporter

Human homologue of Complete gene candidate EMBL:D87449 protein
 KIAA0260_id:BAA13390
 gi:166578 Similar to a
 35 C.elegans protein encoded in
 cosmid C52E12 (U50135) and
 Ensembl predicted gene
 ENSG00000024527
 Clone:AL133320
 40 Contig:AL133320.00001
 8.10E-95

Putative function Sugar modification protein similar to proteins involved in
 Drosophila cytokinesis and signalling

45 **Confirmation by RNAi** Marked increased G1 and S peak indicating mainly arrest in
 G1

Example 2 (Category 1)

	Line ID	1066/5
5	Category	Male semi-sterile, Meiotic defects in testis: cytokinesis defects, segregation defects. (Seg-01/62)
	Reversion	?
	Map Position	89B
10	Rescue ID	F9E
	Rescue Sequence	GTATACCATTAGAGAATATGATGAAGAAGGACTGTAAGAAGATCCTTCAGTG AATTTGACTGCTGACGTCGATCGGAACCTTGCTGCGCTGACGTACAAAATCGCG AAGTGAATAAATAATATGGATGAGACCTGTTTCGCCGACATATACAATAGTG 15 CTCAAGACCTAATGGAATTATACGTTAATAACCAGCCACATTTCTTAGATATTT CTAATATGAGCCATCTGCTGCAGGTTCTTTCCAATATCTAATTCTAGATCTTCT TCGAATACGACCTTTTTGGCCATGAAACGATGATTTGCCACTTCATTACAAAG CATTAAATTTGTCATGATTCTCTTAAGCGTGCACTTTATCTGAAAGTCTGAACAG CTGGCTGCGAAATGGATCCCCGGGATTGGAGATGGCAAGTAAATCTGTCCCTCG 20 CTACAAACAAGTGGGCACCACTGGGCATTCGGGGAATAGGGATATGGGTTGG GAATGGGGATATATTGTGGCATTGGCGAAAGGTCGCTATGC

Genomic hit, Accession No. CSC:AC019750

25	Associated ORF	>16:04:57 GENSCAN_predicted_peptide_4 418_aa MKPIPNESKGTAAVGDATVVHVDVCTLFAVELDPYLRSSMGMRTTRRAQSGALLL QLLAVADGGFAAHICACKCRLRLPHVTCCNRPFKATAKAKGQAVSSTKPNQL CFHGCCGWIITTKGETFTENSPTSMSGFAWERHSLGECVVVAGTEQILLIGRTLIGR 30 MSHTQTDSTSPFVVDCHSQLCGSKCKCICVSVGFCVRPSCQRFDMKIVWANLAM QKRFLLGAAIADMCCRNSVIWCKLQLDPVKPIDERADGSLALVTKVCDNNNIV HYVVVAGVTGSQSRSLQPLRSGQNESTEQWPRTKGEGGFNNNSRNNKHSAPT QEQQELWQKQLLQDQRDDCHASGSFQSASFAETRSFTFDDTTAHSEFCFRTRAEK RRILVLLETSIKLKPDKYATSGHTRRCAIGLLHSII
35		>16:04:57 GENSCAN_predicted_CDS_4 1257_bp atgaaccattccaacgaatcaagggaacccttgcggcagttggagatgctactgttgcacgtgtgtactttgttgcg tagagcttgatccctatctcaggagcagcatgggaatgaggacgctagagctcaaagcggcgtctgttattacagctccttgcg gttgcgcatggaggtttgtgctcatattgtgctgcaagtgtcggcttgccttgcacatgtcacatgttgcacacgggaatcct 40 ttcaaggcaactgcacaaaggtcaggcggcagctccactaaacaaaccagcttgccttcacggctgctgtgctggat aattactacaaaggtgaaacgttcacgaaaactcggcagcatcatgagcggtttgcgtgggagcggcatagccttggtagt gcgtggttgcgtggaacggaacaaatcctgctgattggcaggacattgattggcgcagcagcactcaactgattcgacc agcccccttgcgtgactgtcactgcacactgtcggctcgaagtgcacatgtatctgtgtatctgttaggtttctgtgctgcccgtct tgtcagcgccttgacatgaaaatagtttgggcaacttggctatgcacaaagcgatttctattaggagccgccaatgccgacatgtgct 45 gccgaaatcggtgatttgggtgcaaaactgcagctagatccagtcgaagcaattgacgaaagagccgacggcagcggtcttgcact ggttacaaagtatgcgataacaataacatcgtccactatgtgtcgttgcgtgggggttacgggcagtcagtcacggtcacggctgc

aaccctccgctccggccaaaacgagtcacagaacaatggccaaggacgaagggggggaggggggattcaataacaaca
 gcaggaacaacaaacattctgtcccacgcaagagcagcaggaactgtggcaaaacagctgctgcaggatcaacgagacgat
 tgtcatgccagtggagcttcagctgtcgtcattcgggagacgcgtagtttcacgttcgacgacacaaccgctcacagcgaattt
 5 tgtttcggactagagctgagaaacggcgaattttggtgcttctggaaacatcgattaactaaaacccgataagtatgcgacaagc
 ggtcacactcggcgatgtgcgataggattgctgcattcgattatag

- Drosophila* Gene Hit** rescue sequence: mitotic heterochromatin fragment clone CH(2)6
 (L36595) and subtelomeric heterochromatin repeats (L03284).
 TBLASTN with ORF1: nebula (nla) (AF147700)
 10 **Human Homologue** BLASTX with nebula: Down Syndrome candidate region 1-like
 protein 2 (AF176117)
***Drosophila* EST** rescue sequence: CK01138 (AA141069)

- 15 **Annotated *Drosophila* genome genomic segment** AE003712
Annotated *Drosophila* genome Complete gene candidate CG6072 - nebula
 CG6046 - sap18

- 20 **Human homologue of Complete gene candidate** CG6072- 8e-36 'ZAKI4 a thyroid
 hormone responsive gene in human
 skin fibroblasts' also known as
 DOWN SYNDROME CANDIDATE
 REGION 1-LIKE 1; DSCR1L1
 EMBL:D83407
 25 protein_id:BAA11911 gi:143504

- 30 CG6046- 3e-45 2108210 (U96915)
 sin3 associated polypeptide p18
 [Homo sapiens] and gi5032067
 C7E479FFE9CA5774
 [ref]NP_005861.1| sin3-associated
 polypeptide, 18kD [Homo sapiens]
 (1.90E-43)

- 35 **Putative function** Nebula unknown function, Sap18 transcription factor

- Confirmation by RNAi** Both show reduction in G1 and G2/S peaks indicating fewer
 cycling cells

Line ID 234/50
Category Meiotic defects in testis: cytokinesis defects, abnormal spindles.
(Ab-02/12)
Reversion R
5 **Map Position** 89B

Rescue ID 2C5E
Rescue Sequence
10 GTTTGACTGCTGACGTCGATCGGAACCTTGCTGCGCTGACGTACAAAATCGCGA
AGTGAATAAATAATATGGATGAGACTCCTGTTTCGCCGACATATACAATAGTG
CTCAAGACCCTAATGGAATTATACGTTAATAACCAGCCACATTTCTTAGATAT
TTCTAATATGAGCCATCTGCTGCAGGTTCTTTCCAATATCTAATTCTAGATCTT
CTTCGAATACGACCTTTTTGGCCATGAAACGATGATTTGCCACTTCATTCACAA
GCATTAATTTGTCATGATTCTCTTAAGCGTGCACTTTATCTGAAAGTCTGAACA
15 GCTGGCTGCGAAATGGATTCCCCGGATTGGAGATGGCAAGTAAATCTGTCCTC
GCTACAAACAAGTGGGCACCACTGGGCATTCTGGGGAATAGGGATATGGGTTG
GAAA

Drosophila EST rescue sequence: CK01138 (AA141069)
20 All other entries as for 1066/5.

Example 3 (Category 1)

Line ID 1104/16
Category Mitotic defects in brain: cytokinesis defect
 5 (no overcondensation of diploids, high polyploidy)
Reversion R
Map Position 92A

Rescue ID B5P
 10 **Rescue Sequence 1**
 CTCCGGACACGCAGTAGCTAAATAACAACTCATTACTAGTATATTACTGCCG
 CCGATTTGCAAGCGCGTACCGATCCCCGATACCAGGCCAATCGCACTCCCCAGT
 TGTACGTCATCACTTAAGTAATAAATCAGCGGCAAATCGCATAAATTGCTATT
 GATATTCGCCCCGCTGTGTGTGCGTGTGTATTTGCAAGAGAGTGTGTGTGTGT
 15 GTGTGCATATGACTCGTGCGTTTAGCCGACAATTGGAGAAAAAGCATTACCAA
 TCCCAATTGGCTAACTAACTAAAGTTGGCTTGGCCAAACATAAACAAAAAGT
 GCGGGCGCAGCGATTTGGCAGCGAAACATATACACCAAAGCGCTATTGGCAG
 ATATATATGTAGATTAAATATAGAAAGTGCGTGCGAAGGTTAAGAGTCGAGT
 GCAAGTGCATTTATATTTGGAAATAATAAATGCTACAAT
 20

Rescue ID B5E
Rescue Sequence 2
 GTCCGGAGCGGAGCTAAAGTTCGATGTTTCGTGCAAAACACTTCGATTCCGATA
 GGCGGATGCTATCGATTTCCGGCGATGCCCGTTGGTCACACTTGGTGGTGGGCG
 25 CTGCCCCGTCGCCGACTATCGATAGCACAAGCGGGTTATTTAGGTGTGCGCAGC
 TTGTAAGGGTGACTCATGCTGTATAAAATTATTATAAAAAGTTAATGAATATAA
 TATAGTTATAATAAAATTATATATAAATCTATAAGATCAAAGATCATCAGTTA
 TCATTTATCATTTGATTATATGAAAAACAAGAACAGAAACAAGATTTAATAGG
 TTTTTGAAATGTGAAAATGTGGGTTACCCCCAATTCTTATTCGAAATTAAATAA
 30 CCTAAAGAACAGTTATACACAGATAGGTAATTTGCACATAAGCCAAATTTTGT
 CTAGAATTCCGCGGAATTAATTCTTGAAGACGAAAGGGCCTCCGTGATACGCC
 TATTTTATAGGTTAATGTCATGATAATAATGGTTTCTTA

Genomic hit, Accession No. AC006589
 35 **Associated ORF**
 Genscan: ORF1 predicted sequences
 >/tmp/aaaaainga|GENSCAN_predicted_peptide_2|850_aa
 MATRGANVIWFRHGLRLHDNPALLAALADKDQGIALIPVFIFDGESAGTKNVGY
 NRMRFLLDSLQDIDDQLQAATDGRGRLLVFEGETPAYIFRRLHEQVRLHRICIEQDC
 40 EPIWNERDESIRSLCRELNIDFVEKVSHTLWDPQLVIETNGGIPPLTYQMFLIRCTH
 HNGDVNGDEDTGEGEGTGGRIDWAKEGACWRAGNSDEQECQACQSVSSVIMM
 VLQYSNPAHHCQLLECLMTLKHNVVKDILCVVAYGTAVSRTSAKLLFFYYWP
 AFNANLFDKRVLLSKLTNDLVPFTCQREHCPNSGNAEAAKVCYDHSISIAYAPDC
 PPPLYLCIECANEIHRHGSLEFGDILHPMQQVSMVCENKNCRSNEKSAFSICFSTE
 45 CASFNGNHPIRYCSQCHSNRHSRRGGDHVVHRSLOPAWQMDPEMQMHMVESV
 VSLLREAKPLNFEPGKESSSESCKNGSGITADNISLEERQRLGRYGIWLLVGRCTP

TADTPVEVLGRILSMLFHWFHVTAYSYDGFISCLVPHPPPEYARVGGHWETLASRT
 SHLKEGLQRLICLVPEVITSEIWDYVMPHWMEAITNDVAEKELNELKIVLSKILD
 PEMSPGLGDAKTMYNFVAIRFEKTTAKVQQQALHWLQILTKLEILPLVQLFAMF
 GDGVRIMKYGIQHELMREKDAQSLSLAKAPKTPCKESKETKADMANPPRPPVVE
 5 DDSGNTSAISDDEAPNHRHTEFSTDAEHNLTCCILMLDILLKQMELODVEQHMGI
 HTSVCENVSRLIKCMVTAARVGLSSHVCALKVPIEDIIIEEKSSRKSPPESDKEKTR
 DRDVSLSMAPLPIPLGLGGFADP

>/tmp/aaaaainga|GENSCAN_predicted_CDS_2|2553_bp
 10 atggccacgcgagggcggaatgtgattggttcgccatggattgcgcctccatgataat
 cccgctctattggccgcccgcgataaggatcagggtatagccctaattcccgtttcatattcgatggagagagtgcagggtacc
 aagaatgtgggttacatcgatgcgtttcctcctggactcgttcaggacatcgatgatcagctacaggcggaactgatggagc
 tggacgcctcctggtctcgaggcggaaccggcttatctccgctacatgagcaagtgcgtctgcacaggattgcatag
 agcaggactcgcagccaatttggaatgagcgcgatgaaagcatccgttctctatgctgggagctgaatacgcacttgcgagaag
 15 gtatcacacacgctttgggatccgaattggtgattgagaccaatgtgtgcattccaccgctgacctaccaaatgttctgatacgt
 gcacgcaccacaatggagatgtgaatgggatgaggatacgggagaaggagaagggaaccggcggaaggatcgactgggcta
 aggaagggcgctgttgaggggcggaactccgacgaacggaatgtcaggcctgccaatcagtgctcctggctcatcatgatg
 gtgtccagtgactccaacaatccagcgcattgccagctcctggagtgcctgatgactcttaagcacaatgtcgtcaaggacatc
 ctctgctgttggcatacggaaaccgctgttcccgacctcggctgccaagctgctcttactactgcccagcctttaacccaatc
 20 tgttcgatgcgaaagtctactctccaaactaaccaatgacctagtgcccttcacctgccaacgggagcactgtccgaactccggg
 aatgoggaggcagcaaggtgtgtctacgaccacagcattagcatcgatcgcccgattgtccaccgcccctttacctgtgca
 tcgagtgcgccaacgagattcatcgaggacaggaagcctggagtgcggacattctgcatccatgcagcagggtatcgatgg
 tgtcgaaaacaagaactgtcgtccaacgagaagtcgccttctccatctgcttccacggagtgtgccagcttcaatggcaac
 catccgatccgctactgcagccagtccacagtaataggcacaattcccgcgagggtggcgatcacgtggtccatcgagctctg
 25 agccccgctggcagatggatccagatgcagatgcacatggtggagtcgggtgtaagccttctgcgagaggcggaagccacta
 aactttgagcccggaaggagtcctcgtcgtccgagtcgcaaaaagaacggctccggcatcacagctgacaatatttctctggagg
 aacgccagagactgggacgctatgtatctggctactggtgggctgctgtacaccactgcagatactcccgtagaagtctggg
 caggattctgagcatgctctccactggttcatgtaaccgcttactcatacagatggttttatctcctggtgccacatccccgga
 gtatgcccggttgaggagccactgggagacctggcgctcgcaacaagccactgaaagagggtcttcagcggttatatgcctg
 30 gtgccatgatagggtatcacttccgaaattgggactatgtgatccgcactggatggaggccatcaccaacgacgtggccgaga
 aggaactgaacgagctgaagattgtgctcagcaagatccctgatccggaaatgtcgctctgggctttgatccaaaacatgtac
 aactttgtggccattcgattgagaagacaacggcaagggtgcagcagcaggcactccactgggtgcagatcctcaccaagctgg
 agattctcattccactggtccagttgttcgcatgttcggcgatggtgttcgcataatgaatacggcatccagcacgagctgatgcg
 cgagaaggatgccaatctcagtccttggccaaggctcccaagaccccggtgaaagagagcaaggagaccaaaaggatg
 35 gccaatccgccaggcctcctgtgtcaggatgactctgtaatacgtctgccatttcggatgacgaggcgccacgaatcgta
 cacggaattctccaggtgctgagcacaatctcactgttgcatcctcatgctggacatacttgaagcaaatggaactacagga
 cgtggagcagcacatgggcatccatagctgtcgcgagaacgtctccaggctgatcaagtgcattggtcactgcagctcaggt
 ggggtcagtagtcatgtctgcgccttaagggttccatcgaggacatcattgaggaagaaaagtcctcgcgcaaatctccaccg
 aatccgacaaggaaaagacccgtgatcgagatgttccctctcgatggctccactaccattccgctgggacctttaggaggatttg
 40 cagacccttaa

Human Homologue BLASTX with EST: Phosphatidylinositol transfer protein
 (P48739)

45 **Drosophila EST** SD01527 (AI530804), GH18602 (AI387906)

Annotated Drosophila genome genomic segment AE003725

Annotated *Drosophila* genome Complete gene candidate CG5269 – vib PIP transfer protein

- 5 **Human homologue of Complete gene candidate** 1e-90 1346772 P48739
PPI2_HUMAN
PHOSPHATIDYLINOSITOL
TRANSFER PROTEIN BETA
ISOFORM
- 10 **Putative function** phosopolipid transporter involved in lipid metabolism
- Confirmation by RNAi** Slight reduction of G1 and increase in G2/M peaks
 indicating arrest in G2/M

Line ID 418/32
Category Meiotic defects in testis: cytokinesis defects. Dark bands in eyes, dominant.
Reversion ?
5 **Map Position** 69C

Rescue ID G2E

Rescue Sequence

AGCTAAATAACAAACTCATTACTAGTATATTACTGCCGCCGATTTGCAAGCGC
10 GTACCGATCCCGATACCAGGCCAATCGCACTCCCCAGTTGTACGTCATCACTT
AAGTAATAAATCAGCGGCAAATCGCATAAATTGCTATTGATATTCCGCCCGCT
GTGTGTGCGTGTGTATTTGCAAAAGAGTGTGTGTGTGTATGTGCATATGACTC
GTGCGTTTAGCCGACAATTGGAGAAAAAGCATTAGAATCCCAATTGGCTAACT
AAACTAAAGTTGGCTTGGCCAAACATAAAACAAAAAGTGCGGGCGCAGCGATT
15 TGGCAGCGAAACAAAAACACCAAAGTGTTATTGGCAGATATATATGTTAATTA
AATATNAAAAAGTGCGTGCGAA

Genomic hit, Accession No. AC006589

20 **Drosophila EST** SD01527 (AI530804), GH18602 (AI387906)

Rest of results same as line 1104/16

Example 4 (Category 1)

Line ID 1285/1
Category Meiotic defects in testis: cytokinesis defects
Reversion ?
Map Position 85D1-5

Rescue ID D8E

Rescue Sequence

10 GTTCGCAAAAAATATATCTCACCGTGAGTGCAGAAAGAGAAAAAGAGAAGCGG
 AGAGGTGGAGAGCAAGTGGACATGAATCGTCGAGAGTCAGAGAGAGAGAGG
 TGGAGAGGGTGAGCAGCTGTTGTCTGACAATAACATAATCAGCAACAATTTAT
 GCTGTTTAAAAAGAGCAAGAGAAACGCTAATGAAGGGGAACACGGGCAGGGT
 CAGGGGTTGGTGGATCCCCTACATATCTCTCTCTTTACCGCCCCCGCTCTGGC
 15 ACCCTCTCTGTCTGCTCTCCATTAGCCGCACACGTGCAAGCTTAGCATTTCTATC
 TGTCTGTCTCTGTTTGTGTTTGTGTTGCTAAGCCGAATTCT

Genomic hit, Accession No. CSC:AC014256

Associated ORF

Genscan ORF1 predicted sequences

>/tmp/aaaaakfaa|GENSCAN_predicted_peptide_1|702_aa

MIQRCVLLWVFCDFLGLLFLKRKRNAHTPPPPQFTTYRHLLCYCFRNGEIM
 ANICLSRLSVLEEVLLLRVPCAFYFVDYYYVPCLLSVLSESFLYHDQLKVFNRK
 25 QQHQQQQQQQQQQLYQQHQQQQQQHYGPPPPYFQQLHQHQHQQQQQQQQQQ
 HQQHMKFLGGNDDRNGRGVGVGTDAIVGSRGGVSQDAADAAGAAAAAAGV
 YVFQQRPSGGVGVGVGGVGGVGPVGAVGSTLHEAAAAEYAAHFAQKQQQT
 RWACGDDGHGIDNPDKWKYNPPMNPANAAPGGPPGNGSNGGPGAIGTIGMGSG
 LGGGGGGGAGGGNNGSGTNGGLHHQSMAAAAANMAAMQQAALAKHNHMI
 30 SAAAAVAAQQHQHHPHQHPQQQQQQQQAQNQGHPHLMGGGNGLGNGNG
 LGIHPGQQQQQQQQQQQQQHPGQYNANLLNHAAALGHMSSYAQSGGSMYDH
 HGGAMHPGMNGGMPKQQPLGPPGAGGPQDYVYMGQTTPVPMGAAMMPPQNG
 YMNSSAVAAANRNAAITSTAKKLWEKSDGKGVSSSTPGGPLHPLQIPGIGDPSS
 VWKDHTWSTQGENILVPPPSRAYAHGGASDTSNSGNAGILSPRDSTCAKVEYVF
 35 SGSPTNKDSSLSGLEPHLRNLKFDDNDKSRDDKEKANSPFDTNGLKKDDQVTNSN
 GVVNGIDDDKGFK

>/tmp/aaaaakfaa|GENSCAN_predicted_CDS_1|2109_bp

atgattcagcgctcggtgttcttctatgtagtctgcttctgcgactgttcttggggtcctgttctcaaacgtaaacgcaacgca
 40 cactactccccccccccccgcaattcaccactatcgcatctacttgttattgtttcgtaatgggaaatcatggctaattttgc
 cttagtctgttctttagaagaattgttttctttacgctgtgcttattttgttattattattgtccctgtctgtctgt
 ctgtgttctcggaattcttcttaccatgaccagctcaagttttaatcgcacaaaacagcaacaccaacagcagcagcagcagca
 gcagcagcaactctatcagcaacatcaacagcagcagcagcaacattacggtccaccaccgacctactttcaacagctacacca
 gcaacaccaacagcagcagcaacaacagcagcagcagcaacaccagcaacacatgaagttttgggtggtaacgatgatcgca
 45 atggcgcgaggcgctggcggtggcacggatgcatgtaggatctcaggtggcgctctcaggtatgccgccgatgcagctg
 gtgccgccgagccgccgctggctatgtctccagcagcgtccatcgctgtgggtggcgctggcggtggcggtggcggtg

ggtggcgggtgtgccaggggtcggagccgtaggctcgcacttcacgaggccgcccgcggagtagccgcccactttgcc
 agaagcaacagcagacccgatgggcgtgcggcgacgacggccatgggatcgataacccggacaaatggaagtacaatccgc
 cgatgaatccggccaatgcgcctcctggcggtccaccgggaaatggcagtaatggtggcccgccgattggaaccattggc
 atgggcagcgggattgggtgtgtggcgcgggagctggcgcggaataatggcggtctgtacgaatggcggtctgc
 5 atcatcaatcgatggcgtcagctgcgaatatggcagccatgcaacaggcgcgcggttgccaagcacaatcacatgatat
 cacaggcagcagccgagttgcagctcagcaacaacatcagcatccacaccagcagcatcccagcagcagcagcaacagca
 gcaggcgagcaaacaggggcatccacatcaccttatggcggtggaatggactgggcaacggcaatggattgggcatacaa
 catcccgccagcaacagcagcagcagcagcaacaacagcagcagcaacatcccgccagtacaacgcgaatctgcttaacc
 atgctggctgccttgggtcacatgtcatcttatgcccaatcgggtggcagcatgtacgacctcatggtggagccatgcacccggg
 10 aatgaacggcggtcatgcccagcaacagccattgggtccaccggagccggaggacccaggactatgtctacatgggtggc
 cagaccactgtgccatgggagccgaatgatgccgccacagaatcaatatgaacagctctgtgttcagctgccaatcgga
 atgcagcgattaccacatccactgccaagaaattgtgggagaaatccgatggcaaggcggtatcctcagcactcccggtggac
 cgttgcatccctgcagatcccggtcatcggtgatccctcctcctgtgtgaaggatcacacctgtgtccacacagggcgagaatat
 attggtgccgccccctgcgagcctacgccatggaggcgccctccgatacttcaaacagcggaatgcgggcatactgagtc
 15 ccgcatcgcacttgcgccaagtgttgaatatgtttcagtggtcgcgccaccaacaagatagctcgctttccggattggaacc
 gcatttgcgggaatctaaagttgacgacaacgataagtacgcgacgataaggagaaagcaactctccgttgacacaacgggt
 tgaagaaagacgatcaggtcacaaactcaatggtgtgtcaacggcattgacgatgacaagggttcaagtga

***Drosophila* Gene Hit** TBLASTN of ORF1: pumilio protein (L07943)
 20 **Human Homologue** TBLASTX with pumilio: Soares fetal heart NbHH19W Homo
 sapiens cDNA clone (W77820)

Annotated *Drosophila* genome genomic segment AE003681
Annotated *Drosophila* genome Complete gene candidate CG9755 – pumilio RNA

25 **Human homologue of Complete gene candidate** 1e-154 1944416
 dbj|BAA19665| (D87078)
 similar to D.melanogaster
 pumilio protein (S22026)

30 **Putative function** Putative RNA binding protein which is localised to the cytoplasm.
 Wild-type allele of pum involved in development of the abdomen
 (embryos) and of the imaginal discs (larvae or pupae), perhaps
 35 having a function in signal transport.

Confirmation by RNAi Only wild type profiles observed

Example 5 (Category 1)

	Line ID	1389/1
	Category	Meiotic defects in testis: segregation defect, cytokinesis defect (Ck-09/32)
5	Reversion	NR
	Map Position	93B4-8
	Rescue ID	2C9P
10	Rescue Sequence 1	
	GTTCCGGGGTGTGTGCGTGCTTGCGAGTGTGCCTGTGTGTGTGTAGGAAAGGAG CAAGAAGCAGCAGCAGCGGCAGCAGTAGAAATAGCAAAAGGAGGCAGCAAC AACAATAAGCTAGAGAAACCGCCAGCAGCAGCCCCCTAATAAAGAGCAGAGA AAAAAATGAGTTCAAGTTGTGAAAGGTGTGTGCCGTTACACTACAACTACAA 15 CACCACCATCAGCGGCAGCAAAGAAATACAACAACAAATACGGCAATCTCCA GACAACGCGAATGTCGAAATTGTGTATACAATTTATTAAGAAAGCAAGAGCA GCAACAACAATGACCAGCTGCAGTTCATCAGCGGTGTCCTCCTGAATGCCGCT GTCGTGCTTGGTGTCTGCCACCGGCGGTTCTCAATAATAAGGGCAGGAGGAG CTGCTTAGGTGCACACAATGTAGTTTGGCTTGGTGAATGCTTCTCTTTTGTG 20 CTGCTGGCGCATACGTTCTCTCTCCCTCATGATCTCAGTTGTCTGCATCGA TGAGCCGCCACCAACGGTGGCTTCTTCTGCTCCTCTTTGGCAACGGACTGCTG CAGTCTTGCCAGAATTTTCTCTAAATACTGAGCTTCAACTTGGTCTGCTTGGT AATGGTATACCATAAGCCATGGACTTGATGCCCCTACAAAGCTCTGTGATTG AAATGGGATGCA	
25	Rescue ID	2C9E
	Rescue Sequence 2	
	CCCCGAACGCACTTTATATATATAAATATATATATTATTTTCTTCACTTATTTT CGTTTCGGCCGCGACAGCGAATATGCAATTTTCTCTCAATTGATTTTTTACA 30 CACTCGCACTCCTTTTACATGCGTGCAGTTTATGTTGCTATTGCTGCTACTGC TGCTGTTGTTGTTATTGTTGTTCTGGCTGCCGCTGCAGTGCAACTTGTAACACT TTCACATTTATGACATAATGCACTGGCCATATTTTGGCTTGGCTCTCCGTTTGT GCAACTGCATGTTCCCAGTGCTTTTTTAATATTTATGCTGCAGTGCGTGCAAAT TCGAACGCGAGACGATCCGCTTTTCGCTGCATCTATGCGCTGAAGATGTGCTG 35 CAGTCGATGGGCTCGTCGATAGTGGGAAGGCTCGGTGCCGGCACTATCGATT CCAACACCATACGATAATATCGGCTAAAGTTATCAATATCGAAGTTTACTATA TTTCGGGTTTTTACGTTTTTAAATCTACCTTATCAACATTTTGNAGAAGTAAA AAGTAGTTCTCTTATGGATGCATC	
40	<i>Drosophila</i> EST	several including LD10379 (AA816796)
	Annotated <i>Drosophila</i> genome genomic segment	
	Annotated <i>Drosophila</i> genome Complete gene candidate	
	AE003733	CG3421 - novel protein with weak homology to myosin
45		

5	Human homologue of Complete gene candidate	Ensembl predicted Gene:ENSG00000071333 Clone:AC022505 Contig:AC022505.00011 5.60E-37 (predicted protein with Core domain in kinesin and myosin motors ENSG00000087179)
10	Putative function	Possible novel motor protein involved in cytoskeleton organization
	Confirmation by RNAi	Marked reduction of G1 and G2/M peaks indicating fewer cycling cells

Example 6 (Category 1)

Line ID 293/9
Category Mitotic defects in brain: cytokinesis defect
 5 (no overcondensation of diploids, very high polyploidy)
Reversion NR
Map Position 66B

Rescue ID 2G5E
 10 **Rescue Sequence**
 GTACAAACGAATTATTTGTCTCCTTGTGCGTTCGTTTTATTGTGTTTCGAGTTCT
 GTTGGTGTGTGTTTTTGTGTATGTTCCACGAGTTGTTTCGCATTAATAAAATTAAC
 TGCAGAAGATCCATGGAAATGGAGACCATTGAAGAGCAATCGAAGTGCGGTG
 AGTACTGAAAGAGGGCGCGGGGCGTGGCAGCTCCAAATGGCCGGCGAATTTA
 15 TCATTTTCAATGTCGTCCAAAGGGGTGGGTACGGGGTAAAACCACATTCGG
 GGCCAAAAGATCCTCATAAAAAATGTCGCTGCCAGCAAATGCAAAAAATAAA
 ATAAAATAAGAACGACTATAAGTACATCTTTGTGTGTATTGTGTGACTAAAA
 AAGCAACGGCATCGTGTGCGCANATATTTAATCTTTNTTTCTGAATTTATTTCTG
 GTGTACAAAATATTTATCGCATAAATGCGAAATGCCTCCCTCTCTTCATCATCG
 20 T

Genomic hit, Accession No. AC008303

Associated ORF
 25 Genscan ORF1 predicted sequences >20:53:38|GENSCAN_predicted_peptide_3|261_aa
 MMDNDLALLNNGGPQSGAETVYGTEDNNMVMSEKCRIFPATQRTGFVGATFSG
 VLLLDL GALQHCDVIRIDVNIATLEQIKRERQEELAAERERIRAQIAADRAEQAQRF
 NTPDISSTNSVAATAASNVTDDASVSSVDETRLQIRLPGGIQRKTSFPAGEVLAT
 VRVYVRNEMLAASDVRDFTLATSYPREFQTEDEVKTLNELNLVPNAVVLVLT
 30 EQVNPADDQTAKRSASTKRTKTHRHKRQLMADEPQSDHYKN

 >20:53:38|GENSCAN_predicted_CDS_3|786_bp
 atgatggacaacgatgatgcactgtcaacaatggaggaccacagtcggagctgaaactgtctacggtaccgaggacaacaac
 atggatcatgtcgagaagtgcgcataatcccgcgactcagcgactgattgtggcgcgacgttttcgggagtgctgcttct
 35 gatcttggtgccctccagcattgtgatgtatccggattgatgtaacattgcaacgctggaacagattaagcgtgagcgtcaggag
 gagctggcgccaggagcgcattcgtgccaaattgcagccgatcggcgagcaggcacacgttttaatacggcgacat
 tagcagcagcacaattcgggtggcgccaccgctgcctccaacgtgatcacaacagacgcctcggtagtggtggagcagaga
 cgaggctgcagatccgactaccggtggcattcagcgcaccaaatccttccagccggcgaggtgctggctaccgttcgtgtcta
 cgtgcgaaacgagatgtggcgcgagcgtatcgcgactttaccctggctaccagttaccacgaaggaggtccaaacgg
 40 aggcagaggtcaagaccctgaacgagcctaactagtgcctaatcggtggttctggtgctgaccaaggagcaagtgaatccag
 ctgatgaccagacagcaaacgatcagcaagcacaacacacacagacacacagcggcaattgatggcagacga
 gccacaatctgaccattataaaaactga

45 **Drosophila Gene Hit** rescue sequence: pebble (rho1 GTPase exchange factor)
 (AF136492)
Human Homologue BLASTX with pebble: KIAA0337 (AB002335)

	<i>Drosophila</i> EST	SD09146 (AI542703), SD01796 (AI530981)
	Annotated <i>Drosophila</i> genome genomic segment	AE003557
5	Annotated <i>Drosophila</i> genome Complete gene candidate	CG8114 - pbl pebble rho1 GTPase exchange factor
10	Human homologue of Complete gene candidate	2224615 dbj BAA20795 (AB002335) KIAA0337 [Homo sapiens (3e-21) also mouse homologue 3e-95 42359 transforming protein (ect2) - mouse >gi 293332 (L11316) ect2 [Mus musculus]
15	Putative function	A guanyl-nucleotide exchange factor involved in signal transduction which is localised to the mitotic anaphase. pbl is required for the formation of the contractile ring and the initiation of cytokinesis in <i>Drosophila</i>
20	Confirmation by RNAi	Slightly reduced G1 and G2/M peaks indicating fewer cycling cells

- Line ID** 542/3
Category Mitotic defects in brain: cytokinesis defect
(very high polyploidy)
Reversion NR
5 **Map Position** 66A
Rescue ID 2A1E
Rescue Sequence
GTCCAGTTAATGAAAGTAAACGAATCGAGTACAAACGAATTATTTGTCTCCTT
GTGCGTTCGTTTTATTGTGTTTCGAGTTCTGTTGGTGTGTGTTTTTGTGTATGTT
10 CCACGAGTTGTTTCGCATTAAAAAATTAAGTGCAGAAGATCCATGGAAATGGA
GACCATTGAAGAGCAATCGAAGTGCGGTGAGTACTGAAAGAGGGCGCGGGGC
GTGGCAGCTCCAAATGGCCGGCGAATTTATCATTTTTCAATGTCGCCCCAAAGG
GGTTGGGTACGGGGTAAAACACATTCCGGGGCCAAAAGATCCTCATAAAAAA
TGTCGCTGCCAGCAAATGCAAAAAATAAAATGAAATAAGAACGACTATAAGT
15 ACATCTTAGTGTGTATTTGTGTGACTAAAAAAGCAACGGCATCGTGTGCGCANA
TATTTTAATCTTTTTTCTGAATTTATTTTCGNGTANAAAAATTTTATCGCATA
AATGCGAAATGCCTCCCTCTCTTCATCATCGNTTCCCCTNACTCTCCCTCTCTT
CGCCCGACACTGTACCGACGCAAGAAGAAC
- 20 **Genomic hit, Accession No.** CSC:AC018042
Drosophila EST SD09146 (AI542703), SD01796 (AI530981)

rest of results same as line 293/9

Example 7 (Category 1)

5	Line ID Category	229/30 Mitotic defects in brain: cytokinesis defect. Meiotic defects in testis: cytokinesis defects (Mitotic higher level of condensation, polyploidy, Meiotic:
	Ck05/07)	
	Reversion	?
10	Map Position	91F
	Rescue ID	A7E
	Rescue Sequence	
15		TCTTGGCCAAACAACGCGAGCAGCTGATGTCGCATGGTGGGAAAATGAGGGT GGCGCGAGTGGAAGTTGCCATATCGCTGCGATCACAAGCAGCAAATATGGAA GATTAAGCGGAAAACGAAAGACAAAATAATTACAATCAAACAACCGAATTAT AAAAAGAAAATGGTTTGTCTCCGAGTTCGTTTAAATATGCTTATCTACGTATC AATTAAAAAAACCGTAGAAAGAAATTCACGATTCACCCTAATCTAGCTAAGA CACCAACCAAAAATTTCCGATTTACTTTCAGTTGAAGTTGTTGTTACACACTTT TCTTGTGCGATGTTTTGAAGCGCCCATTTGAAATTGATCATTGAAATGTTTTTCCA 20 AATTACCCACATCCATTACAATAAAATTTAAATTGCTTATTATTTGATTTTTACT TGGGAAAATCCCCTTGCCAAATTGGAATTACAATTCCAGCTTGGAATCCGTCA AACTTTACAACATAAACTTATTGTTCTTTTCCGGACAATGCTTCCA
25	Annotated <i>Drosophila</i> genome genomic segment Annotated <i>Drosophila</i> genome Complete gene candidate	AE003686 CG6284 - novel protein possible sir2 family of transcriptional regulators/telomeric silencing
30	Human homologue of Complete gene candidate	gi7706710 0268A424791DE5BF [ref]NP_057623.1 sir2-related protein type 6 [Homo sapiens] (1.10E-74)
35		
	Putative function	Putative transcriptional regulator
40	Confirmation by RNAi cycling cells	Complete loss of G1 and G2/M peaks indicating fewer

Line ID 1104/16
Category Mitotic defects in brain, Cytokinesis defect (no overcondensation of diploids, high polyploidy)
Reversion R
5 **Map Position** 92A

Rescue ID B5E
Rescue Sequence
10 GTCCGGAGCGGAGCTAAAGTTCGATGTTTCGTGCAAAACACTTCGATTCCGATA
GGCGGATGCTATCGATTTTCGGCGATGCCCGTTGGTCACACTTGGTGGTGGGCG
CTGCCCCGTCGCCGACTATCGATAGCACAAGCGGGTTATTTAGGTGTGCGCAGC
TTGTAAGGGTGACTCATGCTGTAAATATTATAAAAAAGTTAATGAATATAA
TATAGTTATAATAAAATTATATATAAATCTATAAGATCAAAGATCATCAGTTA
TCATTTATCATTTGATTATATGAAAAACAAGAACAGAAACAAGATTTAATAGG
15 TTTTGTGAAATGTGAAAATGTGGGTTACCCCAATTCTTATTCGAAATTAAATAA
CCTAAAGAACAGTTATACACAGATAGGTAATTTGCACATAAGCCAAATTTGT
CTAGAATTCCGCGGAATTAATTCTTGAAGACGAAAGGGCCTCCGTGATACGCC
TATTTTATAGGTTAATGTCATGATAATAATGGTTTCTTA

20 **Rescue ID** B5P
Rescue Sequence
CTCCGGACACGCAGTAGCTAAATAACAAACTCATTACTAGTATATTACTGCCG
CCGATTTGCAAGCGCGTACCGATCCCGATACCAGGCCAATCGCACTCCCCAGT
TGTACGTCATCACTTAAGTAATAAATCAGCGGCAAATCGCATAAATTGCTATT
25 GATATTCCGCCCGCTGTGTGTGCGTGTGTATTTGCAAGAGAGTGTGTGTGTGT
GTGTGCATATGACTCGTGCGTTTAGCCGACAATTGGAGAAAAAGCATTACCAA
TCCCAATTGGCTAACTAACTAAAGTTGGCTTGGCCAAACATAAACAAGT
GCGGGCGCAGCGATTTGGCAGCGAAACATATACACCAAAGCGCTATTGGCAG
ATATATATGTAGATTAAATATAGAAAGTTCGTGCGAAGGTTAAGAGTCGAGT
30 GCAAGTGCATTTATATTTGGAAATAATAAATGCTACAAT

other results same as 229/30

Example 8 (Category 1)

Line ID 343/5
Category Mitotic defects in brain: cytokinesis defect
 5 (very high polyploidy, chromosomes entangled?)
Reversion NR
Map Position 75B

Rescue ID C6E
 10 **Rescue Sequence**
 GGTTTCGAGTTCGTTTCGGCTCTCCGTTTCGGCTCTCTCTCGCCATCCC
 GCTGCCGCACACATTGGCCTCTCTCTCGCAGCTCCACATTGGAAGGTGGCTGA
 CCGAAATGTGGGTCACGACAATGGCGGGGTTTCGTTGAACTGAACCACCGCCG
 CAGTCGCTGCCGTGCTCGCTGCTCTGCCTCTGCTGACGTCGTTAACGTTTTGGG
 15 GCTTTCGGTTACGTAGCTCGTGTGCGAGCGAGAGGGGCTACTAGAGGGACTGC
 GACACACAAGTTGTGTGCATTTTTTGGCCCCAAAAAATCACAATGGGCACAAA
 ATATTATTTAATACATCACATAATTGTTTAAATCATCTGGCTGGAAAGTGTGCGAG
 TTCATCGAACTGCCAGCGATTGACAAATTGCGATTTTCAATGCGGCAAAAATA
 TTTACTCAAGCAAATTGTTTGCACCTCGTTAATTAGGCGGGGAGTGCCGCCAA
 20 ATTGGGTCATATTGCAGAAAGTATCCAAGAAGTTGGAGAAACAAGCTGCTTAA
 ACATTAATTAACACACACCTAAATGGATACATTTGCTACAAACAATTATAAAT
 GTTACCCTTATATTAATTTTCAAATTTCTAAATAATCAA

Genomic hit, Accession No. CSC:AC015427
 25
Associated ORF
 Genscan ORF1 predicted sequences
 MVCAMQEVAAVQHQQQQQQLQLPQQQQQQQQTQQQHATTIVLLTGNGGGNL
 HIVATPQQHQPMHQLHHQHQQHQQHQQQAKSQQLKQQHSALVKLLESAPIKQQ
 30 QQTPKQIVYLQQQQQQPQRKRLKNEAAIVQQQQQTPATLVKTTTTSNSNSNNTQT
 TNSISQQQQQHQIVLQHQQPAAAATPKPCADLSAKNDSSESGIDEDSPNSDEDCPN
 ANPAGTSLEDSSYEYQCPWKKIRYARELKQRELEQQQTGGSSNAQQQVEAKPA
 AIPTSNIKQLHCDSPFSAQTHKEIANLLRQQSQQQQVVATQQQQQQQQQHQQHQQ
 QRRDSSDSNCSLMSNSSNSAGNCCTCNAGDDQQLEEMDEAHDSGCDDELCEQH
 35 HQRLDSSQLNYLCQKFDEKLDTALSNSSANTGRNTPAVTANEDADGFFRRSIQKQ
 IQYRPCTKNQQCSILRINRNRQYCRLKKCIAVGMSRDVLRLEQPKAGAKNKSCE
 PSKNSTVNQINSKLELGNSNEMK

 >21:55:09|GENSCAN_predicted_CDS_1|1533_bp
 40 atggtttgtgcaatgcaagaggtgtgctgccgtgcagcatcagcagcagcaacagcaactccagttgccccagcagcaacagcag
 cagcagcagacaacacagcagcaacatgcaacaactatagtgtgctgacgggcaatggcgcggtaatctgcacattgtcgcc
 acaccgcaacagcatcagccgatgcacagctccaccatcagcatcagcatcagcatcagcaccagcagcaggccaagagcc
 aacagctgaagcaacaacactcggcgctggtcaagttgctggagtcggcgcccatcaagcagcaacagcagcagcccaagca
 aattgtttacctgcagcagcagcagcagcaaccgcaacgcaaaaagactgaaaaacgaagcagcaatctacaacagcaacaac
 45 aaacacctgcaacactagtaaagacaacaaccaccagcaacagcaacagcaacaacacccagacaacaatagtattagtcag
 cagcaacagcagcatcagattgtgttgagcaccagcagccagccggcgagcagcaacaccaagccatgtgccgatctgagcg

5 ccaaaaatgacagcgagtcgggcatcgacgaggactcccccaacagcgatgaggattgccccaatgccaacccggcgggcac
 atcgctcgaggacagcagctacgagcagtatcagtgccctggaagaagatacgctatgcgcgtgagctcaagcagcgcgagt
 tggagcagcagcagaccaccggaggcagcaacgcgcagcagcaagtcgaggcgaagccagctgcaataccaccagcaac
 atcaagcagctgcaactgtgatagtcccttttcggcgcagaccacaaagaaatcgcaatctcctgcgccaacagtccagcaac
 10 aacaggttggccacgcagcagcagcagcaacagcagcagcagcaccagcaccagcaacaacgaaggatagctccgaca
 gcaactgctcgtgatgagcaactcgagcaactccagtcgggcaattgtgcacctgcaacgctggcgacgaccagcagctgg
 aggagatggacgagggcccacgattcgggctgcgacgatgaacttgcgagcagcatcaccagcgactggactcctccaaactg
 aattacctgtccagaagttcgatgagaactggacacggcgctgagcaacagcagcgccaacacggggagggaacacgccag
 ctgtaacagtaacgaagatgccgatggattcttcgcccgtccatccagcaaaagatccagtatgcccgtgcaccaagaatca
 15 gcagtgcagcattctgcgcatcaatcgcaatcgttgtaaatattgccgctgaaaaagtgcattgccgtgggcatgagtcgcatgt
 tctgcgcctagagcaacctaagctggtgccaataaagtcattgtgaaccgagcaaaaattcgaccgtcaaccaataaacagc
 aaactcgaactcggaacagcaatgaaatgaaatga

15 **Drosophila Gene Hit** TBLASTN with ORF1: ecdysone-inducible gene E75B (X51549)
 and nuclear receptor superfamily protein (U01087) BLASTN with
 genomic sequence matches ecdysone inducible gene

20 **Annotated Drosophila genome genomic segment** AE003522
Annotated Drosophila genome Complete gene candidate CG8127 Eip75B ecdysone-
 inducible gene E75B nuclear
 receptor NR1D3

25 **Human homologue of Complete gene candidate** ORPHAN NUCLEAR
 RECEPTOR NR1D1 (V-
 ERBA RELATED PROTEIN
 EAR-1) (REV-ERBA-
 ALPHA) Q15304 (9.40E-74)

30 **Putative function** Ligand-dependent nuclear receptor, putative transcription factor

Confirmation by RNAi Slightly reduced G1 and G2/M indicating fewer cycling
 cells

35

	Line ID	448/23
	Category	Mitotic defects in brain: cytokinesis defect (very high polyploidy)
	Reversion	NR
5	Map Position	75B
	Rescue ID	2G4E
	Rescue Sequence	
10		GCTGGTGGACGCTGCTTTCATTTCGCAAATTGCTCGTCGTTGGCAGCGGTTGTGC AGAGCAAGAAAAGCGCGCGAAAAACCAAGCAAAAAATTAATACAGCTGGAT CAAGCGAAAAGAGATAGAGAGCAGAGTCAACAGCAACAAATGTTCAATAGCA AATGATATCGCATATTTTTGTTGGTGCCAGTGAAGTGAGATCAAAGTGAAGTG TGCAATGTTCTTATTAGCAAATCGTAGAGCAACCAACAATCGAGAGTTCAAG TGTCATTTCTGAAGCCAAAAAGCAAAATCTCTAATTCAAATATGGTTTGTGCAA 15 TGCAAGAGGTTGCTGCTGTGCAGCATCAGCAGCAGCAACAGCAACTCCAGTT GCCCCAGCAGCAACAGCAGCAGCAGCAGACAACACAGCAGCAACATGCAAC AACGATAGTGCTGCTGACGGGCAATGGCGGCGGTAATCTGCACATTGTCGCCA CACCGCAACAGCATCAGCCGATGCATCAGCTCCACCATCAGCATCAGCATCAG CATCAGCACCAGCAGCAGGCCAAGAGCCAACAGCTGAAGCAACAACACTCGG 20 CGCTGGTCAAGTTGCTGGAGTCGGCGCCCATCAAGCAGCAACAGCAGACGCC CAAGCAAATTGGTTACCTGCAGCAGCAGCAGCAGCAACCGCAACGCAAAAGA CTGAAAAACGAAGCACAATCGTACAACAGCAACAACAAACACCTGCAACAC
	Genomic hit, Accession No.	CSC:AC015427
25	Drosophila EST	GM03519 (A801874)

Other results same as line 343/5

Example 9 (Category 1)

5	Line ID	36/1
	Category	Meiotic defects in testis: cytokinesis defects (Ck-04/06) `
	Reversion	R
	Map Position	79C
	Rescue ID	A8B
10	Rescue Sequence	GAGTAAAGTAAACTACAGAGAAAAACGCTTTACGGCGAGAGAACGCTTTAA TTATACTTAATTTGTTGTTAATCAAACGCACAGAGCACACAACACAGAAACAC AAAACACCGCTTGGGAAAAATCTGTAGGTAGANGAAAGGAGCTCACGTTTTT CTGGTGCAGATCGAAATCGGTATCGGGTTTATTTCGCTTTGCCGGATTGTTACTT 15 CACGTTTGTTAATTGCGTTTCTTTGTTTCTTATTCTCCTGCGCACACTTTGATTT GCGTTTGCAACTCGCAATTCGCAATTGGCATTGCTATGCGACAACGCGGTT ATTTCCGGTCCGTTTACTTTTCCAATGGCTTGCCTACACACCGCCAAACTTTGG CTTGAACCTGGGATATTGGTTGCCCGAATTTCTTGANAAATTTTCCTT
20	Genomic hit, Accession No.	CSC:AC013886
	Associated ORF	Genscan partial ORF1: >18:33:59 GENSCAN_predicted_peptide_1 99_aa CICFALLGLLIRKLMVFGSTSRKAQSLESRRAKNTSQKIGNQYPKFSQVCGKPS 25 KSNDNRNNGSCRANANCELRVANANQSVRRRIRNKETQLTNVK
30	>18:33:59 GENSCAN_predicted_CDS_1 300_bp tgtatctgcttcgccctgcttgggctactcattcgcgaaaaattaatgggtggtgttcggttctacgtcgcgcaaggcacagtcctctaga gtctcgagagctaagaatacatctcagaaaatcgcgcaaccaatcccaagttcgcgcaagccatcgaaaagt aacgaccgaaataacggcagttgtcgcatagcaaatgccaattgcgaattgcgagttgcaaacgcaaatcaaatgtgctgcagg agaataaagaacaaagaaacgcaattaacaacgtgaagtaa	
	Drosophila Gene Hit	rescue sequence and TBLASTN with ORF1: nucleic acid binding protein (mub) (X99340)
35	Human Homologue	BLASTX with nucleic acid binding protein: poly(rC)-binding protein 2 (hnRNP-E1) (S42471)
	Drosophila EST	several including LD32520 (AA951799 BLASTN matches nucleic acid binding protein (X99340)
40	Annotated Drosophila genome genomic segment	AE003596
	Annotated Drosophila genome Complete gene candidate	CG7437 - mub mushroom bodies RNA binding protein
45	Human homologue of Complete gene candidate	4826886 ref NP_005007.1 pPCBP2 poly(rC)-binding protein 2

79

>gi|542853|pir||S42471 (4e-75)

5 **Putative function** A putative RNA-binding protein specifically expressed in the CNS of *Drosophila melanogaster*

10 **Confirmation by RNAi** Only wild type profiles observed

Line ID 472/22
Category Female sterile
 (anaphase bridges, lagging chromosomes)
Reversion ?
 5 **Map Position** nd
Rescue ID sau 5'spl

Rescue Sequence

10 GCACGATCNCTAAAGTCTNGCANAGCTAAAAATACATCTNAGAAAATCGGCA
 ACCAATATCCCAAGTTCAGCCAAGTTTGCGGTGTGTAGGCAAGCCATCGAAAA
 GTAACGACCGAAATAACGGCAGTTGTGCGCATAGCAAATGCCAATTGCGAATT
 GCGAGTTGCAAACGCAAATCAAAGTGTGCGCAGGAGAATAAGAAACAAAGA
 AACGCAATTAACAAACGTGAAGTAACAATCCGGCAAAGCGAATAAACCCGAT
 15 ACCGATTTTCGATCGGTGCGGGCCTCTTCGNTATTACGCCAGNTGGCGAAAGGG
 GGATGTGCTGCAAGGCGATTAAAGTTGGGTAAACGCCAGGGTTTTCCAGTCACG
 ACGTTGTAACGACGGCC
 ANTGCCAAGCTCTGCTGCTCTAAACGACGCATTTTCGTA CTCAAAGTACGAAT
 TTTTCCCTCAAGCTCTTATTTTCATTAAACAATGAACAGGACCTAACGCCNGT
 20 AAC

Rescue ID Sau 5'splac

Rescue sequence

25 GTTGTGATCNTCTTGGTNAATCNNNTTGGAAATTCCCCTAANGCTTCCGACAA
 TGACCCNGNCNTACNNAGCAAANAATNGNAGNACNNGCNGNTGGNCGTANT
 ANCAANAACAGGCCCGCACCGATCGAAATNGGNATCGGNTTTATTCGCTTTGC
 CGGATTGTTACTTCACGTTNGTTAATTGCGTTTCTTTGTTTCTTATTCTCCTGCG
 CACACTTTGATTGCGTTTGCAACTCGCAATTGCAATTGGCATTGCTATGCGA
 30 CAACTGCCGTTATTTGCGTCGTTACTTTTCGATGGCTTGCCTACACACCGCAA
 CTTGGCTGAACCTGGGATATTGGTTGCCGATTTTCTGAGATGTATTCTTAGCTC
 TCGGAGACTCTAGAGACTGTGC

Other results same as line 36/1

35

Example 10 (Category 1)

	Line ID	459/2
5	Category	Mitotic defects in brain: cytokinesis defect. Meiotic defects in testis: cytokinesis defects: (mitotic: high polyploidy, no diploids, higher mitotic index, meiotic: Ck-01/05)
	Reversion	NR
10	Map Position	66B1-6
	Rescue ID	2D5P
	Rescue Sequence	GCTCCGTTTCGAAAAGTTGAGAGAGACTTGAAACATATGTTTCGGCGTTGCTAGAG CTGGTCGGCTACCGATAGAAACATCGATAGGTCCGATGTTTTTTACTCGTATAT 15 TGATTCANAGTTTGGCTATCGATGTTTTTAGAGTGCCCGCACATTATCTATTTT CATCTCTATTTTCGTTGGTATTTTTTGTATTTTATGACATTTTCGACTGCAAAAGC AGGATGGCAACGCCAGATTGCCGCGAAAAGTACGTTATTTTTAAATTGGCGCAT TGAATATGAAAAATTGCAGGCACATACAGTTTCTAATAAATAATAGCAATAAT TATTATTTAGCTTGTATCATAACGAAGTGCACATTACAGCTACGCATCTGAAAT 20 AATAATTTTAATATATCGTCTTTTCTCCCATCGATAGAGTTCGCGCCTATCGA TATATCGTTGATCACCAAATAAATAAACTAAATAACGCCGCAATGGAACAC GCGACGAGTGAATTGAGGGAATTTATCTCAGATCTTGTAATTCGACACCACGT TGCAATGGTAACATCAATCCGGATCACATCACAATGCTGGAAGGCACCCAGA TCCAGAACAG 25
	Annotated <i>Drosophila</i> genome genomic segment	AE003557
	Annotated <i>Drosophila</i> genome Complete gene candidate	CG8038 - novel gene ribonuclease P homology 30 CG7892 nmo - protein serine/threonine kinase involved in eye morphogenesis
	Human homologue of Complete gene candidate	CG8038- 5e-24 4309676 gb AAD00893 (AF001176) ribonuclease P protein subunit p29 [Homo sapiens/
40		CG7892- protein kinase mitogen-activated 7 (MAP kinase)' gi:4506093 and gi7706445 D919050533B3C33A
45		[ref NP_057315.1 nemo-like

kinase [Homo sapiens]
(3.30E-174)

- 5 **Putative function** CG8038: tRNA processing enzyme Ribonuclease P protein subunit
CG7892: a protein serine/threonine kinase involved in cell cycle,
possibly targeted to cytoskeleton

- 10 **Confirmation by RNAi** Both showed a marked increase in G1 peak indicating arrest in
G1

Example 11 (Category 1)

	Line ID	623/8
5	Category	Meiotic defects in testis: cytokinesis defects
	Reversion	?
	Map Position	37E1-3
	Rescue ID	2E2E
10	Rescue Sequence	
		CTACGGGCATTTCGCATGTTTCGAACATCTGGTGTAACAAGTTCTGAGCAGTGT TGCCAACTCTTCAGTTAAACAGTTAAAAATAGCTAAAAAATGTTGACGGTAGC TAAATTATAAAGCTAGAAAAGAAATGATATATGATAAAATAAGTATTTCTGACT CACAGCATTTATTATTAAAGACGGTCAGATGAAGTTACAAAAATCCTAAATTG 15 GCGCGCTGTATCTAAGAATTAATACCAAGAAGTTGTCATCAAAGGTCGAACTC GACGGAAATTCTACTTTGAGTTTTTAAATTTAATAAATATGTATTTAAATTAT GTAAATTTGTTTGTAAAACAAAAATAGTATATAGTATAGTAATAGTAGTTAAG TAGTTTTAAAAATGGCCAGATCAAAGACTTTTGAGATATGATACTAATCAAAA GTCGAATTCGCGGAATTAATTCTTGAAGACGAAAGGCCTCGTGATCGCCTATT 20 TTTATAGGTAATGTCATGATAATAATGGTTTCTTAGACGCAGGTGGACTTTTCG GGGAAATGTGCGCGGAACCCCTATTGTTATTTTCTAAATACATTCAAATATGT ATCCGCTCATGAGACAATAACCCTGATAAATGCTTCAATAATATTGAAAAAGG AAGAGTATGAGTATTCAACATTTCCGGGCGCCTTATTCCTTTTTTGGGCGGCAT TTGCCTTCCTGTTTTTGTACCCAGAACGCTGGTGAAAGAAAAGATCTGAAGA 25 CAGT
	Annotated <i>Drosophila</i> genome genomic segment	AE003662
	Annotated <i>Drosophila</i> genome Complete gene candidate	CG17559 dnt - doughnut protein tyrosine kinase
30	Human homologue of Complete gene candidate	Homo sapiens RYKreceptor tyrosine kinase GDB:21773
	Putative function	growth factor transmembrane receptor protein tyrosine kinase involved in cell growth and maintenance
35	Confirmation by RNAi	Only wild type profiles observed

Example 12 (Category 1)

	Line ID	629/14
	Category	Meiotic defects in testis: cytokinesis defects (Ck-06/09)
5	Reversion	NR
	Map Position	64D
	Rescue ID	2A9X
10	Rescue Sequence 1	
	GACGGGAGGAAGTAAGTGGGAGGAGAGAGTAGTGCCTCTTTTTTACTGGAGA AATGGACAAACTCTGGGAACTGCGAACTGCGAACTAACCGAGGCAAAAATTG AGAAGCGAGCTGAAAGCGGAATTCAAACAACGCAGCGCTGACGGCGACGCCG GCAGAAGCAGCGCCGCACAAGGCATGCGCACAGAGAGTAAGAAAGAGCGCG 15 GCTAATGAATGAATGAACGAGGCGGAATGCGGGAAGAGCGCAGAGAGGCGC AATGACAAAATAGTTGTAGAAAAGCGCCGGCAAGCGGAACTCCACACTCTTT CTCACTCTCTCTTTCCACCCACACCCCTAGTTCACCGGAAAAAGAAAATTTCGTT TGCGGCGGGGGTGTATTTTACCAAAAAGAGAGTGTGTGCAAAACGCTAGA GAGAGAGAGAGAGAGAGAAAGAACTGACGTCTGCTCCGTCGACGCC 20 GCTGCCGCGTCCCAAAGCGCCACCACCCAAAAAACGCGAGAAGAAGCAGA ACAAACACACACAAAAATTCGCACAGTGGAGCAGAAATCAAGC	
	Rescue ID	2A9E
	Rescue Sequence 2	
25	CTCCCGTCGTTTTGAGATCAGCTGCTCTCGCAACAACAACACTATAACTGTA GTTACCGTCTCTTTTGCATCGTTTCGTTTTTCGTTTGTGTCGCCAAGTGATTGTGT GTGTGCGTAAGCTTAAAGCTGACTAACAAAACGAAACAAGAAAAAATATAAA TTATAGGAAAATTGTTAAATTATAACCAGAAAAGAGAGCGGCACCTTACGTGTGT TATTGTGTGCGTGTGCTTTAAAAAGATATAAAAAATAGCAATAGAAAAGTTATTA 30 AAGCGTTGGCAAAAAAGTCCAACGAACAGCGAGAGGAAGCGGAGAACGAAA TAGTTAAAGCCAAAGTCGCTGCCGACGTGCGCACTTGAAAACGTCGCAAAAGTT TGTAACACACCAGTGTGTGTTTCGTGTGTGTTTTTGCCGGCGTGCCAGTGTGCG TGCGCCTAGAAAAGAGTAAAGAAGCAGAAGAAAAGGAAGAAGCCGAAGAAG CAGCAAAAGAAGCCGACAGCAAAAAGTAAATAAAATCAAATGCCCCCTGGCA 35 GAATAATATTAAATTAAGACACATACTCAAATTAATAAC	
	Genomic hit, Accession No. CSC:AC015076	
40	<i>Drosophila</i> EST	LP08767 (AI295205)
	Annotated <i>Drosophila</i> genome genomic segment	
	Annotated <i>Drosophila</i> genome Complete gene candidate	
	AE003567 CG10668 - novel with homology to ssDNA/RNA binding proteins	
45	Human homologue of Complete gene candidate	
	CG10668 - 3e-12 4506449	

ref|NP_002889.1|pRBMS2|
RNA binding motif, single
stranded interacting protein 2
>gi|1082

5

Putative function Possible single stranded DNA/RNA binding protein

Confirmation by RNAi Slightly increased G1 and reduced G2/M indicating G1
10 arrest

Example 13 (Category 1)

5	Line ID	653/12
	Category	Meiotic defects in testis: segregation defects, cytokinesis defect (Ck-07/35)
	Reversion	NR
	Map Position	75B
10	Rescue ID	I5E
	Rescue Sequence	GTAAAAGCTTAGCCCATGGCGTCGACGTCGACTGCGACAGCGACGCTAGCCG AGGCAGTGACTGCGACGTTGGCCACTTTTCGCCTTCGTTTCGCTGTCGTTTTCA GTTGTCTCTCGTTGCTCAAAGCGCGCGGCACGCGAACGCTCTGAAATCCCAAG TTACAACAGCAACATCAAGCAGCAGCAACAACAGTGATTCGCTGGCAAACAA 15 ACAAACAAACCAACATATTTTTGTGTATCAATTGTCGGCCTAAAACCTTCACAT AAAAGTGCGTTCAATACGAAACAAATATATTTGTATATATAGAGAGCGAAGC AATCGGTTGCATAAATTGAATTCGTTTCATATACTTCAATATAAATATTATTAA GTACTACAATTTGAAAACATCTTTAAATATACAACATATTTTGAATTAAGTTTA TTTTTTTTTTTTAGCCACATAGAGACATCTTTGTGGCATGCTAAATTCTGTAGTA 20 AAACCTTCTTGGGGAAAGTGAAAGCCACGTATCAGACCAAAATCCACCCAAC CCTGCACACACGCATCCCCATAAAGAACGACCTTGAGCT

Genomic hit, Accession No. CSC:AC014071

25 **Associated ORF**
Genscan ORF1 predicted sequences >16:36:33|GENSCAN_predicted_peptide_2|477_aa
MLILMRPSIKLAANQNAIKAPNGPKNFLDKVLVVRCSLVCLENGHIAVTASGS
NNNNNSNNINLNLKANYQMSATSIRDSFATILLDAQNRVQNATVAAKNFMLPLR
LRSDTSGDTSNNNENNSRRARQAYNCGVNWLTTTHRPKRRRQVHPPLGSTPSCNN
30 NSSKISRNSSSSNNIASATATRIFLGTSAILAIDFDNTRVPGYYQPTGEWIWVSKS
MIKQLFAVAATADDVAAAAASRGNALTFPGKEKGPRKKAEGCGMEWSGVIEWS
GGDVMCVLSSVATVDDDDHHGGGHFDGLLGTPSALIRLNCLINPKKMRMDFEVE
VAWQIARAADLRISMHLNVPIYEMKTMKTMSVIDGGSLYQPTALFGSLFCLVY
SSAADVLLLLANCKSLAHGVDVDCDSASRGSDCDVGHFSPSFRCRFQLSLVAQS
35 ARHANALKSQVTTATSSSSNNSDSLANKOTNQHFVYQLSA

[illegible]

ggaaaaagcgcggaagggtgtggaatggagtgagtgaggatggagtggtggcgatgtgatgtgtgtctctcagatgtg
gccacagtgtgacgatgatgatcatcatggtggcgccacttgcagcgctgttgggaacaccttcagcgctcatccgacttaactgc
ttaatcaaccggaagaagatgaggatggacttggagttgaggttgcacatggcaaatgtctcgagctgctgatctcggtgctgatcga
atgcaccttaatgtgccttatgaaatgaaaacgatgaagacgatggagagcgtgatcgaatgggtggctccctgtaccaaccgactgc
5 tctctcggttctttgtttgcttgggtgtattcttcagctgctgatgtgtgtgtgctgctggcgaaactgtaaaagcttagcccatggcgctg
acgtcgactgcgacagcgacgctagccgaggcagtgactgcgacgttgccacttttcgcttcgtttcgtgtcgttttcagttgtc
tctcgttgctcaaaagcgcgcggcacgcgaacgctctgaaatccaaagtacaacagcaacatcaagcagcagcaacaacagtga
ttcgctggcaacaacaacaacaacaacatattttgtgtatcaattgtcggcctaa

- | | | |
|----|---|---|
| 10 | <i>Drosophila</i> Gene Hit | rescue sequence, ORF1 and genomic sequence: Canton S E78B nuclear receptor superfamily protein (U01088) |
| | <i>Drosophila</i> EST | LP11082 (AI296953 similar by BLASTN to U01088) |
| | Annotated <i>Drosophila</i> genome genomic segment | AE003593 |
| 15 | Annotated <i>Drosophila</i> genome Complete gene candidate | CG18023 - Eip78C
Ecdysone-induced protein 78C nuclear receptor NR1E1 |
| 20 | Human homologue of Complete gene candidate | CG18023- 4e-32 119100
P20393 EAR1_HUMAN V-
ERBA RELATED PROTEIN
EAR-1
>gi 1082832 pir A32608 |
| 25 | Putative function | ligand-dependent nuclear receptor , putative transcription factor |
| | Confirmation by RNAi | Not done due to failure of PCR procedure |

Example 14 (Category 1)

	Line ID	876/2
	Category	Meiotic defects in testis: cytokinesis defects
5	Reversion	?
	Map Position	73A
	Rescue ID	2H1E
	Rescue Sequence	
10	GATCAAACAGAAAATCCAAAAACGAACAGCGCGCGGCGAACGAGAGCCGTT	
	GAAGCCGGCAGAGAAGTGCGCTGCTCGCGTCGCTGCCGGTATGTGCGTGTCTG	
	TGCACTGAGAGAAAATGCTCGATTAAACAGAGAAAATTAATAGTAATATAAAA	
	AAAAAAAAAAATTTGTTTATTATTCTCAATTCAATAAAATGTAATTATTTATTAT	
	ATTGGTTGTATAAGAATTTTTATAAAGTAGTATAAAATTTCAATCAAATAAAT	
15	ATGTACATCTAACAAAAAATGTTATTATCTTATAACAAAGAGGTAAAATCATA	
	AGTAGTACGAAATCTTTAAAAGAGAAAGTGTGTTACGCAAAAAGTATTCAAA	
	GCAGTCTTTTATTTAATTTAATTAATTTATTTGTGCTTTATCCCTTATATATATA	
	TGTACATTTTCATTAAAGCTAATGGTATAATTAGGTATTTACAGTGTTTAGCTAA	
	GGCTTTTCATCTGAAATATTTATTAATTATGTCTAGTTGACCTGTTTTTAGTTTTT	
20	TTGNATAACAATATTTATTATTTATTAAGGAAAACAAGGGGAGAAGAAAAAC	
	CTTAATTGAAGCAAAGCAGTCTTTTGAACCCACTGGTG	
	Genomic hit, Accession No. AC005633	
	Drosophila Gene Hit rescue sequence: argos (M91381)	
25		
	Annotated <i>Drosophila</i> genome genomic segment	AE003527
	Annotated <i>Drosophila</i> genome Complete gene candidate	CG10162 – Egf2 translation facto
30	Human homologue of Complete gene candidate	CG10162 - 4e-11 181969 (M19997) elongation factor 2 [Homo sapiens]
	Putative function	Translation elongation factor
35		
	Confirmation by RNAi	Not done due to failure of PCR procedure

CATEGORY 2: FAILURE TO ENTER M-PHASE**Example 15 (Category 2)**

	Line ID	1216/12
5	Category	Meiotic defects in testis: no division (no meiosis)
	Reversion	NR
	Map Position	82F1-2
10	Rescue ID	2I5X-1
	Rescue Sequence 1	AAACCAAGCAACAGAAATATCTCCAGTAGAGAGCGCCACTGGAAGATCGGAA TTTTAGTGCTCTGCTCTGACTAACAGGTTTTAGTAGTAGTGCTTACTTTCTAC TACGATTTTTGTCGCGGCTAACAAATCTGTTTTCCCACTCCCTCTCTCAGTTTTT GCATGGTAACTTTTCGGTCATTGTACTGTTGTTGTTGTCTTGCACACCGCAAGA GAACAACAACAATCGGAGAAACACTGATAGCGCGGTACAGTGGGGCAGGCCA AACTAGAACCTATACATTTAAGATGTCTCCAATTTGTGATTTTGCCTTTCAAGC ATACTAGTTCATAGTTGATTGTTTTGTTATGTTTTGTCTTGAATGCGATGTTTCA AGAAATCTTATTTTCGAATTACGATATTATTCTTATTCCTTTGACTTATTAAAA TAAATGAAAACGGCGAGTAGAGCAAAAGAGCGACCACTGTGGCTCCACAAGC TCGTTTCTCTGTTTCTCATTTCGCGCCAGCTCCAATTTGCCTTATTCACACACA CACCTCACTGCTTGC GACTGCAAATTTGTGCAGCTGAACTTTG
	Rescue ID	2I5E-1
25	Rescue Sequence 2	CTTGGTTTATCACCTCTCTCTCTCTATCGCGCGCGCGCTCTTTGTGGAA ACAGGTATAACTGTTTGGCGTGAGGGAGCACGAAACTCCAGTGAGACTTCTC CGCATCGCCAGCGAAACAAACGATCAAAATGAATACTCTGATAACGTGTGAA GGTGAGCAACAAAATAAAGTATAAGAAAAATACCGCCACGAAAACAACAACA ATAGAAATGTCGACGCACCCCTTTTCTTTTCTCGCAAAGAACGAGGAAATGGA GAAGCGCAAAACCACATCCCGCTTAAAGAGTCCCTTTCCCCCGCTGGAAGTGG AAGGAAAGGCAGCTTAAAGAGGAGCGGGTGGCTTCCAGTATGTGGAAAACAA AGCAGACGCCATTGGAATGCCGTCGTTTTTTGTTGTTGCTAAGCCGGACATGG CAATTGTTGCTTTTGTTCGAGAGGGGGTGGTGAAACTCATAAATATCAGCT ATGGCGAGGGGGTGGGGGCAGTCTTTGTCTGACGTACCGTACTTTTAATTTCTT GTCGCCCGGTTTAATCCAATTTATCCAGCTTTGAATTCGCGG
	Genomic hit, Accession No. AC007532	
40	Annotated <i>Drosophila</i> genome genomic segment	AE003603
	Annotated <i>Drosophila</i> genome Complete gene candidate CG1116 - novel	
	Human homologue of Complete gene candidate	2495728 HYPOTHETICAL PROTEIN KIAA0258(aa)

Putative function No homologies which indicate function

Confirmation by RNAi Slight loss of G1 peak

Example 16 (Category 2)

	Line ID	1344/15
	Category	Mitotic defects in brain: no mitosis
5	Reversion	NR
	Map Position	83C
	Rescue ID	2F6E
	Rescue Sequence	
10	AGCGGGAGTGAGCCGAAAGAGAGTAATTTGGCCGTCACCAAAAAAGTGGCT GCATAGTGCCAAACCAATGTATGGCCGTTACGCATCTTGTTATTCTAGTGTCTT TGGCTGTAATCAGTTTGCAGTGACAGAGGAGTTCAGTTTCAGTTGACTCGGCT TGGTTCAGGGTTTCTGATTGCCGTCCTCTTCTCCCTCTTCGCCTACAAGTCCGC TGTTCCGGCACCGTGACGTCACCTAGACTTACACCCCTAATCAAAGATCCACTA 15 GTTTAGATTTTCCTGCATCAACGCCATATTAACCTTTATAAGCAGTCGTTATATCT CAAGTAGGCCAAAAAGTGTAATAGATATGTATCTAAATTGTCGTACATTCTAT TTATTAATAATTCGTTTTTACATCCAACAGGTGTTATTTTTGAAGTCTTAGATAA CAAACAATATTCGAATTATGTGGTAGAATACTTAGCAATATACGCACATACAT ATACATATGAACATTATATCCAATGCTTTAAAACCGGAATATCAAGACAACAT 20 AATGCAACATCTGGTCCGAGCTATCCAGGCAATCACATTTTTGAAGTTCCTCCC GGTTATCACACATATATCGATCATACCCCGAAATGTGTAAACACAGATACAGCT CACCATCCCTCTGATAAGATCTTATCAAGTTCGGGCTTGCTCGCTATCGTGAAT TGGGTTGAAGGGTCCGCGATAATTGCATTGGGCATGCCATTGGTAATCACAAT TGGCTGATAATGCTGCTGCTGCAATTCCACGGGTATGAA 25 TTCATCAATTGGTTA	
	Annotated <i>Drosophila</i> genome genomic segment	AE003602
	Annotated <i>Drosophila</i> genome Complete gene candidate	CG1347 - novel protein with myosin homology
30	Human homologue of Complete gene candidate	1503990 dbj BAA13194 (D86958) KIAA0203 similar to mouse CC1.(aa)
35	Putative function	similar to coiled coil protein with ubiquitin like domain
	Confirmation by RNAi	Marked reduction of G1 and G2/M indicating fewer cycling cells
40		

Example 17 (Category 2)

	Line ID	703/16
	Category	Meiotic defects in testis: segregation defects, meiotic failure (Mf-07/75)
5	Reversion	R
	Map Position	83B
	Rescue ID	2E7E
	Rescue Sequence	
10	AAGCAGCCCAACAGCTACGCAAAAAGTTACTTATATTTCGCAGCAAAACAGAT	
	TTTTTTGTTTTAATCGTAAGTATAGGAGTGAAAAATAGCGCTAGAGTAGACCT	
	AAGTACACAGAAAGACAAATAGGGCGAGTAAAATCGCGGTCCTGGTCATTTT	
	TCTGGCCTTGACCAATCCTTTGTCTGCGCTTTCGTTGGAAAAGGGGTTATGTAC	
	GAACTGCGTGCGTACCTAAGGCCAGATTAGTCATCGGGCAGTCATATATTCAT	
15	GCAAAAAATCATTTGGTGGCCGTCGGCCTTTGTTCGACTGTACCTTGCTCATT	
	TTTAATAAGCGCGACAGCAATATACACACTTTGAACCCCCATCCCACATTTTTT	
	CTCACCGTTTCCCCCTAATTTTCGTTTTCCCTGTGCCCATCATTCGCTTTTCGCC	
	ATGTCAGTGTATCGCTTCAAAAATGGCGCCGAACCACATGTCTTCGTTCTCGGC	
	TCGTCCGCTTCGTTTCGTGCGCTCGTGTGTCGTCTCATTCGCTCTCCGAATTCG	
20	TTTAACAAAGTGTTGCGAGCAGAGGGGCCGCTGGATTTCGAGGCAAACAACAC	
	ATATACCTA	
	Genomic hit, Accession No. CSC:AC013960	
25	<i>Drosophila</i> EST	several including LD15903 (AA440858), GH20091 (AI389018).
	Annotated <i>Drosophila</i> genome genomic segment	AE003602
	Annotated <i>Drosophila</i> genome Complete gene candidate	CG2922 – novel
30	Human homologue of Complete gene candidate	286001 dbj BAA02795 (D13630)
		KIAA0005 [Homo sapiens] also
		NP_054757.1 HSPC028 protein
		[Homo sapiens] e-179
35	Putative function	Weakly similar to a region of human and murine EIF4G2 translation initiation factors; may act as a translation initiation factor
	Confirmation by RNAi	Only wild type profiles observed

Example 18 (Category 2)

	Line ID	741/3
	Category	Meiotic defects in testis: segregation defects, meiotic failure (Mf-05/31)
5	Reversion	NR
	Map Position	88D
	Rescue ID	H6E
10	Rescue Sequence	GCCTGGAGCCACCTCTAGAGCCACGGCCAAAAAATTGTGTGCCAAAAAATCG TATGGCGTTACGCATCTTGTTATTCTAGTGTCTTTGGTTCTACAAATCTGGCCA ATGGGATGGACGGATTTTGGGGCTTTTGC GCCCCACATATGTNTCTTACAACC CACTCGGCCCCGCAAGTGGGTGTCAATTACGGACATCGGCAATCCGAAGACC 15 GGAGACCCAGAGACCTCAGACCCCAGGGCCCCATTTCGATTTCGATTTCGAGTT GCGTGGGCCGATCTCACATTAGTCACATCGAAGGAATGAAATAAAAAGAAAA AACATGACGGCCGAAAAGAACTTATCCATCTTCAAAGCTCTCAGAAAATACA AAAAACTAAAAAACTTTTGACTCTTCGTCTTTCACATTTTCGAAATCACAAAAT GTCTGCCATAAATTCCAAAGTGAACAATTGAAATAAATTTTTGCGCCATGAAC 20 ACGCCGACTG
	Annotated <i>Drosophila</i> genome genomic segment	AE003705
	Annotated <i>Drosophila</i> genome Complete gene candidate	CG12600 - novel protein
25	Human homologue of Complete gene candidate	CG12600- 5e-27 4240227 dbj BAA74892.1 (AB020676) KIAA0869 protein [Homo sapiens]
30	Putative function	putative cytoskeletal structural protein
	Confirmation by RNAi	Reduction of G1 and G2/M peaks indicating fewer cycling cells

Example 19 (Category 2)

	Line ID	773/1
	Category	Meiotic defects in testis: cytokinesis defects, meiotic failure (Mf-02/15)
5	Reversion	R?
	Map Position	83F
	Rescue ID	2D9P
10	Rescue Sequence	CCACCGCCCATGCCGCCATTTATTGAAAGGCCTGTACGCAGTTTGTGTTTTGTTT TTCTCTTTTTTGCTAGCTCAAACACAAAATTACTTTTTGTGGCTTGACTGGTGA GGTCTCTCTATCTCGCTTTTTCGTCTTTACCTCGCTCTCATTCCTCTCTATCTG CCCTGCTTCCTCTCACTATCTATCTACAAGTGAAGGTCAACAAAATAAGTGCGT 15 AGTCAAAAATGTAATTGAATTGATTGACAAACACAGCGAACGTAAATTTCCGT AATGTTTAACCTTGAATTCAAATGAACAACTGTATAAATATAATACACGGGTT AAACTCCATTTCAAAGCAAGCTAAAACATTTTAAATACATTTTAGGGAAACGG CCAATTAAAAGAATAATATTGTGGGGATCAATCTGGGGAAAAATGCAGTATC AGTAATGCTGAATATTTATTTTACTAAATTACAATGAAATGTCTCAAACAAAT 20 GGGTTAATCATTTTCTTGCTCCATCTGCTTTTCCCAACTGTATCCAAGTACAAC TACAGCATTATCCTCAACTG
	Annotated <i>Drosophila</i> genome genomic segment	AE003675
25	Annotated <i>Drosophila</i> genome Complete gene candidate	CG10272 - novel protein
	Human homologue of Complete gene candidate	CG10272 - 2995577 AC004490 (AC004490) R29381_1(aa) protein includes HMG-I and HMG-Y DNA- 30 binding domain (A+T-hook) found in HMG non-histone components in chromatin
	Putative function	Chromosomal protein
35	Confirmation by RNAi	Loss of G1 peak indicating arrest in G2/M

CATEGORY 3: METAPHASE ARREST**Example 20 (Category 3)**

5
Line ID 1067/13
Category Mitotic defects in brain: prometaphase arrest
 (overcondensation, polyploidy, scattered chromosomes with
 bipolar spindle)
 10 **Reversion** NR
Map Position 69C4-10
Rescue ID 2F8E
Rescue Sequence
 15 GTTTGGGCACAGGGTTGTATTTCAATTTATTTTTGGGGGGAGTCGATACGCTCTC
 TTGGCGTGGTTCGAACGGTCACACTGGCCGAGAGATAACGGAAAATGTTTCAA
 AGGTAAGTAAAGATTATAAACGTATTAAGCTTAATACTATAATTAGCTTACTA
 TTCCAAGTATGTATAATTATTACACGTTTAAAAGGCATAACGTTAAGTGTAAC
 CAAATTATATCAATGGATTTTGAATACCAATATTATTTATTTTATATTTTGAGC
 20 TTAATATATTAAATCACATATATTTAAGCCTCTTTATATATGTAAATATTTTAA
 TTTTATTTAAAATAAATTATATATTGTTTTGTAAATATGATCGAGGGCTGCCACCT
 TGTGATAAATGCTTACCAACACTTTTAGGTACGCCGTTTAGTGTACGTAAGTTG
 CGTACCTAGATATCCAGCGAAATCAAAACATTGAGTAAATCGTGGAATAATGG
 ATGAAAATAGCTTAATCTACGGACTCGAACTGCAGGCGCGGGCTTTAACACCT
 25 CAGTACGGAGAGAGCAACGATGTGTGCTTCTTCATAGCCACCAACTCCTTGAA
 GCCCACCAATCAGGTTCACTTAATCCAGTACGAAGA

Genomic hit, Accession No. CSC:AC020333

30 **Associated ORF**
 Genscan: ORF1 predicted sequences: >16:51:11|GENSCAN_predicted_peptide_2|178_aa
 MAQNISPEQSGGAGGGGSKHSDDSM PVKDNHAVSKRLHKELMNLMMANERGIS
 AFPDGENIFKWVGTIAGPRNTVYSGQTYRLSLDFPNSYPYAAPVVKFLTSCFHPNV
 DLQGAICLDILKDKWSALYDVRTILLSIQSLLGEPNNESPLNAQAAMMWNDQKEY
 35 KKYLDAFYEKHKDT

>16:51:11|GENSCAN_predicted_CDS_2|537_bp
 atggcgagaatcagccccgagcaaaagtggaggagcggcgaggcagcaagcacagcgatgactccatgcccggtg
 aaagacaatcacgccgtgagcaaaagactgcacaaggaactgatgaacctgatggccaacgagaggggcatctcagcgtt
 40 tccggagcggcgagaacatctcaagtgggtgggcaccatagcgggtccacggaacacgggtgtattcggggcaaacgtatcgttt
 gtactggatttcccaattcctatccgtatgcagcaccggtgggtgaagttcctgacgtcctgcttccatcccaatgttgatctgcagg
 gcgcacatctgtttggacatactgaaggacaatggcggccctgtacgatgtgcgcaccattctgctgtccatacaatccctgctgg
 gcgaaccgaacaacgagagtcactgaatgcgcaggccgcgatgatgtggaatgac

- Drosophila* Gene Hit** TBLASTX with ORF1: poor homology to several sequences including homolog of RAD6 (DHR6) (M63792), bendless (L20126) and Ubc D1 mRNA for ubiquitin-conjugating enzyme (X62575).
- 5 **Human Homologue** TBLASTX with ORF1: ubiquitin carrier protein E2-C (UBCH10) (NM_007019.1) and ubiquitin-conjugating enzyme E2B (RAD6 homolog) (NM_003337.1).
- 10 **Annotated *Drosophila* genome genomic segment** AE003541
Annotated *Drosophila* genome Complete gene candidate CG10682 – vihar ubiquitin-conjugating enzyme
- 15 **Human homologue of Complete gene candidate** gi5902146
0B6F58A1F0665D9A
|ref|NP_008950.1| ubiquitin carrier protein E2-C [Homo sapiens] (2.50E-50)
- 20 **Putative function** Cyclin specific ubiquitin conjugating enzyme
- 25 **Confirmation by RNAi** Complete loss of G1 and G2/M peaks indicating fewer cycling cells. Immunostaining shows metaphase arrest with condensed chromosomes

- Line ID** 1105/1
Category Male sterile, Female sterile, Mitotic defects in brain: prometaphase arrest
(Overcondensation, polyploidy, fewer anaphases, high mitotic index, scattered chromosomes with bipolar spindle)
Reversion R
Map Position 69C
- Rescue ID** A5B
Rescue Sequence
GTACATATAATCACAATTGAGAATCGAAAACCCGACCGCCACGAAGCGCGCT
AAATTACACGCACATACTGAAAGCCAAACAGCGGATAGCACTAGCATCCTAC
ATATATAGACGTAGATATATAGTCATGGCGCAGAATATCAGCCCCGAGCAAA
GTGGTGGAGCAGGCGGCGGCGGCAGCAAGCACAGCGATGACTCCATGCCCGT
GAAAGACAATCACGCCGTGGAGCAAAAGGTGAGTATCATGGTGCAGCCTA
AGATAATCCGCCAATATACACACACACTCACACTACCCACAGACTGCACAA
GGGAACCTGATGAACCTGAATGAATGGGCCACCGAAAAAAGGGG
- Rescue ID** A5E
Rescue Sequence 2
ATATGTACTGTATAGTGGAAATTTAGTTTGATCGGTCCGAATACGCGTCTGTT
GCTTTTTTCTAGATATTTTTTTTTTCACTTTTGTGTGAAAACAAAATGGAAGGAGA
ACGAGAAGAACTGTGTTTGGGCACAGGGTTGTATTTCATTTTTTTTTGGGGG
GAGTCGATACGCTCTCTTGGCGTGGTCCGAACGGTCACACTGGCCGAGAGATAA
CGGAAAATGTTTCAAAGGTAAGTAAAGATTATAAACGTATTAAGCTTAATACT
ATAATTAGCTTACTATTCCAAGTATGTTATAATTATTACACGTTTTAAAAGGCA
TAACCGTTAAGTTGTTAACCCTAATTATATCAATGGATTTTGAATACCAATATT
ATTTATTTTATATTTTGGAGCTTAATATATTAATCCACATATATTTAACCCCCCT
TTATATATGTTAAATATTTTAATTTTATTAATAAATTATATATTGTTTGGTTA
AAA
- Genomic hit, Accession No.** AC007328 69B-69C
- Associated ORF**
Genscan: ORF1 predicted sequences
>/tmp/aaaaanjda|GENSCAN_predicted_peptide_1|357_aa
MGKKAKHKKKKGKGPKEKTAMKADKKQAARQKKMLEKLGEANIADIQLLEAKEG
KIEAISESVCPPTPRSNFTLVCHPEKEELIMFGGELYTGKTTVYNDLFFYNTKTV
EWRQLKSPSGPTPRSGHQMVAVASNGGELWFPNFACISRNQSWFVFNCRKAA
SREKVLNLFNGTVLHPANNIIVHVKLFKKANGFKPWLLDVKLDACRFVRTNFHPF
VRIIFDLFKDFSTINHTCPYVVLRSRMRYIVRRSPRLVHPIVDVPAIGHTRPRRKA
RGIGCAHRCPLIRMATPCRTNVMMTMLMRGSRVSRVMAICCYRRPAIAIARRRHP
TAIAHSQEVAERLGGLLYPDIQRTNP
- >/tmp/aaaaanjda|GENSCAN_predicted_CDS_1|1074_bp
atgggcaaaaaggccaacacagaagaaggcgcaaggcccgagaaaaaggccatgaaagcggacaaaaagcaggcgg
cgcggaagaaaatgctgaaaaactgggagaagcaatatagctgatatcatccaattgctggaggccaaggaggcgcaag
attgaagccatcagtgaatccgttggccgccaactccacgatccaattcaccttagtttgccatccggaagaggagctc

atcatgtttggcggcgaactgtacactggcacaaaaaccacagtgtataacgattgttctttacaacacccaaaaccgtcgagtgg
aggcagctgaaatcgccatcgggaccacgcccagaagtggacaccaaagggtgctgtggcagcaatggaggagaactct
ggtttccgaacttcgctgtataagtcgcaatcaatcctggtttgtgtccacaattgtcgtctgaaggcggccagtcgtgagaaggt
cttactcaactttaatggaacgggttacatccggccaataacataatgttcacgtcaagctgtttaaaaaggccaacggtttaagc
5 cttggttattagacgtaaaactcgatgcttgcgtttgtgcggaccaacttccatccgtttgtacgcattatattcgatctcttcaaagat
ttctccaccataaaccacacgtgcccataatgtggtcctccgatcgatgcggtatattgtccgccgatccccacgacttgtgcacccc
atcgtagatgttccggctattgggcacactcgcctcgacggaaggccgccgttcgtggcatagggtgtgtcatcgctgccctct
gattcggatggcgactccgtgtgtaccaacgtggtgatgatgacgctgatgaggggctcggtgagatcgaggggtgatggcgatt
tgetgtaccgccgaccgccattgccatagcccgtcggcgccaccccactgccattgccactcccaagaagtgtctgaacgc
10 ctcgggtgtcttcttaccggacattcagagaaccaatccgtag

Drosophila EST several ESTs including LD04777 (AA201675)

All other entries as for 1067/13.

Example 21 (Category 3)

	Line ID	1407/13
	Category	Mitotic defects in brain: (weak overcondensation, metaphase with bipolar spindle)
5	Reversion	NR
	Map Position	92B1-3
	Rescue ID	2D3P
10	Rescue Sequence 1	ATCACGAATTTGACATTGCTACCACATTCGGTGCGTGGACTCTGAAAGCTCTG AGTGTTTTGTTTATGCAAAGCTTTTTTGGACTATCGCGTGGTAAGTAGCCGAAA GAGAAAGCTCTCTTATACGGAAGATGAAGAGTGTGATTCATGAAAATGTATA AGAACGCGGGTCCAAAAAGTCAAGGGAGTTCTAGTGAAATGAAAAGTTCCAA 15 AGGTTTTGAAATCGTTTTATTTCTCGTTCGTATAATTATTGGGTGTCGATCTTT GTTGGGCAGTGTAAGCACAACTTTGAGCTTCATCATACATATCATATGTAA AGCCGGGACGAAAGCTTATGATTCTGTAAAGTGTCCGCCCAAGATAACATTC TCCAGCCCTTCAAATCTTCAAATAAATACGGCTTAAGGCGAGCAAATTTGTAA ATCAAATGATTTGTAAATAAACATTATATGTATTTATCATGCCAGGTTAGAA 20 CACATTGTGCTGATGCAAATAAAATTCCAATTAAACGCCCTGAATGGGAAGA TGACGCATCTTTAATGGGAATATTATGGTAAATTTAATA
	Rescue ID	2D3E
	Rescue Sequence 2	TNCGTGATTATCAGCGTTAATTGTACAATATTATGATTTATTCGAGCTGTAAAT 25 CTTCACAGCAAGCACAACTGTAATTATACCACTTAGAATTCCGCGGAATTAA TTCTTGAAGACGAAAGGGCCTCGTGATACGCCTATTTTTATAGGTAAATGTCAT GATAATAATGGTTTCTTAGACGTCAGGTGGCACTTTTCGGGGAAATGTGCGCG GAACCCCTATTTGTTTATTTTCTAAATACATTCAAATATGTATCCGCTCATGA 30 GACAATAACCTGATAAATGCTTCAATAATATTGAAAAAGGAAGAGTATGAG TATTCAACATTTCCGTGTCGCCCTTATCCCTTTTTTGCGGCATTTCCTTCCT GTTTTGCTCACCCAGAAACGCTGGTGAAAGTAAAAGATGCTGAAGATCAGTTG GGTGCACGAGTGGGTACATCGAACTGGATCTCAACAG
35	<i>Drosophila</i> EST	LD05707 (AA246767)
	Annotated <i>Drosophila</i> genome genomic segment	AE003727
	Annotated <i>Drosophila</i> genome Complete gene candidate	CG7444 - very short ORF with EF hand homology
40	Human homologue of Complete gene candidate	none
	Putative function	Possible calcium binding protein
45		

Confirmation by RNAi Slight loss of G1 peak

Example 22 (Category 3)

5	Line ID	1439/7
	Category	Mitotic defects in brain: prometaphase arrest. (overcondensation, polyploid, no anaphases, scattered chromosomes with bipolar spindles)
	Reversion	?
	Map Position	96F10-14
10	Rescue ID	G3X
15	Rescue Sequence	
	GTCGGATGTAGAAGACGTGCCCCGAAACCCAGTTAGAAATCGATGTCAGCGAT	
	GGCGCCGGACTGGAGGATGAGGATGATGACGATATGGAACAGATTACAGCTC	
	AGAAGGTAAGGTAAATCGTAACAGAGCTTTTAAATACGCAAGTAATCACATTC	
15	TGATATCCCTAGGTTCTGGAAATCATAGAAACCGCGTGGATAAATGAAATGTG	
	TGCGCCGGAGATCCTGCCAGCCAGACGGACATGCTGGAGCTGATGGTCTCCC	
	AGGTGGCCCATATGGAGGAGCAGATGCGCGATCTGGACAAGAACGATTTCCG	
	AGCGGTGGTGCACCTCCATGGAACCTGGAGAGGGTGGCTACATAATGGCCAGT	
	TATCTGCGTTGCCGCCTGCAGAAGATCGAAACCTTCACGCAGCACATCCTCAA	
20	CCAGGAGGAGAGCCGTGAGCCGGATGACAAACGTCTGTCTCCCGAGGAGACT	
	AAGTTCGCCCAGGAGTTTGCCAGTAAT	

Genomic hit, Accession No. AC007825

25	Annotated <i>Drosophila</i> genome genomic segment	AE003754
	Annotated <i>Drosophila</i> genome Complete gene candidate CG14549 – novel	

Human homologue of Complete gene candidate none

30	Putative function no homologies which indicate function
----	--

Confirmation by RNAi Only wild type profile observed

Example 23 (Category 3)

Line ID 1466/4
Category Mitotic defects in brain: metaphase arrest.
(overcondensation, no polyploidy, fewer anaphases, metaphase
5 with bipolar spindle)
Reversion NR
Map Position 72F

Rescue ID ESE
10 **Rescue Sequence 1**
GGCTGGATGCGATTTCGCTTTCGGATTTCGGATGGATTTCAGCCGCTGTCTCGACA
CCGCCGCAACCGCTCTCGGGAGTTTGAAAATTTGAAATGAGCGGATTTCGCGTT
GCGAAGGCGAGCTAGCGTTGCAGGCAGTGTGGCCAGATGCCGCGTGCGAACG
TATTCTCGAATGCAATCGGCCGAGTGCAGATGCACTAAAAATAACCCACTTCC
15 AGTGACTGGAAATTAAGATCAAGGNAATAGATTTTATAAAAACTTATATGAGT
AAAAATTTTAAATTTGTGGAGTCAACCTAAATTATAAGCAACTAATTTATAAC
ACAAGTAAAGAATGATATTAAGTAACTTTTTAAATAATATTCCATTATGCTTA
CGCTCAATTTATGAACAAATGTTTTCTCGATCCTTAGGTAAAGTTTCGAGTTTC
GCGACTANATTTATTAATAAATTAAGAACATCTCCATTATGTACACATTTAAAG
20 ATTTATGAGCGGTAATATTAGCTGGTTGAC

Rescue ID E5P
Rescue Sequence 2
ATCCAGCCAAGATATCCTATCGTGCAGCTGAAACCCGAAACCCGAATCCGAGT
25 TCGAAACGAAACGAATCGCAGTGGTGGTTTCTCTCTCGCTCTCTAGCTCTCCCT
CTCTCTCGCGTGTGTGTATGTGTGCGAGTGGCAGGAAAAGTGCGAAGCCGAAA
TCTTTTCTAGCTGAAAGAAAGCGCAACTTCAATTAGCGAAAAGCAAGAGTAGCT
AACAAAAAGAAAAGCGGATCGAAAAGTAAAGAAAAACAAAAAACA
AAAGCAACAAATCGAAATGGCAAGCGAAGTGGCCCAATACCCGCCGAGGG
30 AAACGCCCCGAGTGGCGGCGGCGGAAAAATCAGAGGAGCCGGAAGAGTCAG
CGGCCCCGCCAGCGGACTCAGCGCCGCTCCAGCTGCCGCCCCCGCAGTGGA
GAAGGCTGAGGATGCCGATGGCGAAAAAAGGACGGCGAGGCCGGAAGCA
GGACAAGCAGCAGGATGGC

35 **Genomic hit, Accession No.** CSC:AC020154

Associated ORF
Genscan ORF: ORF2 predicted sequences
>21:06:03|GENSCAN_predicted_peptide_5|415_aa
40 MASEVAQIPAEETPAVAAAEEKSEEPEKSAAPPADSAAAPAAAPAVEKAEDADGE
KKDGEAGKQDKQQDGEEPKKDEAVAAPVATKSEAPPAQKFNVHKTNFEKDIIYL
YQFSRTPLPSLSPYCLKVETWLRVLVGLKYENVVDHKMRFRSKKGQLPFIELNGEEI
ADSAIIKELSSKYEKYLDSGLTAEQRNVSYATIAMLENHLIWIIFYWRAKYPDNV
LKGYKVNQLHALGLRLPNSILNFFFKITFGRKGTKKLKAHGIGVHSAEEIEEFGKD
45 DLKVLSEMLDCKPFFFGDEPTTLDVVAFAVLSQLHYLSKDIA YPLRDYMTKECPN
LIGHVSRMKDKCFPDWDEICTKLDLNAHIPKPEPETKEGKEGGEQKSNEQEGTE

GDKIEKELEKDKSNEKESTEENKEKEETK

>21:06:03|GENSCAN_predicted_CDS_5|1248_bp

atggcaagcgaagtggcccaatacccgccgaggaaacgccgcagtgggcggcgggaaaaatcagaggagccggaaaa
 5 gtcagcgggccccgccagcgggactcagcggccgtccagctgccgccccgcagtgaggagaaggctgaggatgccgatggcga
 gaagaaggacggcgaggccgaaagcaggacaagcagcaggatggcgaggagccaaaaaggacgaggcgggtggcagc
 acccgtggcgacaaaatcggaagccccgcccgccagaaattcaatgtgcacaagaccaacttcgagaaggacatcatctatct
 gtaccagttctcgcgcacccactgctgccctccctgtcgcctactgcctgaaggtggagacctgctgcgtctgtggcctga
 aatacagagaatgtgatcataagatgctgtccgctccaagaagggtcagctgccgttcacgagctgaatggggaggaaatcgc
 10 cgattcgcccatcatcatcaaggaaactgtcgtccaatacagagaagtacctggactcgggactcaccgccgagcaaatgtgtctaa
 ctctacgccacgattgcatgtggagaacctctcatctggatcatcttctactggcgcgccaagtatccggacaatgtgtctaa
 gggctacaaggtaacttcagcacgccctcggcctgcggctgcccaactcgattctgaacttctttaaagatcacctttggtcgc
 aagggcacgaagaagtgaaggcgcacggcatcggtgtccacagcgcgaggagatcaggagttcggcaaggacgacctg
 aaggtgctcagcgagatgtcactgcaagccttcttctcggcgacgagcccaccacctggatgtggtggccttcgctgtcct
 15 ctccagctccactatctgtccaaggacattgcgtatccgtcgcgcgactacatgaccgagaagtgcaccaacttgattggccacg
 tatctcgcatgaaggacaagtgttccccgactgggacgagatctgcacgaagctggacctcaatgcgcacattccaagccag
 agcccagaccaaggagggaaggagggtggcgagcaggagaatcaaacgaacaggagggcactgagggcgacaagat
 cgagaaggagttggagaaggacaagtcaacgagaaggagtcgaccgaggagaacaaagagaaggaggaacaaagtaa

- 20 ***Drosophila* Gene Hit** rescue sequence and TBLASTN with ORF2: failed axon
 connections (U21685)
Human Homologue BLASTX with fax: Metaxin 1 and 2 (Q13505 and AF053551)
***Drosophila* EST** several including LD31362 (AA951078 similar by BLASTN to
 U21685 failed axon connections)

25 **Annotated *Drosophila* genome genomic segment** AE003527

Annotated *Drosophila* genome Complete gene candidate CG4609 – fax failed axon
 connectionsconnections

30 **Human homologue of Complete gene candidate** 4505281
 ref|NP_002446.1|pMTX|
 metaxin>gi|3024205|sp|Q135
 05|MTXN_HUMAN
 METAXIN (4e-06)

35 **Putative function** *Drosophila* fax is a dominant genetic enhancer of the Abl mutant,
 developmentally expressed in axons of the CNS

40 **Confirmation by RNAi** Weak reduction of G1 and G2/M peaks indicating fewer
 cycling cells

45

Line ID 262/20
Category Mitotic defects in brain: metaphase arrest.
(overcondensation, polyploidy, aneuploidy, few anaphases, high mitotic index, metaphase with bent bipolar spindle)

5 **Reversion** NR
Map Position 72F

Rescue ID G6E
Rescue Sequence

10 AGCTGCACGATAGGATATCTTGGCTGGATGCGATTCGCTTTCGGATTCGGATG
GATTCAGGAGCCGCTGTCTCGACACCGCCGCAACCGCTCTCGGGAGTTTGAAA
ATTTGAAATGAGCGGATTCGCGTTGCGAAGGCGAGCTAGCGTTGCAGGCAGT
GTGGCCAGATGCCGCGTGCGAACGTATTCTCGAATGCAATCGGCCGAGTGCA
GATGCACTAAAAATAACCCACTTCCAGTGACTGGAAATTAAGATCAAGGAAT

15 AGATTTTATAAAAACTTATATGAGTAAAAATTTTAAAATTGTGGAGTCAACCT
AAATTATAAGCAACTAATTTATAACACAAGTAAAGAATGATATTAAGTAACTT
TTTAAATAATATTCCATTATGCTTACGCTCAATTTATGAACAAATGTTTTCTCG
ATCCTTAGGTAAAGTTTCGAGTTTCGCGACTAGATTTATTAATAAAGAACAA
TCTCCATTTATGTTCCC

20 **Drosophila EST** several including LD28084 (AA949260)

All other results as for line 1466/4

25

	Line ID	262/22
	Category	Mitotic defects in brain: metaphase arrest. (overcondensation, polyploidy, few anaphases, high mitotic index, metaphase with bent bipolar spindle)
5	Reversion	NR
	Map Position	72F
	Rescue ID	F1E
	Rescue Sequence 1	
10		AGCTGCACGATAGGATATCTTGGCTGGATGCGATTTCGCTTTCGGATTTCGGATG GATTCAGGAGCCGCTGTCTCGACACCGCCGCAACCGCTCTCGGGAGTTTGAAA ATTTGAAATGAGCGGATTTCGCGTTGCGAAGGCNAGCTAGCGTTGCAGGCAGT GTGGCCAGATGCCGCGTGCGAACGTATTCTCGAATGCAATCGGCCGAGTGCA GATGCACTAAAAATAACCCACTTCCAGTGACTGGAAATTAAGATCAAGGAAT
15		AGATTTTATAAAAACTTATATGAGTAAAAATTTTAAAATTGTGGAGTCAACCT AAATTATAAGCAACTAATTTATAACACAAGTAAAGAATGATATTAAGTAACTT TTTAAATAATATTCCATTATGCTTACGCTCAATTTATGAACAAATGTTTTCTCG ATCCTTAGGTAAAGTTTCGAGTTTCGCGACTAGATTTATTAAAATTAAGAACA TCTCCATTTATG
20	Rescue ID	F1P
	Rescue Sequence 2	
		GTGCAGCTGAAACCCGAAACCCGAATCCGAGTTCGAAACGAAACGAATCGCA GTGGTGGTTTCTCTCTCGCTCTCTAGCTCTCCCTCTCTCTCGCGTGTGTGTATGT
25		GTGCGAGTGGCAGGAAAAGTGCGAAGCCGAAATCTTTTATAGCTGAAAGAAAG CGCAACTTCAATTAGCGAAAAGCAAGAGTAGCTAACAAAAAGAAAAGCGGAT CGAAAAGTAGAGAAAAACGAAAAAAAAAAAAACCAAAGCAACAAATCGAAATG GCAAGCGAAGTGGCCCAAATACCCGCCGATGAAACGCCCGCAGTGGCGGCGG CGGGAAAAATCAGAAGAGCCGAAAATCAGCGGGCCCGCCAGCGGGACTCTG
30		CGGGCGCTCCAGCTGCCGCCCCCGCAGTGGAGAAGGCTGAGGATGCCGATGG CGAA
	Drosophila EST	several including LD28084 (AA949260), LD38479 (AI518768)
35	Other results as for line 1466/4	

	Line ID	262/3
	Category	Mitotic defects in brain: Metaphase arrest (overcondensation, polyploidy, aneuploidy, no anaphases, high mitotic index, metaphase with bipolar spindle)
5	Reversion	NR
	Map Position	72F
	Rescue ID	H3E
	Rescue Sequence	
10	AGCTGCACGATAGGATATCTTGGCTGGATGCGATTTCGCTTTCGGATTTCGGATG GATTCAGGAGCCGCTGTCTCGACACCGCCGCAACCGCTCTCGGGAGTTTGAAA ATTTGAAATGAGCGGATTCGCGTTGCGAAGGCGAGCTATCGTTGCAGGCAGTG TGGCCAGATGCCGCGTGCGAACGTATTCTCGAATGCAATCGGCCGAGTGCAG ATGCACTAAAAATAACCCACTTCCAGTGACTGGAAATTAAGATCAAGGAATA	
15	GATTTTATAAAAACTTATATGAGTAAAAATTTTAAAATTGTGGAGTCAACCTA AATTATAAGCAACTAATTTATAACACAAGTAAAGAATGATATTAAGTAACTTT TTAAATAATATTCCATTATGCTTACGCTCAATTTATGAACAAATGTTTTCTCGA TCCTTAGGTTAAGTTTCGAGTTTCGCGACTAGATTTATTAATAAATTAAGAACATC TCCCTTTATGTTC	
20	Other results as for line 1466/4	

Example 24 (Category 3)

Line ID	238/20
Category	Mitotic defects in brain: metaphase arrest (overcondensation, metaphase with bipolar spindle)
Reversion	NR
Map Position	75E1-3
Rescue ID	D7E
Rescue Sequence	TTCAGTCGCGCATTTTCACCGTTTCCGAATCGGACGAACCGGGCGTGATTGCTC TCCTGCTGCTTTTCGAGATCGGAGTCCCGATAAGGATATAACTACAACCTAAAG AGGAATCCAAGCCTCCTCCTGCCGCTAGTTTCGAAAAGTAAATAGAGTACTTG TTATCAACTGGGGAAGCGGAGATACATAGCTCCGATATTCTGTGAAAGCCAG ACAAACGGATACCAACGAACAATCGCCATGTGCGTCGTCGTCCCTTCTCGTTT CACACATCGTGCGATAAAAAATACCGCTTTGCTTTTTGTGTTTATTTAAAAATTT TGGTTAGGAAGTGAAGTGAAGTGAAGTGAAGTGAAGTGAAGTGAAGTGAAGTGAAGT AGAGCAAAACATAGCAGAAGAACCCAGAAGAAACAGGAACAGAAACCGTT GACCGAGTGCCAGTGTGAAGTCTAGGCACAAAGAACGCTACCAAGAACTCT TGGGAGTTAGGGAGGCTCTTTACAATGACAACATTGCACCAAGATGGACTCT CTCTCTAAAATGCATTTTCATACCAATATTTACTTT
<i>Drosophila</i> EST several including LP04802 (AI260815)	
Annotated <i>Drosophila</i> genome genomic segment	AE003519
Annotated <i>Drosophila</i> genome Complete gene candidate	CG3979 - novel gene with homology to sodium- dependent dicarboxylate transporters
Human homologue of Complete gene candidate	3e-87 4506979 ref NP_003975.1 pSLC13A2 UNKNOWN >gi 2499523 sp Q13183 NDC1 _HUMAN RENAL SODIUM/DICARBOXY
Putative function	sodium/dicarboxylate transporter
Confirmation by RNAi	Only WT profiles observed

Line ID 490/9
Category Meiotic defects in testis: segregation defects, multipolar spindles (Mul-02/29)
Reversion NR
5 **Map Position** 95C1-8

Rescue ID I4E

Rescue Sequence

10 GCTCTGCCGCTTCAACCGCCCGCCTTCTGTGTGTTGGTGTGCCGCGACGTAGG
TG TAGGGTCCGCTGCACACGTGTGTGTGGGAGCGCGAGAGCGGGAGAAGA
GCAGAACGTTTTTGGGCGGCTAGTGGTGGCACCGTGAGCATGCCGGTCGTCGT
AAGATAGGCTTAGGAACACTCAGAGAAAATTTGTTTAGCTCAGCATTTTCCTA
TTATTGAAATCATTATTTGATGGTCTATGGGGGTTTCTTTCGTAGTTATTCAT
AGATCGGCGATTAAAGCTACGCTTAAAGGGTAATTTGTCTGAGATATCTTTGT
15 CATTTAAAGTTAAGTCTCAGCTTATCCAAAAGTCAGTTATTGGAAAAAAGGAG
CCAGCTTTTCAGCAGAGTTCGGCTTAAAGCGCTTATTATCATATTAACCAGCTTA
ATTAATGTATCTTTTAAATTGTTATATGCATTAAATCACTAATTAAGGTGATTA
CCATTTGTACGTTTTTAAATTAAAGTATTTTGATTTTCACTAATACAGGCTCTAA
GCTGATCCAAATCTACAAGCTTAGTTTTTGAATAGTCTTCACATGTTGACTTTT
20 ATTCTCT

Genomic hit, Accession No. CSC:AC015160

25 **Other results same as** 238/20

	Line ID	660/3
	Category	Meiotic defects in testis: cytokinesis defects, abnormal spindles. (Ab-01/03)
	Reversion	R?
5	Map Position	75E
	Rescue ID	H8E
	Rescue Sequence	
10		GCTCTGCCGCTTCAACCGCCCGCGTTCTGTGTGTTGGTGTGCCGCGACGTAGG TGTAGGGTCCGCTGCACACGTGTGTGTGGGAGCGCGAGAGCGGGAGAAGA GCAGAACGTTTTTGGGCGGCTAGTGGTGGCACCGTGAGCATGCCGGTCGTCGT AAGATAGGCTTAGGAACACTCAGAGAAAATTTGTTTAGCTCAGCATTTTCCTA TTATTGAAATCATTTATTTGATGGTCTATGGGGGTTTCTTTCGTAGTTATTCAT AGATCGGCGATTTAAGCTACGCTTAAAGGGTAATTTGTCTGAAATATCTTTGT 15 CATTTAAAGTTAAGTCTCAGCTTATCCAAAAGTCAGTTATTGGAAAAAAGGAG CCAGCTTTTCAGCAGAGTTCGGCTTAAGCGCTTATTATCATATTAACCAGCTTA ATTAATGTATCTTTTAAATTGTTATATGCATTAAATCACTAATTAAGGTGATTA CCATTTGTTTCGTTTTAAATTAAAGTATTTGAATTC
20	Genomic hit, Accession No.	CSC:AC015160

Other results same as 238/20

Example 25 (Category 3)

	Line ID	273/18
5	Category	Mitotic defects in brain: metaphase arrest (overcondensation, very high mitotic index, few polyploids, metaphase with bipolar spindle)
	Reversion	NR
	Map Position	75E
10	Rescue ID	D1E
	Rescue Sequence	
	AACTGGGCTAAAACCAGCTGAAAAGTGGTAAAAAGTAAAATATTTGGAGAAG GAAAGCCTTAAGTTCCTCTCTACGCTTCGTACACGTAATGTGCGTGGTTAATC TACGTAAACAAGTGGAAACCATGTTACGTGCCGTGGCTTTGTGTGTGTCAG 15 TGGTGCTCATAGCACTATATACGCCAACTTCTGGGGAATCCAGTCAGAGCTAT CCCATTACCACGCTAATCAACGCGAAATGGACGCAGACGCCCCTATATCTGGA AATCGCCGAGTATCTGGCCGATGAGCAGCGGGCCTCTTCTGGGATTACGTTT CGGGGGTGACAAAGTTGGACACGGTTCTCAACGAATATGGTTTGTGTTTATAA GTCATGGAGAACCCGCATTAAAGAGCTTTTTATATTCTCCTCAATGTGAATCC 20 GAATCCATATAAAATC	
	Genomic hit, Accession No. AC015160	
	Associated ORF	
	Genscan: >ORF2 predicted sequences	
25	>16:57:34 GENSCAN_predicted_peptide_5 1548_aa	
	MLRAVALCVSVVLIALYTPTSGESSQSYPTTLINAKWTQTPLYLEIAEYLADEQA GLFWDYVSGVTKLDTVLNEYDTESQQYNAALELVKSHVSSPQLPLLRLVSMHS LTPRIQTHFQLAEELRSSGSCQSFTFAQVGSELACSFNELQKKLEVPLAKDSLDA VVTYSFDHIFPGSENNTRTVVLYGDLGSSQFRTYHKLLEKEANAGRIRYILRHQLA 30 KKDKRPVRLSGYGVELHLKSTEYKSQDDAPKPEAGSTSDLANESDVQGFDFK VLKQKHPTLKRALDQLRQRLQGNDEIAQLKAWFQDLGLQAAAAIAEIQGD LQILQYTAHNFPM LARTLLAHKVTDGLRAEVKHNTAFGRSLNVAPPD GALFINGLFFDADTMDLYSLIETLRSEMRVLESLSHNNVRGSLASSLLALDLTASSKKEFAIDI RDTAVQWVNDIENDVQYRRWPSSVMDLLRPTFPGMLRNIRKNVFNLVVVDAL 35 QPTARSVIKLSESVFIHQAPIRLGLVFDARDANEDNLADYVAITCAYNYVSQKKD ARAALSFLTDIYAAVGETKVVTKKDIVKQLTKEFTSLSFAKAEFLEEDSTYDYGR ELAAEFIQRLGFGDKGQPQALLNGVPMPSNVVTADSDFEAAIFTEIMTHTSNLQKA VYKGELTDNDVAIDYLMNQPHVMPRLNQRLSQEDVKYLDINGVAYKNLGNVG VLNRLSNRDMTATLMDNLKYFGGKKSTELIGRASLQFLTIWVFADLETQGRDLL 40 THALDYVQSGESVRVAFIPNTESSASSRRNLNRLVWAAMQSLPPTQATEQVLK WLKKPKIEIPTQLEDILGSTELHLKMLRVYSQRVLGLNKSQRLVIGNGRLYGPL SSDESFD SADFALLARFSSLOYSKVRQVLKESAQDVNEEFNSDTLLKLYASLLPR QTKTRFKLPDLDKTDHSVVKLPKQENLPHFDVAAVLDPASRAAQKLTPILILLRQ VLNCQLNLYLIPVPQHSDMPVKNFYRYVVEPEVQFEANGGRSDGPLAKFSGLPAN 45 PLLTQQLQVPENWLVEAVRAVYDLNKLTDIGGPVHSEFDLEYLLEGHCFDAA SGAPPRGLQLVLGTQSQPTLVDTIVMANLGYFQLKANPGAWSLRLREGKSADIYA	

- ISHIEGTNTHHSAGSSEVQVLITSLRSHVVKLRVSKKPGMQQAELLSDNEQAAQS
GMWNSIASSFGGSANQAATDEDTETINIFSVASGHL YERLLRIMMVSLKHTKSP
VKFWFLKNYLS PQFTDFLPHMASEYNFQYELVQYKWPRWLHQQTEKQRTIWGY
KILFLDVLFP LNVRKIIFVDADAIVRTDIKELYDMDLGGAPYAYTPFCDSRKEMEG
5 FRFWKQGYWRSHLMGRRYHISALYVVDLKRFRKIAAGDRLRGQYQALSQDPNS
LSNLDQDLPNNMIHQVAIKSLPDDWLWCQTWCSDSNFKTAKVIDLCNNPQTKEA
KLTAARIVPEWKDYDAELKTLMSRIEDHENSISRDSA VDDSVDDSVETTTPS
HEPKHGEL
- 10 >16:57:34|GENSCAN_predicted_CDS_5|4647_bpatgttacgtgccgtggcttgtgtgtctgtgtgtctca
tagcactatatacgcgaacttctggggaatccagtcagagctatcccatcaccacgtaatacgcgaatggacgcagacgcc
cctatactctggaatcgcgcgagtagctggccgatgagcagggccctctctgggattacgtttcgggggtgaccaagtgggaca
cggttctcaacgaatatgataccgagtcgcaacagtaaatgccgccttgagctggtcaagagccatgtgagttctcccaattg
cccctgcttaggtggtggtatccatgcatagcttgacgccccggatccagaccacttcagttggccgagggaactgaggagca
15 gtggctctgtcagagctttacttttggccaggtgggttcggaactggcctgcagcttaacgagctgcagaagaagctggaggtgc
cgctcgcaaggatagcttgatgcttctgtgtcacctacagcttgatcacatttccctggcagtgagaacaatacccgactgt
ggtactatacggcgatttgggaagctcgaattccgcacatcacaaactattggaaaagggaagccaatgctggccgattcgtta
catcttgcgtcatcaattggcaagaaggacaagcgaccggtacgacttccggctatggagtggaactccatctgaagtaacg
gaatacaagagtcaggatgatgctcgaagcccgagctggttccacttctgatgaggatttggctaataatcggacgtccagg
20 gctttgattcaaggtgctgaagcagaagcatctacactaagagagcgctggatcaactgcgtcagaggcttctcagggaac
gatgagatcgcccaattgaaagcatgggagttccaggatttgggtctccaggcgccgctgctattgcagaatacagggtgatg
aaacctacaaattctcaatatactgccataatttcccatgttggccagaacctgctggccacaagggttagcgatggttaag
ggcggagggtaaagcataatacggaaagcatttgaagaagctgaatgtagcgctccagatggtgcccctttcatcaatggactct
cttcgatgctgacacaatggatctgtattccctgattgagacgctgcgctcgagatgctgttctcgagagctgcacagtaataat
25 gtgaggggaaagccttccagctcctgctcgccttggatctgacgctccagcaaaaaagaattcgccatcgacatccgtgaca
ctgcagtagcagtggtgaacgatattgaaaacgatgtgcagtagccgagtgccctcatcggtgatggatctttgctccaacct
ttctggcatgttaaggaaatccgaagaatgtgtcaatttggctctagtgtagacgcgctgcagcccacagctagaagtgttat
taaactgtcagagctgttgcacatcaagctccattcgttgggttgggttctgatgcgaggagcgaacgaggataatcttg
cagattacgtagccatcagctgcgcctataactatgtgagtcagaaaaaggatgccgagctgttaagtttctcaccgacatct
30 acgcagcagttggtgagacaaagtgtcacgaaaaagacatagtaagcaactaacgaaggaattacatcattaagcttgc
caaaaggaggagttcctggaggaaagattccagctacgactatggcaggagctgcgagcagagttcattcagcggtgggatt
cggagacaagggacaacctcaggccttgtgaatggtgttccaatcccagcaacgtgtgaccgccgatagcgacttcaggga
ggctatttccaggagattatgaccacaccagcaatctccaaaaggctgtgtacaaagtgaaactgacagacaacgatgtagcca
ttgattatctgatgaatcaacctcagctgatgccagattgaatcagcgaatcctaagccaggagatgtgaaatatcttgatattaac
35 ggctgtgacctacaaaaatcttggcaatgttgagttttaaactgtctgttaaccgggatatgaccgctacgtaattgataatcttaa
atacttgggtggcaagaagtctacggagcttattggccgagcatccctacagttcctaacgatttgggtgttctgatttggaaactg
accagggtcgagatctgtcaccatgccctggactatgtccaaagtggagagagtggtgcgagtcgattcattccaaactga
aagctctccgctcaagccggagggaatctaatcgattgttgggtgccatgcagagcttccaccaactcaagccacggagc
aggttctcaagtggctaaagaaccaaaggagaaaattgagataccactcagctcaggatatacctgggactctacagagctgca
40 cctgaagatgttgagagtttattccagcgagtggttgggtctaaataaatccagcgttgggtcatcggtaatggcggttattggg
ccccttctgctggatgaaagctttagatgcgcccgttgccttggatgccaggttcagttctctacagtatagcgataaggtgctca
ggctctgaaggaaatctgtcaagatgtcaatgaggaaatcaacagcgatacttgccttaagtgtatgccagcctgctcccaggga
aaccaaaactcgctttaaagctaccaacggacttaaaaaccgatcactcggtgtgaaaactaccgcccacagagaaatctcccc
atthtgatgttgcggcgttggatcccgcctcccgagcagctcaaaaactaacgcaatacttatttgcctcgtcaagtgtgaact
45 gccaaattgaactatcctgattccgctccccagcacagcgatagcccggtgaagaactctacagatacgttgggaaccggag
gtccaattcaggcggaatggaggccgatctgatggtccttggccaaattcagtggttgcagccaatcctctgctgaccagca
gctgcaggttcccgagaactgttggctgaagctgtgagagcagtttacgatctggacaacattaagttgaccgatattgtggac
ctgtgcacagcgaattcgatctggagatctgctgttggagggtcactgctttagctgctagcggcgctccgccagaggacttc

112

agttggtgttgggtaccagagtgcaacctaccttgtagatactattgtgatggcgaatttgggttattccaactaaagccaatcca
 ggagcttggccctacgcttgcgtgaaggcaaatcgccgataattatgcaatcagccacattgaaggaacaaatcccacattc
 ggctggctcttctgaagtcaggttctataacctccttgcgatcccatgttgtcaaattaagggtgtctaagaagccaggcatgcag
 caggcggaaactcctgtcagatgacaacgaacaggcagcgcaatcaggcatgtggaacagcatgccagcagtttggcggcgg
 5 cagtccaaccaagcagccactgatgaggatacggaaacctcaacattttctgtggcatcgggacactgtacgaacgtcttct
 aaggatcatgatgtttcgtgctaaagcacacaaaatcacctgtgaagttctgttctgaagaactatcttccgcaatttacgg
 atttcttctcacatggccagtgagtacaactccagtagaattgggtccagtacaaatggccccgctggctgatcagcaaacgg
 aaaaacagaggaccatttggggctacaagatcctttcctggacgtgctcttcccgtgaatgtgaggaaaatcttctgtgatgc
 cgatgccatcgtagaacggatataaaggagttgtatgacatggacctcggaggagcaccctatgcctacacgccattctcgatt
 10 cccgcaaagagatggagggttccgattctggaagcagggtactggcgaagccatctgatgggcaggcgttaccacatttccg
 cctgtacgtgtggacttgaagagattccgaagattcggcaggagataggctaagaggccaataaccaggcacttagccagg
 atccgaacagcttatccaatttggatcaggacttggccaacaacatgatccaccaggctgccatcaaatccctgcccgcagactgg
 ctatgtgtccaaacgtgtgacgacgacgaactcaagactgctaaagtattgtgcaacaacccgagaccaaggagg
 ccaaaactcacggccgccagaggattgtcccgaatggaaggactacgatgccagctgaagaccctgatgtctcgcacgcag
 15 gatcatgagaattcgcatagcagggactcggcagttgatgattcgggtgacgattcgggtgaggtcaccactgtgacgccttctcat
 gagcccaagcacggcgagctgtga

***Drosophila* Gene Hit** rescue sequence and BLASTX with EST and TBLASTN with
 ORF2: UDP-glucose:glycoprotein glucosyltransferase (U20554)
 20 **Human Homologue** BLASTX with UDP-GGT: hypothetical protein (AL133051)
***Drosophila* EST** several including GH16576 (AI293351)

25 **Annotated *Drosophila* genome genomic segment** AE003519
Annotated *Drosophila* genome Complete gene candidate ugtUDP-glucose-glycoprotein
 glucosyltransferase

30 **Human homologue of Complete gene candidate** CG6850-
 IGI_M1_ctg14521_41
 D65BCE6EEC187AE3
 TRANS:SEPT20T.ctg14521.2
 2 FPC_ctg:ctg14521
 FPC_start:1284609
 FPC_end:1284696
 35 FPC_strand:+ (1.20E-215)

Putative function ugtUDP-glucose-glycoprotein glucosyltransferase

40 **Confirmation by RNAi** Only wild type profiles observed

Example 26 (Category 3)

	Line ID	430/5
	Category	Mitotic defects in brain: metaphase arrest (overcondensation, polyploidy, metaphase with bipolar spindle)
5	Reversion	NR
	Map Position	98B5-8
	Rescue ID	2C2E
10	Rescue Sequence	GTGCGGCCCATGGATGTGCGAACGTGTACGAAGACCAAGATCGGCATCGCCA TCGGCGGCAGCACGACGGACGATAACGAAAAAGCTACAGCCGCCGCCACAGA TACAGATGCAGATGCCATGCCGCTGTTATCAGCGCGAGCGGGAGAATGATAA GGGATGGGATCGCTCAGCGCGGCAGGCAAGACGACCAAAAAGAGAGCCAAC 15 TAAATGATGTGCCTAAGACTAAGAGTTTAAATGAGCATTACTGTGCGCACTCT ATGTATTATGAATAAAATTCATACAACTTTGTGGTTTATTATAATAAAAAGT GTGTCAGCTCTACTCGGGGAAAGTAAGTTTACTTCTTGGCCGCTGGCTTCTTG GCGGCGACCTTCTTCTTGCGGGCGGCCAGCAACTTGGCGCGATTGGCGCAGCC TTGGTGGCCACATTGGCGAAGTGCAGCTTGGCCAGCTCGACGTTCTGCTTCTT 20 GGCTTGGCCAGCACCTTGGCCACGGTGCCTTCTCGGCGGCGAGGGCGGCAC GACGCTTGAGTACCTCGGCATAAGGGTTCAACTTGATCAACTTGCGCACGGTT GGTAAGGGGGTT
25	<i>Drosophila</i> EST	several including LD45359 (AI513164)
	Annotated <i>Drosophila</i> genome genomic segment	AE003763
	Annotated <i>Drosophila</i> genome Complete gene candidate	CG5502 RpL1 - Ribosomal protein L1
30	Human homologue of Complete gene candidate	1e-126 432359 dbj BAA04887 (D23660) ribosomal protein [Homo sapiens]
35	Putative function	structural protein of ribosome involved in protein biosynthesis
40	Confirmation by RNAi	Marked decrease in G1 and G2/M indicating fewer cycling cells

Example 27 (Category 3)

	Line ID	472/12
	Category	Mitotic defects in brain: metaphase arrest. Meiotic defects in testis: segregation defects. Abnormal spindles (mitotic: High mitotic index, meiotic: Ab-08/24)
5	Reversion	R?
	Map Position	96C7-9
	Rescue ID	2B6E
10	Rescue Sequence 1	GTCTGACGTTCTCTGAGGGCAAAAGTTTCGAGTTAGTTGAAGGTGAGGGTGCT CGATCACCGATTTGCGGTGAGACGAAAGAAAAGTATGCATTGTTGCGTTGTAA AGAGAGCCGGCGCTCGTCTTGTTACATTGTCGCTGAGAACGTATGTTGTGCT TCATCATTTTCCTTGTTGATTTCCCTTTGACGTGGCAACTTGACCATGTATGACA 15 ACTCTTTGGTGGTGCCATCTGGAAGGCAGAAATTTGATGTCAACGGTGCTCCC AGCCAGTCCACTCCCCAACTCACCTGCAGCTCCACTTCGATATTAACTCGCA ACATATTAGTGGCGTAGTTGTACCTGCCGCGGATCCCATTTCCGCTTTGAAAT TTCGCACTTTCGAATATCCGTCCACATTCGATTTGAGAACATCTTCGAAACGTT CAGCGGTGACCCAATCGGGTATTTTGCCAGCCGCCATTGTAGATAATCGGGAT 20 AAGTATTTTGAAATCGAGCAGAAAACACATATACGTCCAGTGTGACGGTCTTG CGTAGACTGATGAAAGCCGAGTATTAGACTCTACACATCTGTGGAGCTTTTAA ATTCGTAGTGC GCGGCCGATTCTCTCGATCTTCTCTCAAAGCTCCGCTAAT
	Annotated <i>Drosophila</i> genome genomic segment	AE003751
25	Annotated <i>Drosophila</i> genome Complete gene candidate	CG10618 - novel
	Human homologue of Complete gene candidate	none
	Putative function	no homologies which indicate function
30	Confirmation by RNAi	Only wild type profiles observed

Example 28 (Category 3)

	Line ID	571/15
	Category	Mitotic defects in brain: metaphase arrest (overcondensation, few anaphases, some polyploids)
5	Reversion	NR
	Map Position	93D
	Rescue ID	2A8E
10	Rescue Sequence	GGCGGCGCTACATTTGTTGTTGTCGCTGCTGCTCACAGCTCCACCACCATTTGC ACAGTTATATTACCTCGCTCAAGTCGCCCTCTCCCTCTCGCCCACTCGCTGTG TCAATCGAATTAACGAATGCTCTTCGGCGAATAATTGGGTTTAGATACTTT TCCAGCAGACAAAGTTGTATTTTTGCACTTCTTATTGATATTAGGCAAAACGC 15 ATCGGCCGAATCACACGCACACAAAGCACACACGCGAGCAGCGGTTTTTCAA TCTGCAGTACACCAAACAACACACACTATTTCCCTAATGCCTGTTCTTATCCCTC TGATATTCCCAATGAATCGCTGGGCAATTGGCGATTCTGAACCGATTTTCACTT GGCTCTTTGTTTTATTTAATTTTCACCGAAACGCTCTCACACGCAGAGACGCTT TTGCTCGTTTCGCTGATGCTTCTGCTGCAATACACACCACCTACGAAACGAGCC 20 AAGGGAAATTGTATCTATGGGCTGTGTATCTGTTTCTACGCGGCACGCGCTGC ACGTCCGCTCGCTTCGGGTTTTTCGAGAGAGAATATACTTTTTCGATACGGTA CGGTAAACGAATTCCGCGGAATTAATTCTTGAAGACGAAAGGGCCTCGTGATA CGCCTATTTTATAGGGTAATGCATGATAATAATGGGTTCTTAGACGTCA
25	Drosophila EST	LP07504 (AI294185), LP06548 (AI293427)
	Annotated Drosophila genome genomic segment	AE003734
30	Annotated Drosophila genome Complete gene candidate	CG15802 – novel homology to Doom, a product of the Drosophila mod(mdg4) gene, induces apoptosis and binds to baculovirus inhibitor-of- apoptosis proteins
35	Human homologue of Complete gene candidate	none
	Putative function	inducer of apoptosis
	Confirmation by RNAi	Only wild type profiles observed

Example 29 (Category 3)

	Line ID	736/15
	Category	Mitotic defects in brain: prometaphase arrest (overcondensation, fewer anaphases, metaphase with bipolar
5	spindle)	
	Reversion	NR
	Map Position	73C
	Rescue ID	H5E
10	Rescue Sequence	CTAATGAGTAAGGAAAACCAATCAGCCTTGCTAATCGCTTGGCAGTATTGGCT TCTATGCAGGGGGGCGTGTCCCGCGCCCCCTTGAAGCTCAAATTTTGAAGGG CACAGGTCGTCCCCTCCTCCTCCGCGTGGGTGGCGTTCGGCCGAACGAACCGG CGCCTACTTTGCGTCCGGCTAGCGAGGATCTCTGGGTGCCACCCACGGCTGG 15 GTGTTGCGATCTGCCCATTGATAATCCATGCGTGAGAAAGCTTTAGAGAATC TGCCAGATTATTACTCCCCGCATACTCAGAAAAATGTATCCTTCAGATATG TTTATGTTTATGAAGTGAAAAAAGTCCTTTGAAATACTACAAAAAGTGAGGAT CTGACCAATGATTGATTCTATAGAAATATACTATAAACTATAAACTAC
20	Genomic hit, Accession No.	CSC:AC014181
	Annotated <i>Drosophila</i> genome genomic segment	AE003526
	Annotated <i>Drosophila</i> genome Complete gene candidate	CG3971 baldspot - with homology to membrane glycoprotein
25		
	Human homologue of Complete gene candidate	CG3791-9e-08 4680391emb CAB41293.1 (AL034374) dJ483K16.1 30 (novel protein) [Homo sapiens]
	Putative function	membrane protein, function unknown
35	Confirmation by RNAi	Slight reduction of G1 and G2/M peaks indicating fewer cycling cells

Example 30 (Category 3)

	Line ID	82/24
5	Category	Mitotic defects in brain: metaphase arrest (condensation, no polyploidy, no anaphases, metaphase with bipolar spindle)
	Reversion	NR
	Map Position	100D
10	Rescue ID	2E3E
	Rescue Sequence	GGTCAAGCCCGATGGCGTCCAGCGCGGGCTCGTCGGCAAGATCATCGAGCGC TTCGAGCAGAAGGGCTTCAAGCTGGTCGCCCTGAAGTTCACCTGGGTAAAGCGG ATAATTGAATTAGGAAGAAATCAATAGATATACATACGTGGAAACGGGTTC- 15 CCCACGCGGGGTTGCTATCGGACCTAACCTCAAAGGCTGGGTGCAGGCGTCAT CGCGGAATGACATGTGTTAGAGGTCAGAACTGCAATTAAGTATAACGAACC GTTTTGTAACCAGGCCTCCAAGGAGCTGCTGGAGAAGCACTACGCTGATCTGT CCGCCCCGCCCTTCTTCCCCGACTCGTGAAGTACATGAAGTCCGGCCCCGTG GTGCCCATGGTGTGGGAGGGTCTGAATGTGGTCAAGACCGGTCGCCAGATGCT 20 CGGCGCCACCAACCCCGCGACTCGCTGCCCCGCCACCATCCGCGGTGACTTCT GCATTACGGTCGGACGCAACATCATCCACGGCTCCGATGCCGTCGAGTCTGCC GAGAAGGAGATCGCCTGTGGTTCAACGAAAAGGAGCTGGTCACCTGGACCCC GG
25	Genomic hit, Accession No.	CSC:AC012727
	Associated ORF	Genscan ORF1 predicted sequences >16:43:49 GENSCAN_predicted_peptide_7 172_aa MKLLMLGTLAFFSVISATMAANKERTFIMVKPDGVQRGLVGKIIERFEQKGFKLV 30 ALKFTWASKELLEKHYADLSARPPFGLVNYMNSGPVVPVMVWEGLNVVKTGRQ MLGATNPADSLPGTIRGDFCIQVGRNIIHGSDAVESAEKEIALWFNEKELVTWTPA AKDWIYE
	>16:43:49 GENSCAN_predicted_CDS_7 519_bp	atgaagctcctgatgctcggcacaaatgttgctctttctgtaatctcgccgacaatggcggctaacaaggagaggactttcatcat 35 ggtcaagcccgatggcgtccagcgcgggctcgtcggaagatcatcgagcgttcgagcagaagggttcaagctggcgcgc tgaagttcacctgggctccaaggagctgctggagaagcactacgctgatctgtccgcccccttctccccgactcgtgaa ctacatgaactccggcccggtggtgcccattggtggtggagggtcgaatgtggtcaagaccggcgcagatgctcgccgcac caaccccgccgactcgtgcccggcaccatcgcggtgacttctgcatcaggtcggacgcaacatcatccacggctccgatgc 40 cgtcgagtctgccgagaaggagatcgccctgtggttaacgaaaaggagctggtcacctggaccccgccgccaaggactgg atctacgaatag
	Drosophila Gene Hit	rescue sequence and TBLA; abnormal wing disc (awd) (X13107)
45	Human Homologue	BLASTX with awd and TBLASTN with ORF1: tumor metastasis inhibitor nm23-H2 (A49798) non-metastatic cells 2, protein (NM23B) (P22392) and nucleoside diphosphate kinase B.

- Drosophila* EST several including LP05977 (AI257573 similar by TBLASTX to X92956 B.taurus mRNA for nucleoside diphosphate kinase (NBR-A)
- 5 **Annotated *Drosophila* genome genomic segment** AE003779
Annotated *Drosophila* genome Complete gene candidate CG2210 - awd abnormal wing discs nucleoside diphosphate kinase
- 10 **Human homologue of Complete gene candidate** gi4505409
1A5C3F84D7AD272C
[ref]NP_002503.1| non-metastatic cells 2, protein (NM23B) expressed in [Homo sapiens] (1.90E-61)
- 15
- Putative function** human nucleoside diphosphate kinase, transcriptional regulation of c-myc expression.a candidate suppressor of tumor metastasis
- 20 **Confirmation by RNAi** Only wild type profiles observed

CATEGORY 4: ANAPHASE DEFECT**Example 31 (Category 4)**

	Line ID	1132/8
5	Category	Mitotic defects in brain: anaphase defects (overcondensation, high polyploidy, some lagging chromosomes)
	Reversion	?
	Map Position	86F3-6
10	Rescue ID	2C3E
	Rescue Sequence	GGCCGGAGGTACCATTTTGGTAGGACCGTTTTTCGGGCCAACGAAAATACCAC AAGACGGCAGCGATAATAGTGTTTTTTGCTTCAAATGTAGTATGGCTACGCAA CTCACATATGGTTAAGAACTTCGCTGTTTATTTGGTGGTTAACTAGCTAAATA 15 CAATAAGAGTGGCAACGCCGTCACGTTTTCTACATGTATTTTACTTGGCGTAGT GCGCCAAGCTTATAAACCACAGTTGGGCGGTTCTTTTGAATTGTTTAAATTTACA CCCCACTATGAACTTATTAGCCTTCTTTATTTATTTTATATTTTATTTTATTTT AGAATACGTTTACTCAAGGTTTCGCAGCTTGTCAATCAGTATTCGCAAAATATCA ATAATAAAAGGCATCAATTTTCCAATCAGCAGTTGAAAAGAACTCCCCTCGAC 20 ATTTGAACAAAATGCATTTTGGGTGATTATAATTTATTAGAATTTTATTGAC TTAAGGTAAATATAAATAAAATATTATTCAAGTACAAAGGTATATATACTCAT TAATANTATTTGGATTCAAGGAAAATATATTTCAAATGGCGGGGGTTTAATA AAACAATTTTCAAATTAAGG
25	Genomic hit, Accession No.	AC007805
	<i>Drosophila</i> EST	several ESTs including LP09688 (AI295922)
	Annotated <i>Drosophila</i> genome genomic segment	AE003693
30	Annotated <i>Drosophila</i> genome Complete gene candidate	CG6929 - Lk6 kinase
	Human homologue of Complete gene candidate	gi4505191 DB39E49EC0BED990 [ref]NP_003675.1 MAP 35 kinase interacting kinase 1 [Homo sapiens] (6.20E-113) and gi9994197 551A82FA3D09FD58 [ref]NP_060042.1 G protein- 40 coupled receptor kinase 7 [Homo sapiens] (1.70E-106)
	Putative function	Protein kinase associated with microtubules

Confirmation by RNAi
cells

Complete loss of G1 and G2/M indicating fewer cycling

	Line ID	483/19
	Category	Meiotic defects in testis: segregation defects
	Reversion	?
	Map Position	86F
5	Rescue ID	H2S
	Rescue Sequence 1	
		CTCCGGCCACACGGATGAATTCGTCGTCATTTCGTCGGAATCATTCGAACTTTG
		AAAATGGATCGGTAGCTGGGAAGGAACTTAAAGCGAAATACGCAAAGAAA
10		ACGGCTTTTGTCCGCTATTTCAGCGATTTTTTTTTGTGTTGTAATCAGCAGAGGAA
		ATTTTAACGACCAACTCCACCGCCACACCAGCCATCTCCAGCAGCCCCGGAAA
		ATAAAATAGAACTAAATTAACGCCACCATCACTACAACAACCATCTCACCAAC
		AACTACAAGAGCAACAACCACAGCAACAGCACTACTGCACCAAGCCCACAAA
		GAAGAGGTGAAACGCAATAATCGA=CAATACCCGAAGAAAAAACAACAAA
15		ATATCGCAGATAACCGAAAAAAGCGGTGCAATAGATAAAACCCCATTTTTTGCT
		TGAGCTTTTTTCGCCTGTGTGATGAGAGAAATCAGCAGCAGCCATCGATTACA
		ACAACAACAGCAGCCACACCAACGACGACTCACCACCAAACGAAGAATAATA
		ACCAGCGGANAGCGATAGATA
20	Genomic hit, Accession No.	CSC:AC018284
	<i>Drosophila</i> EST	several including GH28825 (AI517767), LP04213

Other results same as 1132/8

Example 32 (Category 4)

	Line ID	1422/14
5	Category	Male and female sterile, small wings, meiotic defects in testis: segregation defects, elongation defect
	Reversion	NR
	Map Position	90B4-8
10	Rescue ID	2F1E
	Rescue Sequence	GGCCAGCTGCTCAAACATTCTGCAGCTATTTGGCCGCCAGCGAGTAGAACGAT ATTGCCAAATATTTTATAATAGTAACCAATACGTTACCAGTATGACCGCGCCG ATAACGATAGAAAATACCACACGGTCTAAAAGTAAATACCATTGTTGGGATTC 15 CCTAATCTTTTGAATTATTTACCGTTAGGTTTCGGTCGTTTTTTTTTGTGTCAGCTG TTCTTTGTATGAAACGGATTAGTAATTTTATTTGTTGTTTTTGTGCATTTTTGCA TATTAAGCCTTGAACATGCCTTAAATCGTTAAAATAGATTATAAGAGGGA TGGACTGTTTGTAAACCAATTGGAAAATTTGTAATCGCTGGTAATAACTAT CGAGATAAGCTTAATTATCGCTGTTTCTTTGTATCTAGTTATAAATAATAATA 20 ATAAACTGGTAATTAACAAAAGTAAAAAGTTACTTAAGTTATACAAAAATAT TTAGTTATTGNATTCAATAATAAGATGGTAATAATAGATGGTAAGATAGTAAT ATTTTAATAATTGAATTCATCACACATGCTGGTGCACGTTCCACAACCTACAA TCAAACGAAA
25	Annotated <i>Drosophila</i> genome genomic segment	AE003718
	Annotated <i>Drosophila</i> genome Complete gene candidate	CG7623 - novel with homology to UDP-galactose transporter.
30	Human homologue of Complete gene candidate	2136348 UDP-galactose transporter related isozyme 3 - human >gi1669564 dbj BAA13527 (1e-36)
35	Putative function	sugar modification protein

Confirmation by RNAi Slightly reduced G2/M

Example 33 (Category 4)

	Line ID	1479/10
	Category	Mitotic defects in brain: anaphase defects (overcondensation, anaphase bridge, metaphase with swollen chromosomes and bipolar spindle)
5	Reversion	NR
	Map Position	69F3-7
	Rescue ID	2D6E
10	Rescue Sequence 1	CCACGGGCAAAATGTGGTCCGGAGGTCCACGACAACGTGCCGCTGACCATATC CCAGATTGAGCGCGCAACTCAGGATCCGGAGAACGAGAATGTGTTTCATCACA GACGACGTGCATCCGATTCACCTTCTGCACCTGCATCATCTACGCCTTTGTAAC GGCAATGGAACGCACAACGAGTCGTTTCATGAAGTTCATGATCGATGATGGCA 15 CCGGCTCCCTGGAGGCCAGCATCACCAAAAAACCCTTCAATGGACGCGTGATC AGCAGCCTGTACAGTGAAGCCAGTTCGCTGGCCTCGTCCGAGGCCTACAAGA GCATTGCCGTGAGCATGATGCGGCTGCTGCAGGTCTCCATGGAGTACATTGAT CCCACGCGCATCTCGAGGGGCCACAGCCTATTCCTGCGCGGTCTCGGAATAG GTTCCGCGGCAAGATGGGTCTGGACGCTTTTCAGTTCTTCATAGACAGCGGCC 20 GATCGCGGAATATGGAAATTGGCTTCGTGGACTACCTAACCGACTGGCAACG AAGGCATAAAACAATGCAAAATAC
	Rescue ID	2D6P
	Rescue Sequence 2	GCCCCGTGGACTTTTCACTCTGTTGATTCTTGCGTATCACGAAGTTATCCAGCTG GCTTTCTATGTCCTCGAAACTCTGATTAATAATCCATTCTATTTGCTTAGTCTGC GATTTCAAAGGGGATTTCTTTATTGCAGTGCATTTTGCATTAGCGCCAAAAAA AAAAAAGTTGTGAGCATGGGCGTAGACTTCGTATTTTCTTACAAATAATATTA ATTAAAATTAATTTTGTGAGCAATTTTCACACAATTGTATTATAAGTTAAACC 30 AGGGTCACATTAATTTGCAGAACCGCGCAATATTTTCTTTTAAACCCCTTACA AATTTTCAGTTGTTTGTGACTACGCCCTGCTAATTTTACTTATTAAATTCAA GTCTAAAAACATTGTCACCAGATAATACGAGTATACTATATGGACAAACGT AAAATCGTTAATAGAATATATATTCAACCATTATTTACCACCGAGAGAAA TTCATTTGCACAAAACGCCAGGTTGGCAGCACCATCATTGCGCACAGCAAGTG 35 GGCAAACTCGTTGTATCGCTTG
	Genomic hit, Accession No. AC007328	
	Associated ORF	
40	Genscan ORF1 predicted sequences >17:42:01 GENSCAN_predicted_peptide_2 1507_aa MKLAPT VKLNNGYEMPILGLGTYNLKKSRCEAAVCHALEMGYRHIDTAYLYRNE GIIGKVLAKLIGDQKLKREQVFLVTKLWDIYHEPKMVKYACDMQLKLLGVVDYID LYLMHSPVGVVDYISDEDLMPHENGQLRTNDVDYVDITYRSMEQLVHLGLVRS LG LSNFNANQLKRLLENCQIKPANLQIECHPELVQVPLIELCKFHNITVVAYSPLGRSQ 45 TCNPLPDYYTDSKLLALAAKYGKTPAQIILRYLSKDNEGEAAVKHAIDVGYRHID TAYFYQNEAEVGKAIRDKIAEGVVKREDIFLVTKLWNIFHDPERVEGICRKQLSNF GLDYIDL YLMHMPVGYKYVDDNTLLPKNEDDVLQLSDVDYLDYKAMEKLVKL	

GLRIEQLAGLSHLSTHSDGMQFRIRMFLTFQRGGPSHNNMQQQQRRGGGSGTDF
 YNQQRDRRDSGRQMDNNYSNNYNNNNNNNNQNRNRGGGNGMQQQRRGGNGGSGG
 GGGNGGGNNPAWNMHRGNQNSNNMMNMRNRGMGSRGPMRPNQVHLLVHTT
 AIDGLLNPGFHILQGYRQSANQNKPRNKIKFEGDFDFEQANNKFEELRSQAKL
 5 KVAEDGAPKPATNATAATATATNEQVGEKVEGVHTLNGETDKKDDSGNETGAG
 EHEPEEDDVAVCYDKTKSFFDNISCEAAQDRSKNKKNDWRQERKLNTEFTGVSS
 TRRGSAVHQLNVFQAVTADATNTTTIMATAALTRDMEERQATTGTIIAVWGGGG
 NFRNRSNNRNNGGGRGGNGMPNITNGNTAAALKAANNAAGHGSNATDSSAPNA
 TTATTKSTSLLEQTQQVAAVSLPVLLPSIGWLFIVMDGPPDIPRSADIAILFVSFEQ
 10 SVLFLKFHCRYNEFAHLLCAMMSFEDIESQLDNFVIRKNQQSEKSTGKCGPEVHD
 NVPLTISQIERATQDPENENVFITDDVHPIHFCTCIYAFVTGNGTHNESFMKFMID
 DGTGSLEASITKKPFNGRVISSLYSEASSLASSEA YKSIAVSMRLLQVSMYIDPT
 RISRGHSLFLRGRPNRFRGKMGVCTNATAPSVSSINRILNRNAAERAAAEFARAAS
 YGYAIHPHTHPYTSFPTWPAHPLWGA VPLATPPGGGPAGAGGALQPGGSGSSY
 15 GSDGNMSSNPSSNSNTTHSNGHNTNSGSGCGDSSAGSGRLSLPALSPDSGSRDS
 RSPDADANRMIDIEGEDSESQSDQPKFRRNRTTFSPEQLDELEKEFDKSHYPCVN
 TREKLAARTALSEARVQVWFSNRRAKWRRHQRVNLIKQRDSPSTSSSPTPLVNPV
 VSPVSPVPVPVAVPESGQKQPYPYSTSNMCNTSSSSSSNSQPCNTINPGSKMSSK
 TSSVSSNQHMEEPA AAVATASPTASAPLSMGGENSAFRALPMTLPMPTLPTASA
 20 AAFALS FARQYIAKTLLGSPPRSQPPTTNQHKPEPNREFLNEACSSAASVQNSTTP
 ATTADTPTAKSAMCVHCEKKGGAMEWM

>17:42:01|GENSCAN_predicted_CDS_2|4524_bp

atgaagctcgctccgactgttaagctaacaatggctacgagatccaattctgggcctaggaacctacaatttaagaagctcgc
 25 tctgagctcgccgtgtgccacgccctgaatggctatcgccatagacaccgcatactgtacaggaatgaaggcattatag
 gcaaggcttagctaaactattggcgaccagaaactgaacgcgaacaggtgttctgtcacaaagctgtgggacataccac
 gaacccaagatggtgaatacgcctgtgatatgaattaaagctactgggcgtgactatagatctatatctgatgcattcgcg
 gtggcggtgactacatctctgatgaagatctgatccccacgagaatggccagctgaggaccaacgatgtgactatgtggac
 acctacagaagtatggagcaactgtgcatctggggctgtgctgcagctgggattgtccaaacttaagccaatcagctgaagag
 30 attactggaactgccaatcaagccggcaaacctacaatagaatgtcatccgaattgtgcaagtcacattaattgagctctg
 taaatttcacaatatcaccgtgttgcctatcgcactggggcgttccaaacctgcaatccgctgcgggattactacactgattcc
 aaactactggcgttggcagcgaatacggcaagacaccagctcaaatcatcctaagatactgtcgaaggacaacgaaggcgaa
 gccgctgtgaacatcgcatgtgtggctatcgcatatagatacggcctatttctacaaaacgaggccgaagtgggcaagg
 cgattcgggacaagatcgacagaaggtgtgtcaagcgagaggatatttttggctactaagcttggaaactttccacgatccag
 35 agcgcgttgagggcatttgcgcaagcagttaagcaattttgcttgactatctgatctgtatctgatgcataccagtggtgcta
 caaatatgtagatgacaacacctgctgccccaaaatgaggacgatgtgtccaactgagcgatgtcgactatctggatactaca
 aagccatggaaaagctgtgtaaaactggcctgctatcgaaactgtgtgacctgagtcatttcaactcattcagatggcatgc
 agtttcggatacggatgttttaacattccaacgtggcgaggaccagccacaacaatatgcagcagcagcagcaacgaggcgggc
 gcagtggaacggacttctataaccagcagcgggatcgtcgggactcgggacgtcaaatggacaacaactatagcaacaactaca
 40 acaacaataataataatcagcgcaatcgcggcgggcggaacggaatgcaacagcagcagcagggagggaacggcgggcagc
 ggcgcgggcggtggaacggaggtggaacaacccggcctggaacatgcatcgggcaaccagaaactgaacaacatgatg
 aacatgcgcaaccgcgcatgggatcccgcgccccatcgacccaatcaggtacacctgctgtgactcacactgctatagat
 ggtttattaaacctggtttcacattttgagggctatcgtccgcagtcggccaataatcagaacaagccggaacaagatcaa
 gttcagggcgacttctgattcagcagggaacaacaaggtcgaggaaactgcgtcccaactggccaagctcaaggtggcgga
 45 ggatgtgtgacccaagccagccaccaatgaacggcgccactgcaactgcaaccaatgagcaggtgggtgagaaggttgaa
 ggcttcacacactgaatggcgagaccgacaagaaggtgattctgcaacgagaccggcgctggagagcacgagcctgagg
 aggatgatgtgtgtgtgtctacgacaagaccaaatgttcttgacaacatctcgtgcgaggtgcccaggatcgacgaagaa
 caagaagaacgattggcgccaggagcgcaagttgaacacggagacctcggagtgctctccacacgacgtggcagtggtgctc

125

atcaactgaatgtattccaagcagttaccgcgacgaaccaatactacaacaataatggcaacggcgccattaactcgggatatg
 gagggagcgccaggctacaacagggaacaattatcgcatgggtggcgggcggaacttcgaaacaggagcaacaatcgc
 aacaacggcgggcggtcgtggcggaacggaatgccaacatcaccaatggcaacacggctgctgcgtgaaggcgccaac
 aatgctgctggccacggatccaatgccacggactccagtgaccaaagccacaaccgacgacaaaagtcgacgtccctcttg
 5 ccagagcagacgcaacaggtggcgagtttctggtgcccgtgtgttaccatcgattggttggttttatcgttatggatggaccac
 cagacattccaagatcggcagatattgcgattctcttctgtagtttgaacaaagtgtacttttctaaatttcacaagcgatacaacg
 agtttgcccacttgctgtgcgcaatgatgagtttcgaggacatagaaagccagctggataacttcgtgatacgcaagaatcaacag
 agtgaagagtcacgggcaaatgtgtccggaggtccacgacaacgtgccgctgaccatattccagattgagcgcgaactca
 ggatccgggagaacgagaatgtgttcacacagacgacgtgcacccgattcacttctgcacctgcacatctacgccttgaactgg
 10 caatggaacgcacaacgagtcgttcacgaagttcatgacgacgtatgacgaccggctccctggaggccagcatcaccacaaacc
 cttaatggacgctgacagcagcctgtacagtgaaagccagttcgtgctcgtccgagggcctacaagagcattgcccgtgagc
 atgatcggtgctgacgagtcctcatggagtagcattgacccacgcgcatctcagggggccacagcctattcctgcgcggtcgtc
 cgaataggttccgcggaagatgggtgtctgcaccaatgccatgctccttcggtgagcagcatcaatcgcatattgcgtaatcga
 gcggcggaaggcgagctgcggaattgtcgcggcgaggttacggctatgccatccatccacacatccgcatccgtacacc
 15 agtttcccacttgccggcgcatcatccgctgtgtgggagccgtgcccctggccacgccacctggtggcgccctgctggagcc
 ggtggtgactgcagccggcgagcagtgatggcagcagctatggcagtgatggcaacatgagctcaatcccaatagcagcaaca
 gcaaccacccacagcaatggccacaataccaacagcggcagtgatggggatagtagtgccggaagtggacgcctctc
 cctgccggcactttcgcggattccggaagttagggacagccgctcccagacgcagatgccaatcgatgatagatatcgaagg
 cgaggacagcgagtcgcaggacagtgaccagccgaagtccggcgcaatcgaccaccttcagtcggagcagctggatgag
 20 ctggagaaggagttcgacaagtgcactatccctgcgtgaatacccgcgagaaactggccgcccggacggcactgagcgagg
 ccaggggtcaggtttgttttcaacagacgagcgaatggcgccgaccagcgggtcaacttgatcaagcagcgcgactcg
 cctcgcacatcgagctcaccacgccgttggtcaatccggtgtcagtcgggtcagtcgaatccagttccagttccagttcagtt
 ccagaatctgccaacagaagcagccatatccgtacagcaccagcaacatgtgcaacaccagcagcagcagcagcaaacgctc
 aaccgtgcaacacatcaatccggcgagcaaatgagcagcaaaaccagcagcgtcagcagcaaccagcacatggaagagc
 25 cagcagcgggcggtggccactgcctcaccacagcatcagctccattatcaatggcggtgagaacagtgcatttcgcgctctgcc
 catgaccttgccgatgccatgaccttgcccacggcatcgggcgcccttcgcgctcagcttgcccgccagtacatagccaa
 gacgcttctcggttctccagatccagatccagccaccaaccaccaaccagcataagcccagccaaatcgcgagttcctcaat
 gaagcctgcagctccgagcatctgtccagaatcgacaacggcggaacaaccgcagatactctacagccaaatcagcaatg
 30 tgcgtgcactgcgagaaaaaggaggggcatggagtgatgtga

30

***Drosophila* Gene Hit** BLASTN with rescue sequence 2: Histone acetyltransferase GCN5
 (AF029776) very small match at end, TBLASTN with ORF1:
 middle domain histone acetyltransferase GCN5 (AF029776).
 Genomic matches histone acetyltransferase

35

Annotated <i>Drosophila</i> genome genomic segment	AE003541
Annotated <i>Drosophila</i> genome Complete gene candidate	CG4107 -Pcaf /GCN5 histone acetyl transferase transcriptional activator protein
Human homologue of Complete gene candidate	gi6382076 72F516F8BD10CD0C [ref]NP_003875.2] p300/CBP- associated factor [Homo sapiens] (1.20E-197)

45

Putative function Transcriptional activator

Confirmation in RNAi Only wild type profiles observed

Example 34 (Category 4)

	Line ID	184/5
	Category	Mitotic defects in brain: Anaphase defects.
5		(overcondensation, aneuploidy, some lagging chromosomes and breaks)
	Reversion	R
	Map Position	71B
10	Rescue ID	C4E
	Rescue Sequence	
		CTCGAGCAGATGTGGGACGAGCTGAGCGGAGCGCACAACTGCCAAGTAAGT
		GGAGCATGTGGATGAAAGGAGTTCCCAGAACAGTGTTGCCAACCAAAAAAAAA
		AAAAAAGTTAAAAAGTTAATTTTAATAGTGTAATAAATATGAATTAAATTAA
15		ATTTTTATGTAAACAGTATTAGCTTTACATGAGATTACCAAATTGTGAGTGTCT
		GTGTTTGTGTTGTCTTTTAAAAACTTTAAAAGCACATAAAGAAATATATTTTAAA
		TTTAATTAATAAAGTTCGTAAAAAGTAAAAGGTAGCTAAATTAAAAAGTTTCCT
		ATTCAAATCAGATTTGGCGAACAAAGAGCCAAGTTGGCAACACTGACAAATGA
		CTCCAAGCGCGAACAAAGCGATTTCTATCGTTATCCCACTCTCTCTCCAGAG
20		ATCGTTCTCAAGGCCAAATGGAAGGGACTTCGAGACAATTTCCGTGTGGAGTC
		AAAAGGATCCGGCGGCCGAATAACGG
	Genomic hit, Accession No. CSC:AC019852	
25	Associated ORF	
		Genscan ORF1 predicted sequences >22:43:26[GENSCAN_predicted_peptide_2 1003_aa
		MAPKKSTIVLNVEQFIHDIERP AIWNRNFHCNKAFLQMWDELSGAHKLPKIVL
		KAKWKGLRDNFRVEYKRIPRADNGDFMVDPATFESKWLHY YALLFLT DHMRHR
		LPKNEQDQSFYFSQSEDCEKTVVEPDLTNGLIRRLQSDDEDYDEEEMEADGEAS
30		EATMEETMPTPPAAHQMNQVSTTPLATGALRAQEEAHQHALIKAGLLRAQLMEL
		EKEAEDLSRKPPPPQMTSPVAPSLQVLVEPPAAHCSPPPMVTTTSAQVQQPGSA
		AVLAPATTTASASSVSSNGAPMGGKRSVSPPLYNKAHHPLATLAAH LAAKDRN
		EDFGPTSAVGGNGDHLSTQHSYANGLIPALKLRPRLSEDSNFNGSSTMDTPLVP
		EDDDYHYLLSLHPYMKQLTAAQKLRI RTKIQLIFKELYKEDLEESNLDREVYVL
35		DDGAEVDLDLGN YERFLDVT LHRDNNITTKIYKLVIEKERTGEYLGKTVQVVP H
		ITDAIQEWVERVAQTPVQGSSKPQVCIVELGGTIGDIEGMPFVEAFRQFQFRVKRE
		NFCLAHVSLVPLPKATGEPKTKPTQSSVRELRGCGLSPDLIVCRSEKPIGLEVKEKI
		SNFCHVGPDQVICI HDLNSIYHVPLLMEQNGVIEYLNERNLQLNIDMSKRTKCLQQ
		WRDLARRTETVRREVCIAVVGKYTKFTDSYASVVKALQHAALAVNRKLELVFIE
40		SCLLEETLHSEPSKYHKEWQKLCDSHGILVPGGFGSRGMEGKIRACQWARENQ
		KPLLIGICLGLQAAVIEFARNKLGLKDANTTEIDPNTANALVIDMPEHHTGQLGGT
		MRLGKRITVFSDGPSVIRQLYGNPKSVQERHRHRYEVNPKYVHLL EEQGMRFVG
		TDVDKTRMEIIELSGHPYFVATQYHPEYLSRPLKSPFPFLGLILASVDRLNQYIQRG
		CRLSPRQLSDASSDEEDSVVGLAGATKSLSSLKIPITPTNGISKSCNGSISTSDSEGA
45		CGGVDP TNGHK

>22:43:26|GENSCAN_predicted_CDS_2|3012_bp
 atggcgccaaaaagtcaccattgtgctcaatgtggagcagtttattcacgacatcgaggagcgcccgccatctggaaccgca
 atttccactgcaacaaggcccttcctcgagcagatgtgggacgagctgagcggagcgcaaaactgccaagatcgtgctcaagg
 ccaaatggaagggaacttcgagacaattccgtgtggagtacaaaaggataccgcggcgataacggtgattttatggtgatcc
 5 gggccacctttgagtcgaagtggctgcactactatgcattgtgttttaactgacacatgcgtcatcgttgccaaaagacgaacagg
 atcagtcattttacttcagccagcaaaagcaggactgtgaaaagacagtgtgtggagccggtttaacaaacgggtctaatacgtcgt
 ctgcaggacagcgacgaggattacgacgaggaggaatggaggcggacggagaggtagcgaagccaccatggaggaaac
 gatgcccacgccaccggctgcgcataaatgaatcaagttagcaccacaccactggccaccggagctttgcgagcccaagaag
 aggcacatcagcagcgttaattaaggcaggattactccgctcagttgatggagctggaaggagggaggacttgagca
 10 gaaagccacctcggccacagcaaatgacatctccagtggcaccctcactacaagtgtagtggaaaccaccagccgcacactgtt
 ctccaccgccaatgtgaccaccacatccgcacaagtacaacaaccgggctcagcagctgttctggcgccggaacgaccaca
 tccgctcatctgtatcctcgaaatggagcgcaatggcggaagagatctgttcgccaccgcctctatacaaaaagcacacc
 atccgctggccactctggcagcagcacatcttggcgcaaaagaccgaaatgaggatttcggaccacctctgtctaggaggaa
 acggagatcacctgagcttactcaacactctacgcaatggactgatacccgcccttaagctgaagcgcccgctctccga
 15 ggatagcaatttaattggttctcgacaatggacactccgctcgtaccagaggacgatgactaccactacttgcagcctacatcc
 gtacatgaagcagctgaccgcagcccagaagctgcgcatacgcaccaagatacaaaagctcatctcaaggaaactctacaaga
 agatcttgaggagtcacaacctagatcgcgagggtttacgttttgacgatggcgccgagggtggaactggaactctggaactatgaa
 ggtttttgagtggtaccctgcatcgggacaacaacataaccaccggaaaaatttacaagttggtcattgagaaggagcgactggc
 gagtacttgggcaaaacgggtcaagttgtccacacatcactgatgccattcaggaatgggtggagcgctggccagacacc
 20 gticagggatcttcaagccacaggtgtgcatcgtggaattgggaggaacgattggtgacatcgaaggcatgcttctgtagagg
 ccttcgctcagtttcagttccgctgaaagagagagaactctgtttggcccatgtgtcgtggttccgttgccaaaggctaccggag
 aaccaagaccaagcccacacaagtctcggtcagagaactgagaggatgtggcctgagtcgccgatttgattgtctgccgatcgga
 gaaaccattggactggaggtcaaggagaagatcagcaactttgtcatgtggggccggatcaggtgatgcatccacgatttga
 actccatttatcatgttccgctgctgatggagcagaatgggtgttgaatacctaaatgagcgccctacagcttaatacgcacatgagc
 25 aagaggaccaaatgcttgcagcaatggcgagattggcgctgaacggagaccgttcgccgtgaagtttgcacgcgcgtcgtg
 ggaaagtacaccaagtctacggattcgtacgcctccgtagttaaagccctgcaacatgccgcctggcagtgaaatcgcaaaactgg
 aactgtctttatcgagtcgtgcctgctggaggaggaactttgcatctgagccgagcaagtaccacaaggagtggcagaagct
 atcgatagccatggcatcctagtccccgggtgattcgggtccgtggaatggagggaagattcgtgcatgccaatggcgcgga
 gagaatcaaaagccattgcttggcatctgcttgggtctgaagcgggcgtcattgaattcgacgaaataaacttggctcaaggat
 30 gcaaacaccacagaatcgatccgaacacagctaagtcccttggatcgtatgccagagcatcacacgggtcaattggcgggc
 actatgcgcttgggcaagcgaataactgttttctgatgtcctagtgtcattcgccagttgatggcaatccgaaaagcgtgcagg
 agcgtcatcggcatcgttacgaggttaatcccaatacgtgcatctgctggaagagcaaggcatgctgatttggggcaccgacgt
 cgacaaaactaggtggaatcattgagtcagcggtcatccctactttgttgcaccaataatcatccagagtactgtcgcggcc
 tctgaagccgtcgcctcttctcggcctgatcctggcctcagtgatcgattgaaccaataatcagcgcggttgcgcctgtcg
 35 ccccgccagctatccgacgcacatcctcgatgaggaggacagtgtgtggcgttggccggagcaacaaaatcgtgagctcctt
 aaaattccattacaccacaatggaatatcaaaaagttgcaatggtagcataagcacttccgacagcgaagggtgcctgcggag
 gcgttgatcctaccaatggccataagtaa

Human Homologue TBLASTN with ORF1: CTP synthase (CTPS) (NM_001905.1)
 40 *Drosophila* EST LD27370 (AA941993)

Annotated *Drosophila* genome genomic segment AE003532
 Annotated *Drosophila* genome Complete gene candidate CG6854 - novel protein,
 possible CTP synthase?

45 Human homologue of Complete gene candidate gi4503133
 C33BD849A0044697
 |ref|NP_001896.1| CTP

synthase; cytidine 5-prime
triphosphate synthetase
[Homo sapiens] (8.40E-217)

5

Putative function

Enzyme important in the biosynthesis of phospholipids and nucleic acids, and plays a key role in cell growth, development, and tumorigenesis. The region of the human gene is the location of breakpoints involved in several tumor types

10

Confirmation by RNAi

Loss of G1 and G2/M peaks indicating fewer cycling cells

Example 35 (Category 4)

- 5 **Line ID** 225/27
 Category Meiotic defects in testis: segregation defects
 Reversion NR
 Map Position 90D
- 10 **Rescue ID** 2D2P
 Rescue Sequence 1
- Rescue ID** 2D2E
- 15 **Rescue Sequence 2**
 GCCTGAACTTAAAACGCTGCCTTCGGCTCTCGCTCGGCACTCGCTCGGCTGCG
 ACGTCGACTGCGACGCTGGCAGCGACAACAACGATTGGCCTCTCTCATTCACT
 TACCTCCTCTCTCTCTCTCGCACTCTCTCTTAGCGGTGAGAGAGTGTTCCTC
 ACATTTGTTTTGCTTTTGCGGTTGCGCAATGGCCCCCAAAACGAAAAGAGCG
 20 CGCAAGAGCTAGCTCCACAGTGGATCCTAAGAGAACGGTCCCTGTGGACTCC
 ATCTAGCTAAGAGAAACGCACTTAGTTAGTTTCTATTTTGGTTGTTAAGTAC
 TGCTAGCTGCCTGCCAGTTGAGTGTCCGTCCAAAAACGGTGGTGGAAATGGGG
 GTGACCACTTCAAACATGAAAGCGAAATGTCCTGAGACCCTACAAAAACTAG
 AAATACGCGGGTGCCTGAGAGAAATTTTTATTTCAAGTAAATTGGCAGAGG
 25 CTACATTTTGAATGTTTACAATGAAAATTGCTGGGGAAGCTAGTGAACAACCA
 TTTCGCCATAATTTACACTATCTAAGCTTTTATTTTATAGCCACATGATATATGC
 ATGCA
- 30 **Genomic hit, Accession No.** AC008361
- Associated ORF**
 Genscan ORF1 predicted sequences >20:36:39|GENSCAN_predicted_peptide_2|515_aa
 MSSTIRLQTSSCQCKLYKYERHPNKPQLPTIPNYPCEILHIDIFALEKRLYLSCI
 35 DKFSKFAKLFLHLSKASVHLRETLVEALHYFTAPKVLVSDNERGLLCPTVLNLYR
 SLDIDLYYAPTQKSEVNGQVERFHSTFLEIYRCLKDELPTFKPVELVHIAVDRYNT
 SVHSVTNRKPADVFFDRSSRVNYQGLTDFRRQTLEDIKGLIEYKQIRGNMARNKN
 RDEPKSYGPGDEVFVANKQIKTKEARFRCEKVQEDNKKNRNGKAAGGKGKTR
 RVARGAQIYQNWAI CRNLFSLACCRVCKVCDIVVEFRKGTNAVNVQIREAI
 40 SHVFHKEDIVIDVQESKEWCIWTDDQVQSPLPELENLWHELWIGPSHAYLIDQIVD
 LFENLLEKYNVQVVDVVRFNFLHRAVVVHSGIIIIIMIGVSGGQRTNAFHHRS
 QRSAIGGDPQQKDSAVQQVQARSSDAFCQIPHRSPRFPGRSQLIPKPNREILRNASA
 TKNLLFRIRSQ
- 45 >20:36:39|GENSCAN_predicted_CDS_2|1548_bp
 atgtccagtagcatccgtctgcaaacttctcatgtcagtggtgcaaactctacaagtagagagacaccctaacaaccaaactta

caacctacgccaaattcctaactacccatgtgaaatacttcacatcgacatitttgcgctcgaaaaagggtatacctaaagttgtattgac
 aaatttagcaagtttgccaaacttttccatctgcagtcacaaagcatctgtgcatttgcgagaaactttggtggaggccctacattacttc
 accgcccctaaggtcttggtttcggataacgagcgagggttgtatgcccacagtgctcaactatctcggctctagatcgcgatct
 gtattatgctccaaccagaagagcgagtaaatggtcaagtcgagagattccactctacgttcctagaaatttatcgttgccftaaa
 5 gatgagctccctacctcaaacccgttgagctggtacacatagcagtgaccgctacaacacttccgttcactcggtaacgaatcg
 aaaaccagcagacgttttttcgaccgctcgtcaagggttaactatcagggtctgacagatttccggcggcagactttagaggacat
 caagggttaattgagtataagcaaattagaggtaatatggctcggataaaaaataggacgagccaaagtcttatggccggga
 gatgaagttttgttgcataaagcaaataaaaaacaaaggaaaaagcgaggttcagatcgaaaaaggtacaggaagacaacaag
 aaaaatcgcaacggaaaagcgccgggttgaaaggggaaaaactcgagagtagcccggtggagctcagatttatcaaaactggg
 10 caatctgccggaatctgtttctgttctgtcttgcctgctgccgagtggtgtaaagtgtgtatagtcgtagaattcagaaaaggaa
 ccaacgcccgtcgtgaacgtgcagatccgtgaagctatcagccatgtgtccataaagaagacatagtcacgatgccaggagtc
 aagggaatggtgtatttggaccgatgcaggtgcagtcgccttgcgcagaacttgagaatctgtggcatgaactgtggataggccc
 tagccatgctgacatgatcagatgtcgcgtcttgcgaaatctgctcgaaaaatataatgtgcaggtgtcgcgatgtgttcggtt
 caatttctccatcgctctcgtagtcgtgatcatttgggtatcatcatcattatcatcatgatcatcgccggttagcggcgcca
 15 aagaacaaatgccttttcacaccaccgatctcagcgatcagcgatcgccggcgaccctcaacaaaagattcagcggtgcaaca
 ggtgcaggcacgatcttcggatgccttttgcagataccccaccgatctccaggttccaggcgcgaccaacttattccgaagc
 caaatcgagaaattctcgaacgcgagtgccacaaaattattgttcgaattcgcagccagtga

- Drosophila* Gene Hit** BLASTN with rescue sequence: couch potato (Z14974).
 20 **Human Homologue** BLASTX with couch potato: RBP-MS/type 2 (RNA binding motif family)(D84108)
- 25 **Annotated *Drosophila* genome genomic segment** AE003720
Annotated *Drosophila* genome Complete gene candidate CG18434 -couch potato RNA binding protein
- 30 **Human homologue of Complete gene candidate** 2224621 dbj|BAA20798| (AB002338) KIAA0340 [Homo sapiens] (2e-19) and Ensembl predicted peptide Gene:ENSG00000070877 Clone:AC009710
 35 Contig:AC009710.00004 (predicted unknown protein)

Putative function Possible RNA binding protein

Example 36 (Category 4)

Line ID 238/37
Category Meiotic defects in testis: segregation defects, multi-stage defects
 5 (P1-02/17)
Reversion ?
Map Position 70D

Rescue ID I7E
 10 **Rescue Sequence**
 GTTCAAACGCACTTTTAAGGTGGCCTATCGGCCCATCAGGAGCAACTTTGTTC
 TGCTGCCGGATCAGTACTACGGCGTCGTTTCGACCTATGTAAGTGTCTAAAGG
 TCTTCGCTCCATTTCAGATTAGATACGCCAAAGATTAATCCGGTCAACCATTCT
 GATTAGGACACGGGCTGCCTGAGCTTGCAGTACAATGGTCGGACGCACTACG
 15 CCTCCTGGGCGCCACAAAAGGGCGGCGGCATTAAAGACACCGAGATTGG
 GATCAATGCCAGAGCGGCCAAGGAGATCGGTAAGCCATTACTTAACGGCCCGG
 ATGTGCATCGGTTGCCAATGTGCCGTAATATTGGACTCCGGCCATCTGCCCCG
 TACCTCGTACGCTAGCAGCACCCACTTACCCTTTCTTGCCGTAGGTCTGCACGA
 GAATGATCTGGTCAAGTGTGCGCTCATCGCTGACGTTCTCAACCTGCGCAGCG
 20 TCCACGTTACCCCCGTCTCGTCCAAGGACTGGGAGATCATAGTGAGTGACGGT
 TTCGCCTGCTTGGCGGCGTGG

Genomic hit, Accession No. CSC:AC017664

25 **Associated ORF**
 Genscan ORF1 predicted sequences >15:26:30|GENSCAN_predicted_peptide_1|1819_aa
 EMVQAKDPPSHYLSKLRTYLDPKASRSHRLYLFYFLCQKRKMVGESTSTQVLRD
 LEISLRTNHIEWVKEFLDDTNQGLDALVDYLSFRLQMMRHEQRLQGVLCASEERL
 NLTNGGDGGEIVMGNSSSVSPGGGGGLLSHGNSTGHGLANGTLDNRQQHTMSYG
 30 FLRPTIADALDSPSLKRRSRHIAKLNMGAAATDDIHVSIMCLRAIMNNKYGFNMVIG
 HREAINCIALSLIHKSLRTKALVLELLAAICLVKGGHEILGSFDNFKDVCQEKRRF
 QTLMEYFMNFEEAFNIDFMVACMQFMNIVVHSVEDMNYRVHLQYEFTALGLDKY
 LERIRLTESEELKVQISAYLDNVFDVAALMEDSETKTSALERVQELEDQLEREIDR
 NSEFLYKYAELESESLTKTEREQLAMIRQKLEBELTVMQRMLOHNEQELKKRDT
 35 LLHTKNMELQTLRSLSRPSASSGDGSLANGGLMAGSTSGAASLTLP PPPPPMPASP
 TASSAAPPPPPPPAPPAPPPPPGFSPLGSPSGSLASTAPSPPHAPPMLSSFQPPPPVA
 GFMPAPDGAMTIKRKVPTKYKLPTLNWIALKPNQVRGTIFNELDDEKIFKQIDFNE
 FEERFKIGIGGALRNGSNGTEVDGSLQSSKRFKRPDNLLEHTRLRNIAISRRLG
 MPIDDVIAAIHSLDLKKLSLENVELLQKMVPTDAEVKSYKEYIIERKDQQLTEED
 40 KFMLQLSRVERISSKLAIMNYMGNFVDSVHLISPQVQSIAGASTSLKQSRKFAVL
 EIVLAFGNYLNSNKRGPAYGFKLQSLDTLIDTKSTDKRSSLLHYIVATIRAKFPELL
 NFESELYGTDKAASVALENVADVQELEKGMDLVRKEAELRVKGAQTHILRDFL
 NNSDKLKKIKSDLRHAQEAFCVEYFGDSSRNADAAFFALIVRFTRAFKQHD
 QENEQRLRLEKAAALAASKKENDQVLMRNKVNQKKQQA VINELKSKAHSVRE
 45 KLLQQDEVYNGALEDILLGLKSEPYRRADAVRRSQRRRIDNNRLSRTLEEMDCL
 HENDLVKCALIADVLNLRSVHVTVPSSKDWEIHELSTEKISGSVLEQTRIVNSTQILI

- VWINKSMQVALTVDRCLKPHMNYGRIDHNTELVVAPNLKGLTNGTSNGVIEENT
 KLSRSKTTAQVKDELTEKLTPLTHSSTVSNVKNTIQRNKRQDHMERLKKDLRRES
 SRSFEFRVIRGLWREQAQESDVFNKGHLPEFFDLDFYCMHTAADKDYVVRVR
 TVEDDIEDDLPETIHPSIELNANLMKLLGIKELERVVLRPKTTVVNFVEKIELFANK
 5 KTHYKIMENAFKRFVIERTQHKPMLFNQEEVVRLEDDLVTVGILPEHFRYCVVD
 AQFLKESKIYAADLVRPVGEIIEETPPTSPLSVQDLIQLPEYDKIVDQVVQELRMN
 LCLSDNSVMRQCNVLLAGASGTGKTVLVERILDQLSRKPDYCHFEFFHGSRSKG
 RKTESIQLDLRNIFTSCLQHAPAIIVLENLDVLAHAAGEQSSQDGEYYNRMADTV
 YQLIVQYTTNNAIAVIATVNELQTLNKRLLSSPRGRHVFQTVARLPNLERADREILR
 10 ELCSHINVAKDDL VKFSNLTEGYRKCDLVQFVERAIFYAYRISKTOPLLNDQLI
 ESLEHTNSYCLQGIQSNQRTGNDADANEMRVEELPGLESVVGVLEEVLMWPSRY
 PTIFNASPLRNQAGVLLYGPPGTGKTYLVSQLATSWNLRIISVKGPELLAKYIGQSE
 ENVRNLFNRARSARPCVLFDEFDSLAPKRGHDSTGVTDTRV
- 15 >15:26:30|GENSCAN_predicted_CDS_1|5457_bp
 gaaatggtgcaggcaaaagatccgccctcacattactgagtaaactgcgcacatatctggacccaaaggcatcaaggagtcac
 ggctttatctcttactttctgtcagaaacggaaaatggctggcagtcacagtcacccagggtctccgcgatctggagatctc
 gctgcgcacgaaccacatcagtggtggaaggagttcctggatgacacgaaccagggtctggacgccctgctgactatctcag
 ctccgactgcagatgatgcgacacgagcagcgccttcagggtgcttctgtgcctcggaggagcgtctgaatctcacaacggc
 20 ggcatggtgcggtgagatagtgatgggaacagtagttctgttagtcctggtggaggtggtggttactatcacatggaacagtac
 gggacatggtctggcaatggcacactgactcaggcagcagcacacaatgtcctatggattcctacgacctaccattgccgat
 gctctggatagtcctagttgaagcgaaggtcacgacatattgccaaattaacatgggtgccgccagggacgacatccatgtgtc
 cattatgtcctgcgagctatcatgaacaataagtagtggttcaacatggttaccagcatcgcgaggccatcaactgcattgccttg
 agtcttatccacaatcgctgaggacgaagccctggtcctggagctgctggcagccatctgtctgtaaggaggagacacgaa
 25 atcattttgggtcgttcgataattttaaggatgtgtgccaggagaagcgcgcttcaaacgctcatggagtagttatgaattcga
 ggcttttaacatagattttatgttgctgcatgcagttcatgaacatcgttgcctcgggtggaggacatgaactacagggtgcac
 ttacagtacgagttacagccctgggttgataagtagtctggagcgaattcattgacagaatcggaggaaactgaaggtgcagat
 atcagcctatttggacaacgtctttgatgttgctgccttgatggaggattccgagacaaaactcagccctggaacgagtgcaaga
 gcttgaggatcaacttgagcagaaatagatcgttaactcagagttcctctataagtagtcggaattagatccgagagtgtaacgt
 30 gaaaacggaacgcgagcagctgctatgattcggcagaagctggaggaggaacttacagtatgcagcgaatgttcagcacaca
 acgagcaggagctgaagaaacgggacacactgctgcacacaaagaacatggagctgcagacgcttgcgttcctgccacga
 tccgctccagcggcgatggttcttgccgaatggtggcctcatgctggttcacatcggggcagcctcttaacattgccacc
 acctcggcccaatgccgcctcgcctactgcaagttcagctgctcctcaccacctccgccagcaccaccggctccacc
 accaccggcggttcagtcgctggcagtcggcggcagcctagcctcgacagcgccatcgccgccacatgccccgcc
 35 atgctaagctccttcaaccgccaccgcctccagtgccggctttatgcccgtcccgatggcgccatgacctcaaacgcaagg
 tgccactaaatacaagttgccacctgaactggatagcactaaagcctaatacaggtacgtgtacaatattcaacgagctggatg
 acgaaaagatcttcaagcaaatcgaattcaatgagttgaggagcgttcaagatcgggattggcggtgcttgcgaatgtagc
 aatggaaccgaggtcgtggtcgtgcagtcagcaaacgcttcaagaggccgacaatgtctcgtctgagcagcacagag
 40 gttagaacaactgcaatctccgtcgcaagctgggtatgccattgatgatgtcatcgccgcaattcatagctggacctgaagaa
 actttccctggagaacgtcagctgctgcaaaaaatggtgccacggatgccgaggtcaaatcctacaaggaatatatcatcgag
 cgcaaggaccaacagctactcaccgaagaagacaagtttatgctgcagttgtcgtgtggagcgtatctcgtccaagctagcca
 ttatgaactatattgggcaatttgcgacagcgttcattatagtcgcaagtgaatcgatagcaggagcgtcgaattccttaaaa
 caatctcgaaaattcaagcggttttgaaattgtcctggttcggcaactatctcaacagcaacaaacggggaccagcctatgg
 45 cttaagctgcaatcgtggacacgctgatcgataaaaatccacagacaagcgatcgtcactgcttactatattgtggccaccat
 acgggccaatttcggagctgctgaacttcgagagcagctgtatggaacagacaaggctgcatcgggtggcactagagaatgt
 ggtggccgatgttcaggagctgaaaggcgatggtgtgcgaaggaggccgagctgcgagtgagggtgccagacg
 catatcctcgtgacttctgaacaacagcgaggacaagctgaagaagatcaagagcgatctcgggcatgcacaggaagcgctc
 aaggagtgctgtgacttctggcgactcctcgcggaatgcagatcgggctgcttcttgcgttgatcgtacgcttcacgagagcg

ttttaagcaacacgatcaggagaacgagcagcgtcttcgcctggaaaaggccgctgcgctggccgcttccaagaaagagaacga
 tcagggtgcttatgcgcaacaagggttaaccagaagaagcaacaggagctgtcataaacgagctgaagagcaaggcgactcgg
 tgcgcgagaaaaagctgctgcagcaggacgaggtgtacaacggagccctggaggacatcctgctcggcctgaagagcagagcc
 gtacaggcgggcgatgctgtgcggcggtcgcagcgccggaggatcgacaataatcgtttatcgcgcaccctggaggaaatgg
 5 attgtctgcacgagaatgatctgggtcaagtgtgcgctcatcgctgacgttctcaacctgcgcagcgtccacgttaccctcgtcgt
 ccaaggactgggagatcatagaacttagcactgaaaagatatcgggcagtgctggaacaaactcgcatagtgaattcaacgca
 gatccttattgtttgattaataagtcgatgcaagttgcgctgacagtggatcgtctgaagccgcacatgaactacgggagaataga
 tcacaatacggaaactcgtggtggcgcccaatctgtacaagggtctgaccaatggaacttcaaatggtttatagaggaaaacaca
 aactctccagaagtaaaaccactgccaggtcaaggtgatgactgactgaaaagtaaacaccgttgaccattctccacggtgtcc
 10 aatgtgaaaaatactattcagcgttaacaagcgtcaggatcacatggagcgtcttaaaaaggacttgcgcgcggaagctcgcgta
 gcttcgaatttcgtgtcattcagaggtctatggcgggagcaggccaggagtcggatgtttgtgaacggaaagcatctgcctgag
 ttctttgatctagatctattctattgatgcacaccgcagccgacaaggattactatgtgagagtgcgcacagtagaagacgatattg
 aggacgatctaccagaacacattcatccatcgatcgaactaaatgccaatctatgaagtgtcgtgggtattaaaggaattggaacgag
 tgggtctaagacctaaataccgtagtttaactttgtagaaaaaattgagctatttgccaacaagaagacgcactacaaaatcatgga
 15 gaacgcatttaagcgatttgtgatagagagaactcagcacaagccgatgctctcaaccaggaggagggtggtcagcgctggaggga
 cgatttactgtgtactgttggaatttaccagaacactttcgttattgctggtggagcgcgagtttctgaaggagtccaagatctacg
 cagcagatctgtgtcgtccggttggcgagattattaaggaggagacgcctccgacatcgccactaagtgttcaggatctcatcca
 gttaccggagtacgataagattgtggatcaggtatgttcagggaattgcgaatgaatctatgcctcagtgctgacaattccgtcatgcgt
 cagtgcaatgtcctactcgtggtgctcgggaacgggtaaaacagttcttggagcgcattttgaccagctgtcacgcaagcc
 20 ggattattgtcacttcgagttcttcacggatcgcgaagcaagggccgaagacggagtcacatccaaaagatcttcgcaacatttt
 taccagctgcctgcagcatgccccgccattgtgtgctagaaaacttgatgtactggccacgcgtcgtggagagcagtcacgtc
 aggatggagagtactacaatcgatggcgatactgtgtatcagttgattgttcagtataccaccaacaacgcatttcagtaatcg
 ccaccgtcaacgagttgcagaccctcaataagcgattgagctcaccaaggggaagacatgtctccagactgttgcgtcgtcgtccc
 aatttggaacgagcagatcgagagataattcttcgagagctgtgcagccatatcaatgtggccaaggacctggatcttgttaagtct
 25 ccaacctcacggagggtaccggaaatgtgatctgttcagttcgtggagcgtgcaatatttatgcttatcgcataagcaagacc
 agcctcttctgaccaatgatcagcttattgagtcctggagcacacaaactcgtactgcctgcagggcattcagagcaatcaaaga
 actggcaatgatgccgatgccaatgaaatgcgctgcgaggagttgcctggcctggagtcagttgtgggagttctggaggaggtcc
 ttatgtggccatcaagggtatccaaccattttaacgcctctccactgcgcaaccaggccggagtacttctatatgggccaccaggaa
 caggtaaaacctatctggtctcagttggccacatcgtggaacctgcgcacatttccgtcaagggtcctgagttgctcgccaaata
 30 tattggtcaaagcgaggaaaatgttcgaaacctgttcaatcgagctcgcagtgcccaccatgtgtgcttttctcgcagagtttgac
 agcttggcgccgaaacgtggtcacgattccacgggggtcaccgatcagtg

Drosophila Gene Hit recue sequence and TBLastn with ORF1: mRNA for l(3)70Da (AJ243811)

35 **Human Homologue** BLASTX with l(3)70Da: peroxisome biogenesis factor 1 (AF026086)

Drosophila EST LD43687 (AI512050)

40 **Annotated Drosophila genome genomic segment** AE003536

Annotated Drosophila genome Complete gene candidate CG6760 mRNA for l(3)70Da
 - novel protein with
 homology to endoplasmic
 45 reticulum ATPases

Human homologue of Complete gene candidate 4505725

ref|NP_000457.1|pPEX1|
peroxisome biogenesis factor
1 >gi|2655141 (AF026086)
(8e-80)

5

- 10 **Putative function** Putative member of the AAA protein family (ATPases associated with diverse cellular activities) including homologies to transitional endoplasmic reticulum atpases, and an E.coli membrane-bound AAA-type metalloprotease which degrades degrades sigma32, an alternative sigma factor for heat shock promoters
- 15 **Confirmation by RNAi** Slight loss of G1, increase in G2/M indicating arrest in G2/M

	Line ID	238/44
	Category	Meiotic defects in testis: segregation defects, multi-stage defects (PI-02/18)
	Reversion	R
5	Map Position	70D
	Rescue ID	F8E
	Rescue Sequence	
10		GTTCAAACGCACTTTTAAGGTGGCCTATCGGCCCATCAGGAGCAACTTTGTTC
		TGCTGCCGGATCAGTACTACGGCGTCGTTTCGACCTATGTAAGTGTCTAAAGG
		TCTTCGCTCCATTTCAGATTAGATACGCCAAAGATTAATCCGGTCAACCATTCT
		GATTAGGACACGGGCTGCCTGAGCTTGACGTACAATGGTCGGACGCACTACG
		CCTCCTGGGCGCCACAAAAGGGCGGCGGCGGCATTAAAGACACCGAGATTGG
15		GATCAATGCCAGAGCGGCCAAGGAGATCGGTAAGCCATTACTTAACGGCCCGG
		ATGTGCATCGGTTGCCAATGTGCCGTAATATTGGACTCCGGCCATTTGCCCCG
		TACCTCGTACGCTAGCAGCACCCACTTACCCTTTCTTGCCGTATGTCTGCACGA
		GAATGATCTGGTCAAAGTGTGCGCT
20		Other results same as for line 238/37
	Line ID	428/5
	Category	Meiotic defects in testis: cytokinesis defects, segregation defects (seg-01/01)
25	Reversion	?
	Map Position	70A
	Rescue ID	G4E
	Rescue Sequence	
30		GTTCAAACGCACTTTTAAGGTGGCCTATCGGCCCATCAGGGAGCAACTTTGTT
		CTGCTGCCGGATCAGTACTACGGCGTCGTTTCGACCTATGTAAGTGTCTAAAG
		GTCTTCGCTCCATTTCAGATTAGATACGCCAAAGATTAATCCGGTCAACCATTCT
		TGATTAGGACACGGGCTGCCTGAGCTTGACGTACAATGGTCGGACGCACTACG
		CCTCCTGGGCGCCACAAAAGGGCGGCGGCGGCATTAAAGACACCGAGATTGG
35		GATCAATGCCAGAGCGGCCAAGGAGATCGGTAAGCCATTACTTAACGGCCCGG
		ATGTGCATCGGTTGCCAATGTGCCGTAATATTGGACTCCGGCCATCTGCCCCG
		TACCTCGTACCTAGCAGCACCCACTTACCCTTTCTTGCCGTAGGTCTGCACGAA
		AATGATCTGGTCAAGTGTGCGCTCATCGCTGACATTCTCAACCTGCGCA
40		Other results same as for line 238/37

	Line ID	848/7
	Category	Mitotic defects in brain: cytokinesis defect. Meiotic defects in testis: cytokinesis defect. Multi-stage defects Polyploidy, no overcondensation PI-01/10
5	Reversion	R
	Map Position	70D1-2
	Rescue ID	G1E
10	Rescue Sequence 1	GGCCACCTTAAAAGTGCCTTTGAACATTCTCGTCGTGGGCGTGTGCGAATTTA GTACGCTCCTTCCTGGTTTAAATCATTTTCGCACTAAACTTCTGCTCTCAGCGG AATTTACTTTTGCTTTATTAGAGATGGGAGCTCGCGCATCAGCTGAGCCGATA CTTGCGCAACAGGTGATACAGCTGATTAGAGATGGCCCTTTTCAACTGTTCCC 15 AGCAGTGACCGCTGCCATAACCGTTTTCAAATTTACGTGAGAACAGACATAA AATAAATATTACAGCTCGTAGTAAATGTTATTCTATATTTAAAAGGAAATTGT AATAGTTAAAACTTGCAATGAATCAGTTACGTTCAAAAAAGGAAACACACTTT AGTTTTTGGCTAGTTTATTGGGTTAATAATTTTATTTAAAATAGTTCGAGTG TTCAATATAGTCATGTAAATGTGTACAGAAAGATCCGGCATTGATATTTAAT 20 ATATCGATTTCCCTTCACTTTTCGCTCCTCGTATACCATGCTGGGGTCTTATCAA ATTTATT
	Rescue ID	G1P
	Rescue Sequence 2	AAGGTGGCCTATCGGCCCATCAGGAAGCAACTTTGTTCTGCTGCCGGATCAGT ACTACGGCGTCGTTTCGACCTATGTAAGTGTCTAAAGGTCTTCGCTCCATTCAA ATTAGATACGCCAAAGATTAAATCCGGTCAACCATTCTGATTAGGACACGGGCT GCCTGAGCTTGCAGTACAATGGTCGGACGCACTACGCCTCCTGGGCGCCACAA AAGGGCGGGCGGCGGCATTAAAGACACCGAGATTGGGATCAATGCCACAGCGG 30 CCAAGGAGATCGGTAAGCCATTACTTAACGGCCGGATGTGCATCGGTTGCCAA TGTGCCGTAATATTGGACTCCGGCCATCTGCCCCGTACCTCGTACGCTAGCAG CACCCACTTACCCTTTCTTGCCGTAGGTCTGCACGAAAAATGATCTGGTCAAG TGTGCGCCTCATCGCTGACGTTCTCAACCTGCACAGCGTCCACGTTACCCCGT CTCGTCCAA
35	Other results same as for line 238/37	

Example 37 (Category 4)

	Line ID	252/40
	Category	Meiotic defects in testis: segregation defects, abnormal spindles. (Ab-03/30)
5	Reversion	R
	Map Position	84E
	Rescue ID	A4B
10	Rescue Sequence 1	TACATGACTCTGCGATTTGACAAAAACAAAATTGAGTTTTGTCAAGAAAATCA ACTATTTTTCTGTGTTTAAAAAACCGAAGCCAACAAATCCGACCAAAATGCCT GCCGAAAACCTTGAGGAGCAGGGTCTGGAGAAGAACCCGAACCTGGAGCTGG CCCAGACGAAGTTCCTGCTTACCCTGGCGGAATACAAGCAGGATGCGGCATTG 15 AAGGCGAAGCTTCTGGAGGCGATTGCGACGGAGAATATGGCCCCGTGGGTAC GAGCACATCCTGCTCCGGAACCTCGGCTTGGACCCGTTAGACAAGGATCTTGCC TGCGCGCGAATTGAAGGAAAAACAATCGCGTTTAAGTTGGGAGCCA
	Rescue ID	A4E
20	Rescue Sequence 2	GTCATGTACTACCAGTGTGACCCCAAAGTTATCGATAAATTATACCGCATATT TTAACATTGCCAAAAATACCAGAGCGATGTCCATCAAGATAGCGACGAAATT AGAACAGTGCAATTGCCAATTGGGAATTTGTATTTTAATTTATTTTAAATTCT GAAAGTAATTTTAATTTAAAAAAAACCTTGAGAGCTGTCTAGAAAAGAACTTAT 25 GTTTCATGATAACTTTGTGCGAAGAATTAAGAAATATTTAGTTGTAATAATT GTNTGAATCTATTTTTTTTCCAATAACACGACTTATATTTTTTTTTTAAATATTC CGAGCTAAATCCCAAGAAAGTTAAACTCCAATCTTGGGATTTTGAAGTGCCCC AGAACTCCAAATTAACACTTCCTTTTTTAAATAATTGTTAAGACCCGTATCA CTTATGGTTATATACTGACCTCGAAAGGGCCACACTAAGGGGGGAGTTTGAAA 30 ATTGATTTTCCTGATAAAAATTTTCGCTTGGAAAGCTACAGCATCGTCCACTGTC CATGTTTATATATCCTTATATTTGCCTATAAATATAT
	Genomic hit, Accession No. AC006494	
35	Associated ORF	Genscan: ORF1 predicted sequences >23:00:28 GENSCAN_predicted_peptide_2 389_aa MPAENLEEQGLEKNPNLELAQTKFLLTLAEYKQDAALKAKLLEAIRTENMAPWY EHICSELGWTVDKDLLARMKENNRVEVEQLDAAIEDAEKNLGEMEVREANLKKK EYLCRIGDKAAAETAFRKTYEKTVSLGHRDLIVFHLIRLGLFYLDHDLITRNIDKA 40 KYLIEEGGDWDRNRNLKVYQGVYSVAVRDFKAAATFFLDTVSTFTSYELMDYPT FVRYTVYVAMIALPRNELRDKVIKSEIQEVLHGLPDVKQFLFSLYNCQYENFYV HLAGVEKQLRLDYLIHPHYRYVREMRLGYTQLLESYRSLTLQYMAESFGVTVE YIDQELARFIAAGRLHAKVDRVGGIVETNRPDNKNWQYQATIKQGDLLLNRQKL SRVINI
45	>23:00:28 GENSCAN_predicted_CDS_2 1170_bp	

139

5	<p>atgcctgccgaaaacttgaggagcagggtctggagaagaacccgaacctggagctggccagacgaagttcctgctaccct ggcggaataacaagcaggatgcggcattgaaggcgaagcttctggaggcgattcgacggagaatattggccccgtgtacgag cacatctgctcggaaactcggctggaccgtagacaaggatctgctggcgcgaatgaaggagaacaaccgcgtagaggtggagc agctagatgcggcaatcaggatgcggagaagaatctgggcgagatggaaagtcgcgagcggaatcttaagaagtcagagta cttgtgccgcatcggcgacaaggctccgcagagactgccttccgaagacctacgagaagaccgtttccctgggtcaccgcct ggacatcgtgttccatctgatccgcttgggactgtttaccttgaccacgatctcatcactcgcaacatcgacaaggccaagtatctg atcgaggaaggcggcgattgggaccgacgcaaccggtgaaggctctaccagggtgtttactcgggtggcggtgcgtgactcaag gcggcgccacgttcttctggacaccgtaagcaccttcacctcatcgaactgatggactacccaccttcgtgcgttacaccgtt tacgtggccatgattgccctgccgcgaatgagctgcgcgacaaagtatcaagggtccgaaatccaggaggtgtccatggc ctgcccgcagtgaaacagttcctgttttctgttacaactgccaatatgagaacttctacgtacacctggccggcgtagagaagcaa ttgcgcttgactacctcattcatccctaccgctactacgtgcgcgagatgcgcattctgggctacaccagttgctggagtcg tatcgtccctcaccctgcagtatatggcgagtcgttcggcgtaacagtgaatacattgaccaggagctggcacgctcatcgc cgccggacggctgcatccaagggtggatcgcgttggcggtgttgagaccaatcgccctgacaacaagaactggcagttacc aggcgaccatcaagcagggcgatctgctgtcaaccgcatccagaagttagccgctgataaacatctaa</p>
15	<p>Drosophila Gene Hit BLASTN with rescue sequence 1 and TBLASTN with ORF1: 26S proteasome regulatory complex subunit p42A (AF145308). Human Homologue BLASTX with EST and TBLASTN with ORF1: Hypothetical protein KIAA0107 (D14663). Drosophila EST several including GH17651 (AI387197)</p>
25	<p>Annotated Drosophila genome genomic segment AE003739 Annotated Drosophila genome Complete gene candidate CG5378 - Rpn7 19S proteasome regulatory particle, non-ATPase protein, subunit S10aHuman Homologue</p>
30	<p>Human homologue of Complete gene candidate gi7661914 8843E6684AE91ACD [ref]NP_055629.1 KIAA0107 gene product [Homo sapiens] (3.40E-149)</p>
35	<p>Putative function component of the 19S proteasome regulatory particle</p>
40	<p>Confirmation by RNAi Marked decrease in G1 and G2/M indicating fewer cycling cells</p>

Example 38 (Category 4)

	Line ID	277/7
	Category	Mitotic defects in brain: anaphase defects (weak, higher condensation, some polyploidy, fewer anaphases, polyploids with monopolar spindles)
5	Reversion	?
	Map Position	71B
	Rescue ID	B8E
10	Rescue Sequence	AGTCGGCGCATGCGGAGAGAGAATCGAAAGAGAAAGAGAAGCAAAGAGAGC GACATACAGCAAAAACAATTCAAAAAGAACTGGTGAAGAATACGAAAATAAG ATAATTTTTTAAGGAAGTCGCGCTTTGATCCGTATCCGTTTTAGCGTCCAAGAT TTATATCTTAAATCGGACCTATATTTTGAGGTACAGTGAAGCTTTGATGCGCCA 15 GTCTTATATGAGTTAAAGTTTTAACGATTGAAAGACACCCCTGAGCTGCTCAT TATATTTCAATATTTATAAACAATCTTATATCAGAGCTTGAGAGACTTGTCATGC GCCACAAAATTCCAATTCCAATTCCAATTCCGGAATAATTCACAATAATCTC AATTAACATACGTATTTTATGTTTCGTAATTTTTTAAAAATCCCAGATTCCCCAC AATTGCCATAATAATCTCGATTATGTTATTATACTCTGAGAAGTAGGAGTGTG 20 TGCAAAGACCACAAACAAATCATTAGGGGCGT
	Annotated <i>Drosophila</i> genome genomic segment	AE003584
	Annotated <i>Drosophila</i> genome Complete gene candidate	CG15383 – novel
25	Human homologue of Complete gene candidate	none
	Putative function	No homologies to indicate function
30	Confirmation by RNAi	Slightly increased G1 decreased G2/M indicating arrst in G1

Example 39 (Category 4)

	Line ID	284/4
	Category	Mitotic defects in brain: anaphase defects (overcondensation, polyploidy (with overcondensation), few anaphases, metaphase with bipolar spindle)
5		
	Meiotic	
	Reversion	NR
	Map Position	89B
10		
	Rescue ID	2C6E
	Rescue Sequence	GTCTACCACTAGCTCTTTGTCTTCGCCTTCTAGTCTCTCTCATCTTGGCAGCCC GTTCTAGTGCGCGTATTTTTAGTCGCAACACATTGCCCAATTGCCAGCCGCTA 15 TTTGTGTCGTCCATTTGTTCAATTCATCGGGCTCTTTTCCGATTTCAAGTGGGTGG CATTTAACAATAATCCCTGCGTTCGCTGTCCACGTCCACATTACGATACGTTTA GTGCACGGAAGAAATAAGCGTGTGGTTTCATAATATTAGCTATTGAAAAAA GTTCTTAAATTTAAGCCTCACTCGATTCTGATGCATGAAATATTATTGGATTGT AAATGAGCGTCATGTTTTGGTATACAAATCTCAAAGTAATTTAAAAATTCTCA 20 TCTTACCGTACCTTGAACCACTACCAATCATCTCAGTACAGCATTTTCAGCGAA TTTCTCACTGTGCACTACAATGCCAGGCGGTACAAGCACCTGTATTTATTTATG GTCCGCTGCCGTAATCGACTGCAGTCGCCGCTTCCCTCTCTTTTTGCTACCAA CAACTTGGGGTAGGGCACCTGAACTAGTTTCAAACGGCGGCGGTTCGGCCTTTT CAGCTTTTTCGCATTTGCCATTTTCCCGCGG
25		
	Annotated <i>Drosophila</i> genome genomic segment	AE003711
	Annotated <i>Drosophila</i> genome Complete gene candidate	CG4275 - mor transcription factor involved in chromatin remodelling
30		
	Human homologue of Complete gene candidate	CG4275- 4507081 [ref]NP_003066.1[pSMARCC 2] SWI/SNF related, matrix associated, actin dependent regulator of chromatin, 35 subfamily c, member 2(aa)
	Putative function	Transcription factor, regulator of chromatin
40		
	Confirmation by RNAi	Decrease in G1 and G2/M and increase in polyploidy

Example 40 (Category 4)

Line ID 407/8
Category Meiotic defects in testis: cytokinesis defects
Reversion ?
Map Position 64B1-2

Rescue ID A9E

Rescue Sequence

10 GACTCACCCCTTTCACGCATTTTCATTGGAACGTTTGTTCGTTTATGCACACGC
 GTGTTGACACTTTTCATGAAACGCAGTGCGTGAAAAGTGCATCGCATAAACGC
 AATAAATGTTTGATGGATGCGTTCTGATGGCTTGAAGTCGCCTATTTGGCCGA
 TTTTCGCACGTCCACTCCCGACGGCAACAGAGTCCTGACTGAATCCCGGAGCG
 GAAGGAGTGTGGATAGCCAGGACTGCCAAAGGACACTGCGCACTTTTACTTTT
 15 TCGAAAGCGAAAGCGAAAGTGGTGGGGCCCAGGCCAAAACAANCCCTTGAGT
 TGAAATTGGAACAAAACCGGGACAGGGATGGGAGCCCAGCTCCAACAAACG
 GTTCCGGATTTCCTTGGGAAAGCCACGCCCTGCGCCTGGAAAAGGAATGCCCTC
 CACCTCATTTGTCTCCGTTTTCGCGTATCTCTCCCCCAAATTTCCGTTAAATG
 AAAAACAACTTTGGGTTTTTGGTTTTTAACAATTTCTCCCCATTTGGTTTTNGGG
 20 TTTCCCTTTCCATTTTGGAAATTGGTTTTTAATTAAAT

Genomic hit, Accession No. AC005814 64A6-64B6

Associated ORF

25 Genscan ORF1 predicted sequences >22:57:22|GENSCAN_predicted_peptide_2|524_aa
 MGRRKDKPRVIPEQDARICRAICLCQLTMVLSCVSIVYLSVAIYSPSLKAFKSGFEL
 DPVMCQTVDRQMPNCPWASCGEWCLTKTSGFCPQIHSIVRRNGTDIQLNNCTR
 VTNTSCAMIDLSRLNKFNCNNGTACNNIRGVFNCSNGHCKNMSEFFLCHHKADG
 LTVNSQKDNTKLNNGFECHGVHCTKIKKPFSCDRYCSKITTTNVNTLIMHEDNLIA
 30 ADCENAVAFNQARGSEHGVRIEPFEFWKEDDGNLLTNCA TVTRESNRRITATDCI
 NGTLLEHDTLPAPFMNFTQFWAIYENSTRSVDPEQRYLPNQANLTIYSWKLFINL
 EGCVNTLRGECKDFVARYGNDGDNNTAQSR YQCYYNKDSNVEFVVARYDLDK
 VYRELLVSLIVPIVLFVISSISLCITKSVKVGDDAKMRCVCAGDDSDNDGPFPGPL
 ANKQQDQMYDTDDD VVDLEHQA VDGQELSDHGLPLDNQELIGSTKSLIPSPVGE
 35 SGTSDQIFDQDQEKATTCDVPEKPLVIL

>22:57:22|GENSCAN_predicted_CDS_2|1575_bp

atggggcggcgaaggacaaacgcgggtgattcccgaacaggatgcgcgcacatctgcgcgccatctgcctgtgccagctgac
 catggtgctgtctctgcgtgtccatctctacctaagcgtggccatctactgccctccctaaaggcctcaagtccggcttcgagct
 40 ggatcccgtcatgtgccagacggtggatgccagatgcccaacaactgccctgggcacatctgcggcgagtggtgcctgacca
 agaccagtggcttttccccagatccactcaatagtgcgtgcgaacggcaccgatatccagctgaacaactgcaccagagtcac
 caacacatcgtgcgcatgattgacctgagtcggctgaacaagttcaattgcaacaacggcaccgcctgcaacaatatcagaggc
 gtcttcaactgtccaatggacactgcaagaatatgtcggagtcttctgtgtcaccacaaagccgatggacttacgtcaattcgc
 agaaggataacaccaagctgaatggattcttcgagtgacggggtgactgcaccaagatcaagaagcccttcagctgcgatgc
 45 ctactgttccaagataacaactaccaatgtgaacaccttattatgcacgaggataatctattgccgccgattgtgagaacgcagtg
 gcttcaaccaagcccaggatccgagcacgggtgtcgtatcgaacctttgagtttggaaaggatgatggcaacctgctga

143

ccaactgcgccacagtcacaagagagtcggacaatcgcatcactgccacggactgcataaatggaaccctcctggaacatgaca
 ccttgcccgcctccctcatgaacttcacccagttttggccatctatgagaacagcaccaggctcggatcccagcagagggtac
 ctgcccaccaggccaacctgaccatctacagctggaagaaactgtcatcaacctggagggtcgtgtaacacactcgtggg
 5 gagtgaaggactttgtggctcgtatggcaacgatggcgataacaacaccgccagtcacgctaccagtgtactataacaagg
 actcgaatgtggagttgtggtgcacgctacgattggacaaggtttacaggagcttctagtctcgtgattgtcccattgtgctc
 ttgtgatctcatctatcgttatgtatcatcaccaaatccgtcaaggtgggtgacgatgccaagatgcgctgtttgtccggcga
 tgattcagataatgatggccccttggcccaggactagcaacaagcagcaggatcagatgtacgatacagacgacgatgtagtt
 gacctggagcaccaagcgggtgatggtaagaactatcgaccacggacttccgtggacaaccaagagctaactggtagcac
 caagtcgttgataccaatcagtcggcgagaatccggaactagtgtatcaaatcttgaccaggatcaggagaaagcaactacgt
 10 gcgatgttcccagagaaaccactagtcatactataa

(corresponds to CG15003)

- Annotated *Drosophila* genome genomic segment AE003480
 15 Annotated *Drosophila* genome Complete gene candidate CG15003- novel unknown
 Human homologue of Complete gene candidate none
 20 Putative function No homologies to suggest function
 Confirmation by RNAi Only wild type profiles observed

Example 41 (Category 4)

Line ID 422/28
Category Meiotic defects in testis: segregation defects, multipolar spindles
 5 (Mul-02/22)
Reversion NR
Map Position 68E

Rescue ID 2I4E
 10 **Rescue Sequence**
 TCGTGGACCCTCAAAGNAACGGATTTCTCCAGTTTCTTCAAAGGGTTAATAAA
 CTTTTCGCACGTTTCGCATTTTTATGCTCAATCCGGTTACAAAATGCTGATAAA
 ACCACTTGAACACACGTTTCCGTACTGATAAGGGCCTTTCTTCTTATCTGACC
 TCTGGAATCCGCGGAATTAATTCTTGAAGACGAAAGGGCCTCGTGATACGCC
 15 TATTTTTATAGGTTAATGTCATGATAATAATGGTTTCTTAGACGTCAGGTGGCA
 CTTTTCGGGGAAAATGTGCGCGGAACCCCTATTTGTTTATTTTTCTAAATACATT
 CAAATATGTATCCGCTCATGAGACAATAACCCTGATAAATGCTTCAATAATAT
 TGAAGAAAGGAAGAGTATGAGTATTCAACATTTCCGTGTCGCCCTTATTCCTTT
 TTTGCGGCATTTTGCCTTCCTGTTTTTGTCTACCCAGAAACGCTGGTGAAAGTA
 20 AAAGATGCTGAAGATCAGTTGGGTGCACGAGTGGGTACATCGAACTGGATCT
 CAACAGCGGTAAGATCCTTGAGAGTTTTTCGCCCCGAA

Genomic hit, Accession No. CSC:AC014962

25 **Annotated *Drosophila* genome genomic segment** AE003543
Annotated *Drosophila* genome Complete gene candidate CG5684 (putative
 transcription factor, human
 homolog
 30 **Human homologue of Complete gene candidate** 1e-100 4758946
 ref|NP_004770.1|pPOP2|
 POP2 (yeast homolog)
 >gi|4106061|gb|AAD02685|
 35 (AF053318) CCR4-associated
 regulator of polymerase II
 transcription

40

Putative function Transcription factor

Example 42 (Category 4)

	Line ID	422/5
	Category	Meiotic defects in testis: segregation defects, abnormal spindles (Ab-04/26)
5	Reversion	?
	Map Position	82D
	Rescue ID	B9E
10	Rescue Sequence 1	ATTGGCTCTTGATGGACTACAACGCTACCAAAATGGGGCTTGAGTTGAATTAC CTGTTGGAAGACACAATGCCACCCACGATCAACAATTCGGCGGTAAACAGTG CCGCCGAAAAGCGACCCAGCGGCAAACGGGAGCGCAAGTAAGTGAACAGAT CCCTAAACAGACCCAGATACTCAGACTGATGTGTACCTTGCAGATCCGAGATC 15 ATTTGCCGCGTGAAGTATGGAAACAACCTGCCGGATATACCATTTGATCTGAA GTTTCTGCAGTACCCCTTCGACAGCCACCGCTTCGTGCAGTACAACCCAACGT CGCTAGAGCGTAACTTCAAGTATGACGTGCTGACGGAACACGATTTGGGTGTC ACGGTGGGACCTGATTAACCGGGAGCTCTATCAGGCCGACTCCATGACGCTGC TGGACCCGCCGATGAAAACTGCTGGAGGAGGAGACTCTGACGCCACAGAC 20 TCTGTGCGTTCGCGCCAGCATTCGAGGACGGTGTCTATGGTTGCGAAATCCGA GT
	Rescue ID	B9B
	Rescue Sequence 2	GGCCAAATCTAGAAATCCTCAAATCTGCGCTTGGCAGTGTGACCGTACTTGAC 25 CGGTACGATAATACCTCCGGTAAAAAAATACTATATTTCCGGGGGACTCAAA TGCAACATCCTCATCGTATATAACACAACATCTATTTGAATTTCAATTTCCACAA CTAATATTATGGATAATGCTTTATTATCATTTTCCAAGTTAGCGATAAATCACC CCACAAGCTGAAAAATCAACGTTTAAAAACGATTGATATTTTTTTTAATACTTT 30 TTGGTTTTACTATTTGAATTTTGTATACTTTTAGATTTTACTATTTTAATTTTC GTTTCTTCTAGCTGACTAACGGGTAAAAAAGGATCCGTCGACCTGCAGATCT CTAGAAGCTTGCCTTGCTGGCGTTTTTTCCATAGGCTCCGCCCCCCTGACGAGC ATCACAAAAATCGACGCTCAAGTCAGAGGTGGCGAAACCCGACAGGACTATA AAGATACCAGGCGTTTCCCCCTGGAAGCTCCCTCCGTGCGCTCTCCTGTTCCG 35 ACCTGCCGCTTACCGGATACCTGTCCGCTTCT
	Genomic hit, Accession No. AC008189	
	Associated ORF	
	Genscan ORF1 predicted sequences >15:53:24 GENSCAN_predicted_peptide_3 211_aa 40 MRNANESSGKPKSKFVSNEFHALFSTICSIADSPAVSREKLKIDLAARKIPSASAPK GDSPLERFSRDLFTYLRVCRWGRFSAALFTAELLIVGGIVSSNRTSESSETGNPLA NEPDPLYMKLVDPMVAGESPKRMKDQKDVGLKSTSSSEELRKLPKTRGRQKRFI RNPNYVKANEFYDKMLSSEYVSKRYKDLPPHPGFGADQPPA	
	45 >15:53:24 GENSCAN_predicted_CDS_3 636_bp atgcgcaacgcaaatgaatcgagcggtaaacaaaatcgaaattgtaagcaacgaattccacgcattgtttcaacaattgttcaa	

5 ttgccgattccccggctgctctcgcagaaaaattgaaatcgatttagctgctcggaaaataccttcggcatcagccccaaagg
gattctccactcgcagcgcttttcgcgggatctgttacttacttgcgctccgtttccgctggggctgttttcggcggcactgtttacc
gccgaattgtgatcgtgggtggcatgtgtcttccaacagAACgtcagagctcttctgaacttggaaccacttgcaaacgagccc
gatccattatatgaaactgttgatcccatgtgtagcaggagaatcacctaaaggatgattaaggatcagaagatgtaggcctt
aaatcaactagcagtgcgaagagctccgaaaattgccaaaaacgcgaggctgcacagaagagattcattcggaatccaaactat
gtgaaagctaacgaattctaataagatgttaagcagtgtaatacgtaaagtaaggatcttcgccgcctcatccggga
tttgagcggatcaaccgccagcatga

10

15

Confirmation by RNAi	Almost no G1 and broadened G2/M indicating arrest in G2/M
-----------------------------	---

Example 43 (Category 4)

	Line ID	423/14
5	Category	Meiotic defects in testis: cytokinesis defects, abnormal spindles (Ab-16/13)
	Reversion	R
	Map Position	67B1-10
	Rescue ID	E9E
10	Rescue Sequence	GTTTGGCGTAAAAGCTTCGGCTGTGTTTGGTGCCCAAATTTTCCACTGCTTCT CTTTTGTGTATCTCTTATATCTTGTGCTTTTTTGTGTGTATGTTTTCTCGTTC TTTTTGCACACGCGCTTCGCGTTGCGGGCCAGCTGTTTTGTGTTGATAAGTGGT TACGGTTTGTGTGTGCCAGCGGGTTTCTTAGTCGAACTGCTCGCGATGACTG 15 ATTTTTCACAAGTGACTCAAAAACAGTCGATCGCCCTTTAAGAAAACCCGCT CAACGCACACAAAAGCGGTTTCTCTCTTTTTGTGCTTCTCTTTTCACACTGA CCACACGGAACGAAAAAATGATTACCGACCACACGGAAGAAAAATTTATGT CCAGACGAACTATTTTGTCCAAGTAGCTGATTTGCATAACAATTTAAGCCA CAAGAACTAGATTAATAATTTACATTTAAATACATTATCAAATCCGAAATAT 20 CAATAATTGTAATTTATCCTTACAAAATGTTA
	Genomic hit, Accession No.	CSC:AC020214
	<i>Drosophila</i> EST	several including LP12306 (AI297868)
25	Annotated <i>Drosophila</i> genome genomic segment	AE003552
	Annotated <i>Drosophila</i> genome Complete gene candidate	CG3967 - novel
	Human homologue of Complete gene candidate	none
30	Putative function	No homologies to indicate function
	Confirmation by RNAi	Only wild type profiles observed

Example 44 (Category 4)

	Line ID	427/5
	Category	Mitotic defects in brain: anaphase defects. Meiotic defects in testis: segregation defects, abnormal spindles (mitotic : Overcondensation, lagging chromosomes/less aligned metaphase with bipolar spindles, Meiotic: Ab-06/20)
5	Reversion	?
	Map Position	67B1-5
10	Rescue ID	H4E
	Rescue Sequence	GTACAGCCTGAAGTGATCGTTGTTGTTTGAATCGGTGCTATCGGCGGTTGCGC TTTGTGGGCATCTTTATCCAATTTGCTATGCGCGCTTGCCTTAAATTTTGAAC TGTATTCCAAGGTTGCTTTGGCGGCTATCGATAGTATCGGCATGGTTACATTT TAGTTTTATAACAAGAATTTTACAGGTATTTTGATTATCTGAGCTTAGTTTTAA 15 GCAANAATATTATTGTTAAAAATTTAAAAAGTAAACAAGCTATTTTAACAAGC ATTTAAACAAATAGTATTAATAATAAAAAATATATCGATATGTGTTGCAAAT GTTCGTTCCCTTAGTATTCTCTCATATTTATTTCAAATAAACTGTATAAAATAT CTGAAAAAGCGAACATATTTATTTAATTTTCATCGCAGATATCGATATCACAGC 20 GCTGCTATCGATGGTGTGTCTGTCTGCGCAGTGCCTATCGCTTACCCTGCCATCGCT AACAAAAA

Genomic hit, Accession No. CSC:AC020120

25	Associated ORF	Genscan: ORF2 predicted sequences >22:06:07 GENSCAN_predicted_peptide_7 464_aa MPSEQHTNIKVAVRVRPYNVRELEQKQRSIIKVMDSALLFDPDEEDDEFFQGA KQPYRDITKRMNKKLTMEFDRVFDIDNSNQDLFEECTAPLVDAVLNGYNCSVFV YGATGAGKTFTMLGSEAHPLTYLTMQDLFDKIQAQSDVRKFDVGVSYLEVYNE 30 HVMNLLTKSGPLKLREDNNGVVVSGLCCLPIYSAEELLRMLMLGNSHRTQHPTD ANAESSRSHAIFQVHIRITERKTDTKRTVKLSMIDLAGSERAASKGIGVRFKEGAS INKSLALGNCINKLADGLKHIPYRDSNLTRILKDSLGGNCRTLTMVANVSMSSLT EDTYNTLKYASRAKKIRTTLKQNVLSKMPTEFYVKKIDEVVAENERLKERKNA LEAKATQLERAGNSGFDPLELKTWYSKIDAVYAAARQLQEHVLMRSKIKNINY 35 RQTLKKELEEFRKLKMCVDQRCQESF
----	-----------------------	---

	>22:06:07 GENSCAN_predicted_CDS_7 1395_bp	atgccttcggaacagcatcacgaataaaaagtgccggttcgctacggccgtataatgtccgtgaattggagcaaaaacagcgga gtattataaaggatcatggtcgttcgacgtcgttcgacccgacgaggaggacgatgagttcttcttcagggcgccaagcaac 40 cgtaccgcgacatcaccaagcggtgaacaaaagttgacctggaattcgacagggtattcgatatagacaattccaaccagga tctgttcgaggagtgacgagcgccgctggtcgacgaggtgtaaatggatacaactgctcgttattgtatatggagccactggcg ccggaaaaacattcacaatgctgggcagcgaggctcaccgggtctgacctatctaccatgcaagatccttcgataagatccaa gcgcagagcgacgtgcaagttcgatgtgggggtatctatagaggtgtacaacgaacatgtgatgaatctgtaactaaatc gggccctttaaacttcgcgaggacaacaatggcgtggtggtcagtggtctttgtctcacgccatctacagtgccgaggagctgc 45 taagaatgctgatgctgggaactctcatcgactcagcaccacacagatgccaatgcagagagttccaggtcacatgccatcttc caggtgcacattagatcacggagcgcaagaccgacacaaaagaacggtcaactatccatgatcgtatggcgaggcagtgga gagggcgccagctacgaaagcattggagtgcgattcaaggaaggcgccagcatcaacaaaagtccttagctttgggaaattg
--	---	--

149

cataaacaagctagccgacggcttaaagcacatcccgtaccgcgactcgaacctgacacgcacccctgaaggactcgttgggcgg
 aaattgtcgcacattgatgggtggccaatgtctcgtatgagctcactgacctatgaagatacctacaacacccttaagtacgctagccg
 agctaagaagatacgacgactctgaaacagaatgtcctcaagtcgaagatgccaaccgagttctatgtgaagaagatcgacgag
 5 gtggtagccgagaacgagcgactcaaagagcgcaacaaggcgctggaggccaaggccactcagttggagcgcgcccggcaat
 agtggattcgatccgctggagcttaagacgtggtacagcaagatagacgctgtatatgcggccgcccgagcttcaggagcac
 gtccttggtatgcgtagcaagatcaagaacatcaactaccggcgagacactgaaaaagaactggaggagttcaggaagctgatgt
 gtgtcgaccagcgagtggtgccaggagagttttaa

***Drosophila* Gene Hit** TBLASTN with ORF2: kinesin like protein 67a (U89264)
 10 **Human Homologue** TBLASTN with ORF2: kinesin family member protein KIF3A
 (AF041853)
***Drosophila* EST** GH22018 (AI402731)

Annotated *Drosophila* genome genomic segment AE003552
 15 Annotated *Drosophila* genome Complete gene candidate CG10923 Klp67a -
 motor protein

Human homologue of Complete gene candidate 2e-58 4758646 kinesin family
 protein 3B
 20 >gi|3913958|sp|O15066|KF3B
 _HUMAN KINESIN-LIKE
 PROTEIN KIF3B and also
 predicted peptide
 ENSP00000166696
 25 Gene:ENSG00000073652
 Clone:AC015936
 Contig:AC015936.00023
 6.70E-91 (predicted kinesin?:
 ENST00000166696)
 30

Putative function motor protein involved in cytoskeleton organization and
 biogenesis

35 **Confirmation by RNAi** Almost no G1 and broadened G2/M indicating arrest in
 G2/M

Example 45 (Category 4)

	Line ID	442/3
	Category	Meiotic defects in testis: segregation defects.
5	Reversion	?
	Map Position	70D4-7
	Rescue ID	H7E
	Rescue Sequence	
10	CGCAAGACTGTCTTCGATAGCAGAAGCGTTATTTGGAACATATCGTTTATCG	
	AAACTACAGTTGCTCAATACTGAACTGTCCAGCTTCGAGTAGCTGTGGCTCAA	
	ACCATTTGTTGTCATCGATAAGCAATTGCAATTTTATTTGTTTGCTTAAAAAATT	
	AAAATATAAACTACGAGGATCAAATATACATACATATTCCCAATATGTTAGCG	
	AAAAAACATTTCTGCTCAAAAAAAGATGTTTAAATACAATGTAAGCTGTTCTA	
15	TGCATTGAACAAATTAACACATTGAGAGGTCGCTCTTATAAGTGACATTTCAT	
	ATTTAAATATATTTTAAATATATTCAAATATAGTATAGCAGTATAGCATTCAAAT	
	GTAAGTGTGGTTGGACTATCGCTGTAGTCCAAGAACTGCAGATAGTGTCATC	
	GCTAGCTTTGAAGCATCTCAAAGGAAAAAGGGCGATAATTCTGATTA	
20	Genomic hit, Accession No.	CSC:AC017664
	<i>Drosophila</i> EST	CK02287 (AA141680)
	Annotated <i>Drosophila</i> genome genomic segment	AE003536
25	Annotated <i>Drosophila</i> genome Complete gene candidate	CG6650 - novel transacylase like
	Human homologue of Complete gene candidate	none
30	Putative function	Transacylase
	Confirmation by RNAi	Marked increase in G1 indicating arrest in G1
35		

151

Line ID 473/22
Category Meiotic defects in testis: no division (no meiosis)
Reversion R
5 **Map Position** 70A1-5

Rescue ID 2B7E
Rescue Sequence 1
10 CGCAAGACTGTCTTCGATAGCAGAAGGCGTTATTTTCGGAACATATCGTTTTAT
CGAAACTACAGTTGCTCAATACTGAAGTGTCCAGCTTCGAGTAGCTGTGGCTC
AAACCATTGTTGTCATCGATAAGCAATTGCAATTTTATTTGTTTGCTTAAAAAA
TAAAAATATAAACTACGAGGATCAAATATACATACATATCCCAATATGTTAG
CGAAAAAACATTTCTGCTCAAAAAAAGATGTTTAAATACAATGTAAGCTGTTT
15 TATGCATTGAACAAATTAACACATTGAGAGGTCGCTCTTATAAGTGCACATTT
CAATTTAAATATATTTTAAATATATTCAAATATAGTATAGCAGTATAGCATTCAA
ATGTAAGTGTGGTTGGACTATCGCTGTAGTCCAAGAACTGCAGATAGTGTCA
TCGCTAGCTTTGAAGCATCTCAAAGGAAAAAGGGCGATAATTCTGATAAGAA
AGTTGGCGTAGCCGGAAGGCGGATTGTCACATACAAAATAGTTTGGAAAGCC
CAAAGTGAAG
20
Genomic hit, Accession No. CSC:AC017664
Drosophila EST LD47104 (AI515336), SD03663 (AI532240)

For other results see line 442/3
25

Line ID 670/6
Category Meiotic defects in testis: segregation defects, abnormal spindles (Ab-12/48)
30 **Reversion** ?
Map Position 70C

Rescue ID H7E
Rescue Sequence
35 CGCAAGACTGTCTTCGATAGCAGAAGCGTTATTTTCGGAACATATCGTTTATCG
AAACTACAGTTGCTCAATACTGAAGTGTCCAGCTTCGAGTAGCTGTGGCTCAA
ACCATTGTTGTCATCGATAAGCAATTGCAATTTTATTTGTTTGCTTAAAAAATT
AAAATATAAACTACGAGGATCAAATATACATACATATCCCAATATGTTAGCG
AAAAAACATTTCTGCTCAAAAAAAGATGTTTAAATACAATGTAAGCTGTTCTA
40 TGCATTGAACAAATTAACACATTGAGAGGTCGCTCTTATAAGTGCACATTTCA
ATTTAAATATATTTTAAATATATTCAAATATAGTATAGCAGTATAGCATTCAAAT
GTAAGTGTGGTTGGACTATCGCTGTAGTCCAAGAACTGCAGATAGTGTGTCATC
GCTAGCTTTGAAGCATCTCAAAGGAAAAAGGGCGATAATTCTGATTA
45
Genomic hit, Accession No. CSC:AC017664
Drosophila EST CK02287 (AA141680)

For other results see line 442/3

Example 46 (Category 4)

	Line ID	460/20
5	Category	Meiotic defects in testis: segregation defects, multipolar spindles (mitotic: High polyploids, no diploids, higher mitotic index Meiotic: Mul-02/59)
	Reversion	NR
	Map Position	78A1-4
10	Rescue ID	2B8E
	Rescue Sequence	
	AGCTGGTCCAATTGGAAACGTTAGCTGCTCCAATGGGAGCAGCTGGCGCTCTC	
	TCTTCGATCGCGCTCGCTCTCATCCTCTCTTTAGCTTGTGCCACAGTAGCTG	
	CCGAAGGCAATTTTCATGTGCTCGTGTGTCGACCCCCACTCAGCCCACTTCTG	
15	ATCGGAATCGGGGATTCGGAATCGTGTAAGGCAGCCTTTGAAGGTCCCTTTTC	
	CAGGTGGCGGCCGTATCCTTAAAGTAAACATAGTTCAACTGACTTGGCAGCGC	
	TCCAAATGCGGTGACTTCTTGGCTATGTCATATATACCCCCACTCCCCTCCTGA	
	CTACCCTGCCACGCCCCACGCCCCACCGTCGGCGACGACAATTCCATTAAAAG	
	TTGTACGTTGTCACTTTGCGTTAACTTATCTGTGGAGCATGTTGTGCGATCGCA	
20	TTTTTATTGTCGCCATTGTCTCTCGCTCTCTCCATCGCTCTTTCGCCTGGCTTCC	
	CTACCCTGCCACACAGGGAAGCCTACACACTCTTAAATCATGCACTTGGAAC	
	AAAAAGTGCAAGCATTAACTTTATTTAAACATTCAAGAGCCGCTTCTCTATT	
	TACCATTGAAAATTTAATTTAAATAGAAGAGGCCTTTTCAGAATAATATAAT	
	ACCTTTAAG	
25		
	Genomic hit, Accession No. CSC:AC020460	
	Annotated <i>Drosophila</i> genome genomic segment	AE003592
30	Annotated <i>Drosophila</i> genome Complete gene candidate	CG10588 - novel gene with homology to proteases
	Human homologue of Complete gene candidate	2e-74 4505453 ref NP_002516.1 pNRD1 nardilysin (N-arginine dibasic convertase) >gi 2462488 emb CAA6369
35		
40	Putative function	Novel protease
	Confirmation by RNAi Marked increase in G1 indicating arrest in G1	

Example 47 (Category 4)

Line ID 477/16
Category Meiotic defects in testis: segregation defect.
5 **Reversion** NR?
Map Position 90C5-10

Rescue ID C3E

Rescue Sequence 1

10 CTGTGGACGGTCGTCAATGCGTGAATATTCTTCTATGTGTAAGTGGTGTGCGT
GTATGTAGATTTCTGGTTAAGAAAAGCCCCAAAAACCAAAGCGCCCCGCAAA
ATATATATTGAGTCTTCTTGGCCCAACAACAAATCTGCCGCCGGACTTTCGCC
GGAGGGCGAGTGAAAAATTCAGTTTCTCTCCTCTCGACGATGCACTTTGGAGG
CTGTGTGAGTGTGTGTGCGAGTGAGTGCGTGTGTGTATACATATGCAAATGAT
15 TGGATGTCGAATCCTTGCATCATCATCTTCATAAAACACTTGGCGAAAAAC
CGCAGGAAAACGCAAGCAGCCGAACAAAAAAGAGAGCCTCTCAAGACAAC
GGCAGCGGCCAAAAGTGAACGCGCAACAAACGCGGCCAAGCAGGCGCGGCA
ATTATTTATAAATCTTAAGCCGTTAGCCCCCTCTCTCTCCCACTCACGAAAAG
AAAATAAGTTAAACCAATTGGTGAAGATGATGCCCC

20

Rescue ID C3P

Rescue Sequence 2

GTCCACAGACTGGCTATATATACTAAAAACGAACTCGCGTGAGAAGACAGGG
ACAGGGCAGCAAACCTCGGTATACGAACGGAACGAAATGAAACGATTCAAGTA
25 GTAGTGTATGCAAGTCTTGTCTGTCTGCGCCTGGCGTTCTTTTCTTTCTTTT
TCGATGGTTTTTCGCCAGGCTGGGCGCTGCCAAAACGCTGATACGGCGGCCAC
AATCACACGCGGCTAATCGCCAGTTGGGCCCTGCACAGGCTGCACATACTTTT
CACTATTAATGCGCTGTATTTCACTTATTTTTCGAACAAATTCGCAGCATGACG
AAGAAGCGAGCCTGTACAAGATTAGAGCGGGTAGCACGCACGATAGTATCGA
30 TACGTACGAGTATTTGGCACTGCGATACATTATCGGTGCTCGTTTCGATAGCCC
CCGATAGCTCTAGCACGAAATTTTATCGCTTTATCCATATTTTATACTATTTTT
ATTTATTGGACTTCAATGAATATTTAATTTACGTCTGGGTCGCTTTTTAAATAT
ATATGGTAATCAATAGCTGGCGAATTAGCGATATTTGAGTGTGACGCAAAAAT
GAGTTGCATCGATATCGATTTCTCGCTACTCTGGGACGCCATCTTTATTGCGG

35

Genomic hit, Accession No. AC007810

Associated ORF

Genscan ORF1 predicted sequences >17:48:58|GENSCAN_predicted_peptide_2|349_aa
40 MSRLFILLLLIVTQLSELQAAAFSVRQNRFDVDPDLQTPAPLATSTESSKKPEKAT
SGLLKKCLPCSDGIRCVPIQCPAHVRMESHEKPQICDLPAKGFGYCCETGQNHT
APKPETSPKERRSGFPTILSPAVLDEARRNFEHLMHGVAQIPVRRGFDPFAHGLVF
HSTAKDDLHNFAISNSAIEQVMTTQLFGKKEQVPVEDFITNNVPIKFTETPLAHHC
QPPPVCGNIRSVYRSMDGTCNNPEPQRSWGAAGQPMERMLPPAYEDVPSASPA
45 AICSYIYGIASRLAPVSVVNCCTFAWQLDWTTGMASGECVCVECMPAEWRLGQC
PLLHEASSEMSRLLAKS

>17:48:58|GENSCAN_predicted_CDS_2|1050_bp
atgagtcgcattttattttttgtgtactattgtgacgcaactgagcaggttgaggcggcagcattttctgtgcgcaaaatcgtt
ttgatgaagttcctgatttcgagactcctgcaccttggccacttcactgaatctctaagaaacccgaaaaagctaccagtggtct
5 gctgaaaaaatgccttccctgcagcgtatggtataagatgcgtgccccaaatccagtgtcccgcacacgttcgatgaaagccat
gaaaaagcccaaatgtcgatctccggctggaataatcggtactgctgcgagactggacagaatcacactgtctccaagccg
gagaccttcccaaggagcgtcgatccggatttccaccattctgtcaccgcagtttggatgaggcgcgtcgcaatttcgagca
cttgatgatggattggcagattccggctgcgcgttggttccagatttggccatggcctggtttccactgcagggccaaggat
gaccttcacaacttcgcatatcgaaacagtccattgaacaagtatgaccaccagttgttgggaagaaggagcaggtgccg
10 tagaagatttcacccaacaatgtgccatcaagttcactgagactccgtggcacaccattgccaaccgccccagtttgcggc
aatattcggctgtttatcgagcatggacggcacttgcaataatccagaaccacagagatctctgtggggtgctgtgtgaaccg
atggagcgcgatctgcccccgctatgaagatgttcgctgaccttctctgtgctatatgtagtatatctatggcatcgcatctcg
ctggcgcctgttctgtgtcaattgttgacatttgcatggaattgattggaccactggaatggcgagcggggagtggtgtgtgt
gtggaatgtatccggcgagtggtgttggccaatgccgttgcttcatgaggcgtcgagtgaatgagccgcctcttggcta
15 aaagctag

Drosophila Gene Hit rescue sequence: eyelid/osa (AF053091)
Human Homologue BLASTX with eyelid: KIAA1235 protein (AB033061) Brain
 protein 120 (AB001895)
 20 **Drosophila EST** several including LD04852 (AA201670), LD24466

	Annotated <i>Drosophila</i> genome genomic segment	AE003718
25	Annotated <i>Drosophila</i> genome Complete gene candidate	CG7467 - osa DNA binding putatively involved in DNA packaging
30	Human homologue of Complete gene candidate	CG7467 - 7e-25 2588991 dbj BAA23269 (AB001895) B120 [Homo sapiens] and O14497 SWI/SNF-RELATED, MATRIX-ASSOCIATED, ACTIN-DEPENDENT REGULATOR OF
35		CHROMATIN SUBFAMILY F MEMBER 1 3e-67

40 **Putative function** transcriptional regulator

Confirmation by RNAi Only wild type profiles observed

Example 48 (Category 4)

- Line ID** 496/4
Category Meiotic defects in testis: segregation defects, abnormal spindles
 5 (meiotic: Ab-08/42)
Reversion NR
Map Position 65E4-7
- Rescue ID** 2C1E
 10 **Rescue Sequence**
 GCACGATCGCTCTCTCTTGGCTCTCTCTATCACTCTCTGGACTCTCTCTCAGCA
 CCTTTGCTACCGTTTCGCAGAACAGGTGTATCGGTTTTCAAGGCAACTGTGATT
 TTTTAACCTCAACATTCTATATCGAAAACCTTGTAAGGTCGGAATTTTTCTTGAG
 CGCCTAAAAGTGTGCAGTGAAATCATTAAATCCACTTCCGGTTGCAAAACAGG
 15 AATCACACATATGAAGTGATTAATAATCATAGAAGGTTTGACACCTTCAAATA
 ATAAGAAAACAAAAATTTGTAACTGTGATAATTTATTTAATTGAAATCTTAA
 TTTAATGGCCTACAAATCTGTTGAATATCCGTTGAATACACTTTTCCAGGGTGT
 GTCCTAGTCGGCTCCTCTTTGTTACCCAGTTTGCTGGTCTTCTTAGCCGCACA
 CCAGTTTATCGCTGTTTGCCTTTGCGCTTTTCATTATATAACAAAAACAATG
 20 TTATTGTTATTGCGGTGGCTGTAGATGTAAATGTAAATGTAGATGTAGAGGCT
 GCTTCTTGGG
- Genomic hit, Accession No.** CSC:AC018039
- 25 **Associated ORF**
 Genscan ORF1 predicted sequences >19:35:36|GENSCAN_predicted_peptide_6|190_aa
 MVSEQFNAAEKVKSLTKRPSDDEFLQLYALFKQASVGDNDTAKPGLLDLKGKA
 KWEAWNKKQKGSSEAAQQEYITFVEGLVAKYDNGMHKQEPNTCQARNATRFR
 KSSECSLDQNTYTSSVTVIPAFHEGPKNSTASWPRIYRCYQRNQQAAANCKWANTN
 30 SVCCKPHGKQSRRIIFAEFLAGHTVQILG
- >19:35:36|GENSCAN_predicted_CDS_6|573_bp
 atggtttccgagcaattcaacgccgccgagagggtgaagagcctgaccaagcgtcccagtgatgacgagttcctgcagctg
 tacgccctgttcaagcaggccagcgttggtgacaacgacaccgccaagccgggtctcctggacctgaagggcaaggccaagtg
 35 ggaggcctggaacaagcagaagggaagagcagcaggccgccagcaggagtacatcacctttgtggaggccctgttgcc
 caagtatgacaatggaatgcacaacaagaacaaacacttgccaagcacgcaatgcgactcggttcggaaaagctcggaatg
 ctgctggatcagaatacgtatacgtccagtgtagcggttatacctgcattccacgaaggtccaaagaactcgacggcaagttggc
 caagaatttaccggtgtatcagcgaaccaacaagcggccaactgcaagtgggcaaacacaaatagcgttcgagggaaccc
 caccgaaaacagagccgccgaatcatttcgcagaatttctggccggccatacggcgagattcttggttaa
 40
- Drosophila Gene Hit** rescue sequence: melt (S144114) P element insertion site
 (AF174669), TBLASTN with ORF1: diazepam binding inhibitor
 (DBI) (U04823) and melted (AF205831)
- 45 **Annotated Drosophila genome genomic segment** AE003560
Annotated Drosophila genome Complete gene candidate CG8624 melt - putative signal

		transduction protein CG8631 msl-3 - acyl-CoA- binding protein/diazepam binding inhibitor
5	Human homologue of Complete gene candidate	CG8624- predicted gene ENSP00000065899 Gene:ENSG00000055889 Clone:AC015904 Contig:AC015904.00014 1.70E-15 (unknown predicted gene 1: ENST00000065899 and AK022666 Homo sapiens cDNA FLJ12604 fis 2e-29
10		
15		CG8631- gi5803104 0C85AE40FDF874CD [ref]NP_006791.1 male- specific lethal-3 (Drosophila)- like 1 [Homo sapiens] (1.70E- 36) and Ensembl predicted peptide ENSP0000006617 Gene:ENSG0000005302 Clone:AC004554 Contig:AC004554.00001 8.70E-19 (unknown predicted gene 1: ENST0000006617
20		
25		
30	Putative function	CG8624: putative signal transduction protein CG8631:acyl-CoA-binding protein/diazepam binding inhibitor
35	Confirmation by RNAi	CG8624: reduced G1 and G2/M Indicating fewer cycling cells, CG8631: Increased G1 to G2/M ratio indicating arrest in G1

Example 49 (Category 4)

	Line ID	523/19
5	Category	Female sterile. Meiotic defects in testis: cytokinesis defects, segregation defects (Mitotic: Less condensed chromosomes, nuclear bridges, Meiotic: Seg-01/02
	Reversion	R
	Map Position	75C1-4
10	Rescue ID	2B4E
	Rescue Sequence	
	ACTGAGAGCATATTTGTGCACCAGAGGGCTGCATAACAACATTCTCTTTGTCC	
	ATTCGTTATACTTCGTATTCAGAATACATGTCATTCAGTTGGTCCCGTTCTTTTT	
	GCGTTCACTTCGTATATAATTCGGCGATCGAAATGAACTAACTGAATGTGTTCA	
15	AAGAATGAATGAAGCCAATGAATTTTCAATAGTAATTCAGAGTGCTTAAATTT	
	CTTCATGTTGTCATTGAGTAAATGAGTTCGGACAGCGCGAAGGTAAGTCGAA	
	GTTTGTGTTTTATTATGTTTATTTGTATTATTATGTACACTAGTCGGCATACTTT	
	TGCGTGCGTCTTATACGTGTGCGTCTTATTTAACAATATTGTAAAATAAAATAT	
	ATAAATTATTTGTTATATGCGTAGGGGCCTTTATTTTGTGTATTGATAGTCTTTT	
20	GTCATAGATATCATTATTCTGACAAGATTTGAACTTTTCAAGTTATTGCCTCTC	
	GTTATTCAATTCCTAGCTGGTCTTACGTTACGCGATATTTCTAAAATATCCTA	
	AAATCGCACAAAACAGTCACGCCCACACTTTTGAAAAACGTGGTAATATTTT	
	CATACTTGCATTAAGTCTGG	
25	Genomic hit, Accession No. AC007691	
	Annotated <i>Drosophila</i> genome genomic segment	AE003520
	Annotated <i>Drosophila</i> genome Complete gene candidate	CG4306 – novel
30	Human homologue of Complete gene candidate	
	4e-25 3242764 (AC005154)	
	similar to protein U28928	
	(PID:g861306) [Homo	
	sapiens]	
35	Putative function No homologies to indicate function	
	Confirmation by RNAi Only wild type profile observed	

Example 50 (Category 4)

	Line ID	666/19
	Category	Mitotic defects in brain: anaphase defects (weak, overcondensation, aneuploidy, lagging chromosomes, metaphase with bipolar spindle)
5	Reversion	NR
	Map Position	64E1-5
10	Rescue ID	I9E
	Rescue Sequence	CCCTCGTCTACGTCGAAATTCTGGATGCTTCTCGGATTTAGGGTTGTATCTCGA AAACGTTTGACTGCGAATGTCAATATCGATATGCTAACCGATAGCTGTTCGATG TTTTAAACACAGTGCAGTGTITTTAAATCGCTCCCCATTTATATATATTTGTGC 15 NTGCTTTTGGCGGTNNTTTCTTTATATGCTTCTTATGCTTTTACGATTATTATT AGCGCTTATTTGATTGCAAAATGCCAAGGAAAGCGTGACTGTGATGGCGAAATG CGGAAAGTACTCCTTAAATCTCATATATCGCATAAAACTATCGGTTCTGGAAT GTTTCGTGTAAGTCTGCGAAGATAGAGATCGATCTATTTTGAGGATACATTTG TTAATATTATAAGGGATTCTTCTACAGGGGTCAGATTGCTTAAAAACACACAG 20 AANAATAAACAAAAATTTCTTTGAAATATTGAAATATTTGAAATANAAAAAA CGTATTGACGAGGTAAGCATATTGAAAAAGATAGGAAGGTGATGGAGAAAAGT GCACTTATATTGGTCCACAAAGAGCTTATAATCAAAAGATCAATAGATATAAA TATCTTTATATGATATAAAATATAATACATATAATATAATATCATATACAATG GATAAATTGCAAGTGGCAAAATGAATTCCGCGGAATTAATTCTGAANCGAAA 25 GGGCCT

Genomic hit, Accession No. CSC:AC014815

Associated ORF

30	Genscan ORF1 predicted sequences >17:46:43 GENSCAN_predicted_peptide_1 334_aa MGKDFYKILGLERKASDDEIKKAYRKLALKYHPDKNKSPQAEERFKEIAEAYEVL SDKKKRDIFDNYGEDGLKGGQPGPDGGGQPGAYTYQFHGDPRATFAQFFGSSDP FGAFFTGGDNMFSGGQGGNTNEIFWNIGGDDMFAFNAQAPSRKRQQDPPIEHDLF VSLEEVDKGCIKKMKISRMATGSNGPYKEEKVLRITVKPGWKAGTKITFPQEGDS 35 APNKT PADIVFIIRDKPHSLFKREGIDLYTAQISLKQALCGALVSVPTLQGSRIQV NPNHEIHKPTTTRRINGLGLVPKPSRRGDLIVSFDIKFPDTLAPSLQNQLSELLPN
----	--

>17:46:43|GENSCAN_predicted_CDS_1|1005_bp

40	atgggcaaagacttctcaagattctgggcctcgagcgcaaggccagcgacgatgagatcaagaaggcctaccgcaaactggc actcaataccatcccgacaagaacaagagcccacagggcggaggagcgcttcaaggagatcgccgaggcgtagcaggtgctg tcggacaaaaaagcgcgacatctcgacaattacggtaggatggattgaaggcgacagccgggaccagatggcggcg gtcagccgggagcgtagacttaccagttccacggcgatccgaggccacatttcccagttcttggatcgtagcgttggc gcgttcttaccggcgccgataacatgttagtgcggtcaggcgccgaataccaacgagatcttctggaacattggcggcgacg atatgtttgccttaatgccaggcaccagtcgcaagcgccagcaggatccgcccacgagcatgatctgttcgtgctggag 45 gaagtggacaagggatgcatcaagaagatgaaatctcacgatggccaccggaagcaatgggccgtacaaggaggagaag gtgctgaggatcacagtgaagccgggctggaaggccggtaccaagattaccttccccaagagggtattcgcgccaaacaa
----	---

159

gacgccagctgacatcgtcttcattcgcgacaaaccgcattcgtgttcaaacgcgaggggaatcgatctaaagtatacagccc
 agatcagctgaagcaggccttgtgcggagcactggttagtgcccacgctgcagggcagcaggatacaggtgaatccgaacc
 acgagatcatcaagcccaccacaacgcgcggatcaacggactgggtctgccggtgcccaaggagccatcgaggcgcggcg
 atctgacgtctccttcgacattaagtttccgacacactggcaccagctcgcagaatcagctgtccgagctgtgccaactag

5

Drosophila Gene Hit rescue sequence: fasciclin I (FasI) (M32311) TBLASTN with
 ORF1: DnaJ homolog (DROJ1) (U34904)

Human Homologue TBLASTN with ORF1: DnaJ-like heat shock protein 40 (HLJ1)
 (U40992.2)

10

Annotated Drosophila genome genomic segment AE003565

Annotated Drosophila genome Complete gene candidate CG10578 - DnaJ-1 a
 chaperone putatively involved
 in protein folding. Stimulates
 activity of HSP70

15

Human homologue of Complete gene candidate 8e-94 1706473 P25685
 DNJ1_HUMAN DNAJ
 PROTEIN HOMOLOG 1
 (HDJ-1) (HEAT SHOCK
 PROTEIN 40) (HSP40)

20

Putative function Chaperone involved in protein folding

25

Confirmation by RNAi Almost no G1 peak, increased G2/M indicating G2/M arrest

Example 51 (Category 4)

	Line ID	714/11
	Category	Meiotic defects in testis: cytokinesis defects, abnormal spindles (Ab-01/04)
5	Reversion	?
	Map Position	66A10-15
	Rescue ID	2A4E
10	Rescue Sequence	AACCAGAACGAAACTCCAATGCAGTTTCATTTTGTCTAGTTTAATCATTAAACA AAGAACTGCGCAACCGATCGCAACTAGCTCGTGGACTCTTGTTCTCCCAATAA TTGGTATGTTTTCCATTTTTCGTTAACATGGAAAATGTGTGAAAAGCTTTTCC CCCTCCAAAAGAAGCGTACTGAACTAAGCTTTCGGTGGTTAGTAATAGTAGTC 15 GTTATATCTTATTTTTCTTATTTACGTGCAGCTGCAATCATTGGCTGCGTCACTT TGGCGTCAGCTATAAACTGGTGGATCAACTCGGCGGCCTCCAAAAGCTGCGCA TCTGCTCCAGACACTTTAGCCAACGCCAGGAGATGGCCAAAACCCGCATCAA GATGACGCCGCTGCGCAAGTCCTCGTCCTCAAAGGGCATTGTGCTACCCATTA ATGCCGCTGGAGGGTTCGGTCATTGCAGGCGCCTTAGCACGAGGAGGAGGTGC 20 A
	Genomic hit, Accession No.	AC012390
	Associated ORF	25 Genscan ORF1 predicted sequences >19:47:45 GENSCAN_predicted_peptide_2 711_aa MRSHQAVGNLLLADEALPAVQSASVYVWMAEQPLSPGQSYDIKIADSPSVSS KSITDNGADVQWFAFEHSQYYQGVQQMFLSALERIDSEFLITLIKRCPYHVDLSLVQ LSEVCKMTEDFSLASELLERALLLLESSLHINFSLTSGNCRLDYRRQENRSFYIVLF KHAQYLEERACSRTAFEISKLLLSLQPDTPDLAMILPNQPDQCTGNMTQLQQAGK 30 IRKRSEKQFPIGTEPRGTDALRFTLQTLASAGRDITWNIKRLQGSRTVGAAGGYLI DKKTAVQYKITHLKDPNIDQLFDSSGDGKADLHGSTPDWGCQAMMADAISR YKEGNPVFYTTWTPYWVSNELKPGKD VVWLQVPFSALPGDKNADTKLPNAGGI EGLIADEEVQVLDALCDAPCVGVSHSCRLLDGNRRGNNELRLFIPGKSQFGVADG CADKQSVMEYHAAKTGHTKFSESEEEKKALTEEEKKAQLALIEEKLKQKRIEREE 35 REKIEALQREKNRIKSGKDMTEAKRRMEELEMKKIVEQRKREKDEEKAARDRVK AQIEADKAARKAREQKELGNAEPAPSVSSTTVSSPPAGVKSPPRDYTETRIQGASA ILAAAAPYYQPPAVPQDVQPDPIGYGAFGVVCGSHISGWHCSAGHYEDGNENFE CLKTFSTSDRIGCEWRWAAATVLAATCISPNGRCGHYKRVRRRIKTNITTT 40 >19:47:45 GENSCAN_predicted_CDS_2 2136_bp atgagatgcgcatcaagccgttgcaatctgctgctggcggcagacgaagcgttaccggcggtgcagagcgctgctgtatgtg gtatggatggcggacagccgcttctcaggcagagttacgacatcaaaattgccgactctccatcggtgtcctccaagtctatc acagataatggagcgggacgttcaatggttgcctttgagcatagccaatactaccaggagtgagcagaaatgttcctttctgctctg agcgcattgactcggaatttctgatcacacttatcaaacgctgccctatcatgctgactccttggttcaactcagcgaagtatgcaa 45 gatgaccgaagacttttcttgccctccgaactgctgagcgcgcccttctccttctggaatcgtcgtgcacatcaacttcagttga cgtcgggcaactgccgactggactaccggagacaggaaaaccgatccttctacatcgtgctgttcaagcacgcgcagtagctgg

161

aggaacgagcttgagccgcaccgccttcgagatctccaaactgctcctgagcttcagccagacacagatcctcttgccatgatt
ctaccaaatcagccggatcaatgtaccggcaatatgacgcagctgcagcaggcgggcaaaatccgtaagcgctcagaaaagca
gttccgatcggtactgaaccgcgcggtactgacgcgttgctcaccctgcagacactggcgtctgcccggctcgcacatcacct
ggaatataaagcgtctgcaagggtcccggtgtaccggcgccggccagggttacctcatcgataagaaaaccgctccagtacaa
5 aatcaccatcatcgctcatctgaagatccgaatatcgaccaactgttcgattcaagcggcgacggaaaagcggtttacacggta
gtacccagactggggctgccaagctatgatggccgacgccatcagtcgctacaaagagggaaccgggtttattacacctg
gacggcgctactgggtgagtaacgaactgaagccgggcaagatgtcgtctgttgaggtgccgttctccgactgcccggcgga
taaaacgccgataccaaactgccgaatgccggtggcatcgaaagccctcatcgccgatgaagaagtcaggctctcgatgcctt
ttgtgatgcgccgtgtgtgggtgtctccactgtgccgactccttgatggcaatcgccgaggggaataatgaactggcgtctttatt
10 cccggcaatccagtttgagtagctgatgtgagcagacaagcagagtggtatggagtaccatgccgcaaaaccgggtcac
accaattctccgaatggaggaggaaaagaaggcgctaccgaggaggagaagaaggcccagctggccctcatcgaggag
aagctcaagcagaacgcacgaacgcgaggagcgagaaaaatgaagccctgcagcgggaaaagaatcgcatcaagtc
ggcaaggacatgaccgagccaagcgcgcatggaggagttgagatgaagaagatcgttgagcagcgcaagcgcgaaaa
ggacgaggagaagggcccgcatcggttaagggtcaattgaggcggaagcagcagcgaaggttagagaacaaa
15 aggaattgggaacgcagagccagctccatcgtagctccaccacagttcgtaccaccggccggtgtgaaatctccggcg
gagactacaccgaaaccgcatccaggcgccagcgcaatcttgccgcagcggctccctactatcaaccgcccgtgttccc
caggatgttcagccgatcgtctatcggtatggagcattcgagttgtctgcggttcccatcagcggctggcattgttctgcg
gggcattatgaagatggtaatgaaatttcgagtgcccaagacatttgcacttctgaccgcattggctgcgaatggagatggcg
gcagcaactgttcttccgcaacctgcattagcccgaacggccgttgcgggcattataaacgcgtacgtcgcattaaaacaaa
20 cataacaactacgtga

- Drosophila Gene Hit** rescue sequence and BLASTX with EST: BIP1 (Y14998),
BLASTX with genomic sequence matches BIP.
- Human Homologue** BLASTX with BIP1: alanine:glyoxylate aminotransferase
25 (X53414) ?
- Drosophila EST** GM04749 (AA695904), GM13608 (AA803601)
- Annotated Drosophila genome genomic segment** AE003556
- 30 **Annotated Drosophila genome Complete gene candidate** CG7574 - bip1 unknown
function
CG13681 – unknown
- Human homologue of Complete gene candidate** none
- 35 **Putative function** no homologies to indicate functions, Drosophila Bip1 interacts with
transcriptional activator Bric-a-brac which is required for ovariole
formation
- 40 **Confirmation by RNAi** Both show reduction in G1 and G2/M indicating fewer
cycling cells

Example 52 (Category 4)

	Line ID	763/4
	Category	Meiotic defects in testis: segregation defects (overcondensation, fewer anaphases)
5	Reversion	R
	Map Position	90F
	Rescue ID	2F5E-1
10	Rescue Sequence	CGGCAATGTCTGCGCCCCCAATCTGAACTTGCCCTCGCCCTCTCCGCCCCCTGATC TCATCTCCTCTTCAAACCCCTGCTCCCCCTTTCTGCACACATTAACGTCAGCCT TTAAGTGTGCTTTCTCAGGTGCTGCCCCCTGCGCCCACCATCCCCCGCTCCATG CTCTTTCCATCTTGCGCTCTCTGCGTTCTATCTACATTTTTTTTCGAGGTCGCGCG 15 CTGCTTTTTCCGTTGATGTTTCGTTCTCGTCAATGTCGCAATATGCGCAAAAGGC AGACAAAAAATGAGTGGAAAAAGTACATACATACCGGTGATTGATGGG CGGTGGGTGGCGGTGGTGTAGGNGTGGTTTG
	Genomic hit, Accession No.	AC006495
20	Associated ORF	Genscan ORF1 predicted sequences >22:47:02 GENSCAN_predicted_peptide_3 283_aa MTERENNVYKAKLAEQAERYDEMVEAMKKVASMDVELTVEERNLLSVAYKNVI GARRASWRIITSIEQKEENKGAEKLEMIKTYRGQVEKELRDICSDILNVLEKHLIP 25 CATSGESKVFYFKMKGDYHRYLAEFATGSDRKDAAENSLIAYKAASDIAMNDLP PTHPIRLGLALNFSVFYIEILNSPDRACRLAKAAFDDAIAELDTLSEESYKDSLIM QLLRDNLTLWTSDMQAEEIPIKLPDRQSKTTLIFSPRSQVNPILHKNNITIIRVIC SVFA
30	>22:47:02 GENSCAN_predicted_CDS_3 852_bp	atgactgagcgcgagaacaatgtgtacaaggcaaaagctggccgaacaggccgagcgtacgacgaaatggaggccatga agaaggctgcctccatggacgtagagctgaccgtcgaggagcgaatctgctgtcgtggcgtaagaatgtgattggagcac gccgtgcctcgtggcgcacatcacctgatcgaacagaaggaggagaacaagggggccgaggagaaattggagatgatcaa aacctaccgcgacaggtggagaaggagctgcgcgacatctgctcgatatactgaacgtgctcgagaagcatctcattccatg 35 cgccacatccggcgaaagcaaagtatttactataagatgaaggcgactaccatcgctacctggccgaattgccaccggctcc gaccgcaaggatcgggcagagaactcgtgattgcctacaaggcggccagcgatattgccatgaacgatctgccaccaacaca ccccatccgtttggccttgccattgaacttcggtgttctactatgagattctcaactcgccggaccgcttgccgcttggcgaaa gccgtttcgtgatgatccattgcgagttggatactgagcgaagagagctacaaaactcgacactcatcatgcagctgtgc gcgacaacctcacattatggacgtccgatatgcaggcagaagagattccgattccaaaactccccgacagacagtccaaaacca 40 cattgatatttagccccgaagtcaagtaaacccaaagattctccacaagaacaacaccatcatcggcagagttatctgtagcgtgtt tgcgtga
	Drosophila Gene Hit	rescue sequence: 14-3-3 epsilon isoform gene (U84898) TBLASTN with ORF1: 14-3-3 .
45	Human Homologue	TBLASTN with ORF1 and BLASTX with 14-3-3: epsilon isoform 14-3-3 protein (U43430.1)

	Annotated <i>Drosophila</i> genome genomic segment	AE003721
	Annotated <i>Drosophila</i> genome Complete gene candidate	CG8045 complex gene
5		appears to encode 3 things : Transcript: CT24102 unknown Transcript CT24072: transcription factor RNA polymerase II transcription factor ,
10		Transcript: CT24092: diacylglycerol- activated/phospholipid dependent protein kinase C inhibitor /14-3-3 protein epsilon (suppressor of ras)
15		
	Human homologue of Complete gene candidate	CT24092: e-119 NP_006752.1 tyrosine 3- monooxygenase/tryptophan 5- monooxygenase activation protein, epsilon polypeptide; 14-3-3 epsilon [Homo sapiens
20		
25	Putative function	transcription factor, or 14-3-3 proteins which associate with cdc25 phosphatases
	Confirmation by RNAi	CT24102: wild type profile only, CT24072: Loss of G1 peak CT24092: Increase of G1 peak

Example 53 (Category 4)

	Line ID	951/8
	Category	Mitotic defects in brain: (some overcondensation, anaphase bridge, metaphase with swollen chromosome and bipolar spindle)
5	Reversion	NR
	Map Position	73D
10	Rescue ID	2E8S
	Rescue Sequence	GTATAACAAGATCCCGAGACACCGGTCAGTTGGTGCTACACGCTCTTGGAGA GCGCTGTGTTTGTTCGGTTCAGCGATTAGCGATAGTTTGTTCGAGCCGGTTGT GTTAACTTGGCTAGCTTCGGGTTTATTGTGACACTTTCCCCAAATCGATCGTTT 15 GCGAAGCGTGCATAGCGGAACATACATACATAGATAACCAGCGTGTCTGGGT GTTTCATGAAAAAGAGTGCGTGATATGGGATTTCGATATGGCAACACGCTTTATG GATATACTAAAGCTGACCTTTAAGTGAGTTTTCCCAGTCAGTGTCCGCTTCTTG CTCTTGCGGAGCGTTAAACGGTTTTCTGTGTTTTGAGGTCTCGCGTCTTGTTTT TGCAACAGCTTCTGCCCAGCATGCACACATACGTGTGCACTGGGAAAATAGTG 20 TTGCAGAAAGTGCTTGATTTATAAATATTACAAAAAATGTGATGAAACACTTTT TATTTTCTTCAAAAAATCAAGAATAAATTAACACTATCCTGCTCTTAAACAT GGAGATTAATTCAATTTTAATTAATAAATAATTTTTTTTACAATTTATGATTTA TGAATTTATGCACTCCTTGAAACTATTAAAGACTCAACAGTGA
25	Genomic hit, Accession No.	CSC:AC015272
	Associated ORF	
	Genscan ORF1 predicted sequences	
30	>23:03:05 GENSCAN_predicted_peptide_1 602_aaMGFDMATRFMDILKLTFKPFKTN YTEEKYFNDKLRSSKNIERRYILDVGRGPTAVTYNPIWVISFKYEQRKLSTAIYSV IKTKSGPVRGVKRNTIWGGSYFSFEKIPFAKPPVGDRLRKAPEAVEPWDQELDCTS PADKPLQTHMFFRKYAGSEDCLYLN VYVKDLQPDKLRPVMVWYGGGYQVGEA SRGLDVVIVTVAYRLGALGFLSLDDPQLNVPGNAGLKDQIMALRWVQQNIEAFG GDSNNITLFGESAGGASTHFLALSPQTEGLIHKAIVMSGSVLCPWTQPPRNNWAY 35 RLAQKLGYTGDNDKKAIFEFLRSMMSGGEIVKATATVLSNDEKHHRLFAFGPVVE PYTTEHTVVAKQPHELMQNSWSHRIPMMFGGTSFEGLLFYPEVSRRPATLDEVGN CKNLLPSDLGLNLDPKLRENYGLQLKKA YFGDEPCNQANMMKFLELCYREFW HPIYRAALNRVRQSSAPTYLYRFDHDSKLCNAIRIVLCGHQMRGVCHGDDL CYIF HSMLSHQSAPDSPEHKVITGMVDVWTSFAAHGDPNCESIKSLKFAPIENVTNFKC 40 LNIGDQFEVMALPELQKIEPVWNSFYAPNKL	
	>23:03:05 GENSCAN_predicted_CDS_1 1809_bp	
45	atgggattc gatatggcaacacgctttatggatatactaaagctgacctttaagccatttaaacgaactacactgaagaaaagtattt caatgacaaactcagatcttcgaaaaatattgaaaggcgttatatcttgatgttgcttcgcggaacccacagcagtcacgtacaat ccaatctgggtaataagcttcaagtagcagcagcgaattgtcaacagcaatatatccgtcataaagacgaaatcaggctcctgtg cggggagtgagagaaacacaatctggggaggaagctacttcagtttcgagaagatacccttcgcaaagcctcgggtgggagat	

165

ctgcgcttcaaggccccggaagcagtgaggccatgggatcaggaattggattgcacttcgccggcagacaagccccttcagaca
 cacatgtttttcagaaaatacgcgggctcagaggactgcctctacttaaatgtgtatgtcaaagatctgcagccggataaactgcgtc
 ccgtgatgggttgatctacggaggaggctatcaagttggcgaagcttctcaggagtgatgtggtcatagtcaccgttgcttatcg
 actgggtgccttgggcttctcagcctgatgatcccaactaaacgttccggaaatgcagggtctcaaggatcaaatcatggccc
 5 tgcgatgggtgcaacaaaacatcgaagcattcggcgggtgattccaacaatattacactcttggcgaaagtgcggcgaggcctc
 gaccacttcttgactaagtccccaaactgaaggtcttatccacaaagctatcgttatgtcgggcagtggtttgtgcccctggacg
 caaccaccgagaaataattgggcttataggctggccaaaaatgggatacaccggtgacaataaggacaaggcgatcttgagt
 ttctgcgatcaatgagtgggcggggagattgtcaaggccaccgcaacagttctcagcaacgatgaaaagcatcatcggatcctttc
 gccttcggacctgtcgtagaaccatatactaccgagcacactgtggtcgtlaaacaaccgcatgaactgatgcagaatagctgga
 10 gtcacaggatacccatgatgtttggaggcacgagcttcgagggattgctattctatccagaggttcaaggcggccagcaaccctc
 gatgaggtgggttaactgcaagaatctgtaccgagcgatctcgggttaacctagatcccaactgcgtgagaactacggcttgca
 actgaagaaggcgatattcggcgacgaaccctgtaaccaggcaaacatgatgaagtttctcagctatgctcatatcgagagttctg
 gcaccctatatacagggcagcttgaaccgtgtccggcaatccagcgcaccacgtatctgtatcgattcgatcacgattccaaact
 gtgcaacgccattaggattgtactttggccatcagatgcgaggtgttgcacgtgacgatctgtctatatattccacagcatgtt
 15 gtcgcatcaatccgctccgattctccggaacacaaggtataaccggaatggtcgacgttggacgagtttcgcagcccacgga
 gatcccaactgcgaaagtataaaatcactcaagtttgacccatcgaaaacgtaaccaactttaagtgtctcaatattggggatcagt
 ttgaagtcatggcgcttcagaattgcagaaaatcgaaacctgtgtggaatagttctacgccccaaacaaactgtag

***Drosophila* Gene Hit** TBLASTN with ORF1: alpha esterase (aE10) gene (U51054)
 20 **Human Homologue** TBLASTN with ORF1 and BLASTX with U51054: bile salt-
 dependent lipase (S79774)

Annotated *Drosophila* genome genomic segment AE003671
Annotated *Drosophila* genome Complete gene candidate CG1131 - alpha esterase 10

25 **Human homologue of Complete gene candidate** 4e-48 4557239
 ref|NP_000656.1|pACHE|
 acetylcholinesterase (YT
 blood group) precursor
 30 >gi|113037|s

35 **Putative function** alpha esterase
Confirmation by RNAi Only wild type profiles observed

CATEGORY 5: SMALL IMAGINAL DISCS (BLOCK TO PROLIFERATION)**Example 54 (Category 5)**

	Line ID	113/20
5	Category	2nd chromosome, small imaginal discs
	Reversion	R
	Map Position	50D/E
	Rescue ID	EcoR1
10	Rescue Sequence 1	
	CTGAGGCNCTTTGCCAATATGTGTATATTGGGCGGGGNACATGCGTNAATCGG	
	TTAAAGCCGCTACTTACATTCTGTTCTTTGCATCTCCCCATCCACAGCTATAA	
	AGCAAGATGAGCTACGCCGCTGATGTGCTGAACTCGGCCCATTTGGGAGCTCC	
	ATGGTGGTGGCGACGCCGAGTTGCGTCGTCCATTTCGATCCCACGNNCCATGAT	
15	TTGGATGCATCCTTCCGCCTTACACGCTTCGCANATCTAAAGGGGCGCGGCTG	
	CAAAGTGCCCGCAAGGAAGTNGCTCCCCCACCT	
	Rescue ID	BamH1
	Rescue Sequence 2	
20	CCACCTGGTACCACAGCGCTCANACGTGTATGTACACGGATTTTCTGCCGCGT	
	GTGTGTAGCGCGGCCCGTGATTGGCTGCAGTCGCGATGGCGGCTAAAACGGG	
	CGAAGTCAGTATTTCTCCCTGTGCGACGANGCGAGCAACGTGAACAATGCCAC	
	TCATTTCAATTGCAAAATGCCAAAAAGTGCGCGCTTTGAATTGGCCATTTGGT	
	TCGTTGCGTTCGTTTGTCTTTTGGTACTTACGTTTGCTTGTGCGATTGTACAAA	
25	GATAATTGTAGAGTAACGTTAGCAAATTATATTTATTTTGGCCTGGTTTTTGC	
	TTTTCCAACGANCGAGATGTCACAACAGGGTTGTATTANCGTGTGCGGCTGAT	
	TCGATATTTGGGATGCCGATTGTCTGAAGCGANGGTTCAACGGGGCTGCCAAC	
	TCCCCCGAAAATCTATCNATGGTATTGTGCGCCAAGGGTAAAATAAATAAAAA	
	TATGTTAAAACCGCGGAATAAATGGGGGAACCGAAGTGGAAACTGTGGTTCA	
30	CAGTGCTCTGACTTTCGGGAGCAGTTAATATAGTTGGCATTAATTCAATTAGA	
	GCTCCAAAGTGCTGGTCACAAAGAACGCACAAGAACGGGCCATGAAAAACCT	
	GTTGCGCCAGCAGAACGAAAAGTAAAAATTAGAAGAAACCAT	
	Genomic hit, Accession No.	CSC:AC017131
35	Drosophila Gene Hit	rescue sequence: selenophosphate synthetase (ptuf1) (U91994)
	Human Homologue	BLASTX with U91994: SELENIDE, WATER DIKINASE 1 (SELENOPHOSPHATE SYNTHETASE 1) (SELENIUM DONOR PROTEIN 1) (P49903)
	Drosophila EST	LD46437 (AI514756 similar by BLASTN to U91994
40		selenophosphate synthetase (ptuf1) gene)

	Annotated <i>Drosophila</i> genome Complete gene candidate	CG8553 selD selenophosphate synthetase
	Human homologue of Complete gene candidate	1711372 P49903 SELD_HUMAN SELENIDE, WATER DIKINASE (SELENOPHOSPHATE SYNTHETASE (1e-159)
5		
10	Putative function	selenophosphate synthetase
	Confirmation by RNAi	Only wild type profiles were observed

Example 55 (Category 5)

	Line ID	121/1
5	Category	2nd chromosome, small imaginal discs
	Reversion	NR
	Map Position	60B
	Rescue ID	BamH1
10	Rescue Sequence	TCCTGTGCACTCATATTGATTTGCCTTGTC AAGTGGCTAAAGAAATATTTAAATG TTTGTTATTTCTGTTGCTAGCGCTCCGACAGTCTGGCAGCACTGCTCGCTGTCG ATAGTTCAACTGAGTTGCTGTTTCATCGAACAGAGCTGCCAACTCTATTTTTGT AGCTGGCCAGCCAGGATTGCCAGAGTAAGGCCCTCAAATCAGCTGTTTTGTGT 15 TTTGATTTTATTTTGAAAGTCCTAGTTTAAAATTATGCTTTCTCCGACAGATCA GCACAAATAATACTAATAAAGCTCACAATGCTAAGGTTGTGCCTTCCAACTCG AGCTGGATATGTGCGTAAGTAAGGACTTTACGTCTATAAACTTGTTATGTAA AGTAAATGTTTGCCTATTGCGCAATTTCTCCAACGAAAAACCCAGAAAACCNA AACCCCTTNAAANTTTGGAATATNCCCAATGAATGCAGCACCCGTGAAATCC 20 GTAATGCCTTTGTCCAGCTCTCCAAATTGGTAAGTAACTCCAAGATCCAAAGG AGCCTCCTAAACCCTGCCCTTTCCACAGTACCACCCAGATGTTAAGAGCAATG CTGCGTGTCCGGAGCGCACAGCCGATTTGTTTCAGATCTCCGAGGCGTACAAG AACCTGATAAAGCCGGAACGGAAGGAAAAA
25	Genomic hit, Accession No.	CSC:AC020499
	Drosophila Gene Hit	rescue sequence: DnaJ60 gene for dnaJ-like protein (Y11900)
	Annotated Drosophila genome genomic segment	AE003463
30	Annotated Drosophila genome Complete gene candidate	CG12240 – DnaJ60 CG13570 – spaghetti ser/thr phosphatase
35	Human homologue of Complete gene candidate	CG12240- 4827026 ref NP_005138.1 pTID1 tumorous imaginal discs (Drosophila) homolog >gi 3372677 (AF061749) 7e- 08
40		CG1116- 2495728 HYPOTHETICAL PROTEIN KIAA0258(aa)
45	Putative function	CG12240 : Chaperone involved in protein folding CG13570 : serine/threonine phosphatase

Confirmation by RNAi

CG12240: Marked reduction in G1 and G2/M peaks
indicating fewer cycling cells
CG13570: Marked increase in G1 peak

Example 56 (Category 5)

Line ID 127/2
Category 2nd chromosome, small imaginal discs
5 **Reversion** NR
Map Position 57F

Rescue ID EcoR1

Rescue Sequence 1

10 GCCGGTGGGCCCACACTTGTNCGCCCGCGCATCGGCTGTCTGTGGGAGTGCGA
NCGAGTCAGATAGTAGATCCGATGCGCTCTCCAGATACTTTTTGAACACTGAA
GAAAACGCGCAGTTGTGGGTGAATTCAGCATCATCAGATTGAATCACACACA
ATCCTAGTCGCCTCACGCGAAGAGAACTATGTCATGATCAGATATCGGTGTAT
GCATTCTATATTATGTACTTCGAAATATGTAATTTATTAAGTTTTCGCTATACT
15 TTTCAATCAAATTGGCAAAAACCAATTCAAAGGTTTTCAATATTTTCGAAAAG
CATTTTAGGCTTTCTATGTAACGTATGTTTTCAAACAAAATATTAGTTTTTGA
AACTTTATTATCGGATAAAACAAATGTAAGCCAAATNACAACGTTNTATGATAC
TCCCAAAGATCCGCNCTNTTAAAGTGGCCTAAAAATAGCTGACGCATTAANCC
ATAGGCGCTTCGCTTCTCAAGATAAAACCTGGCGTGCTCAACTCAAGAAACAA
20 ATATGTGGTTATATACATATATACATATATGTTGGGCATATAACCGATGTGTGAC
GTGACATTGGCTCGTTCTATTACATACTTAAACACTAAATGCAAACCTATCA
AAAACCNACTACACTAAGCGAAAAACGGCAGANATAGTTAAGGAAAGTGGTC
CA

25 **Rescue ID** BamH1

Rescue Sequence 2

CTTCTTTTCTCAAAAAACGTCGCTCGNGTCCCNCAATCGTTTTACAAACTTCGC
TCGGAACGGACGTGTGCGCGCTCTGAAAGGAAAAAGTGAAAAAGTGTGTGAC
AAAGTGCAAATAAGCCACAACGCGCATGTGAGAAATCAAATTTAATTGAGAA
30 GCATCAAAAATTGTATACATATCGAGCGTATCCACATCGCTGTATGTGTGAGT
GTGCCAGTGCTAGTGTGGTTTTCCCTTTTCGCCGTGGAAAATATGAAAACCTGA
ATGAAAAACTGAATCGCAGTCAGCCAGAGCCGAATTGGAAAAGAGTAACTCG
CATTGGGGACACGAAGAGGTGTCTCGAAAAAGGTAAAATCTTTTACACAGAA
ACGACGCCAGAAAGCGATTAGCGATTINTGACTATGTGTGAGTGTGTAATTC
35 GGTCTACGGCTGTGTGTCTGCATTTTATTTAACNTTTTGTTCCTCCNGTTNGNTC
CACNGTAAAAATAGCTAAAAAAAAGGGCAAGTACTCTTGGCGCGCTCTCCC
TCTCTCTTTGTTGGTCTGACTGCGACGTCACCGTTCACGTAGAATCGTTTTCA
AGTGGCGTTTCTTTCTTTCTTTTAAATGTGCTGCTTCTTGCTTCTGCCTCTTCTTC
TTGCCTTTGGCTATCTGCTTTGTTTTGAAATACGTCCATGTTATCCAGTGTCTG
40 TGCCAAATGTGTGCGANATGATCTCTACTT

Genomic hit, Accession No. AC009732

Associated ORF

45 Genscan ORF1 predicted sequence
>/tmp/aaaaafrla|GENSCAN_predicted_peptide_2|456_aa

171

- MQTKGPITDADCIRGMACRALAGLARS DRVVRQIVSKLPLFASGQLQTLMRDPILQ
 EKRAEHVIFQKYALELLERVSGKTKPLNNPLDPSLSNMHKANVIAQTRIQYNKQQ
 LYQLIFEHLESNGLSQT AQMLQREVGLPLQTPTRSFHQSPFDYKSLPSGSSSLSRN
 RLRSRMQDVNAAIMGNGLNRSFGEDSSPAGAGGSNAGDGVSPNPFSSSLNTTQTP
 5 IKIRRTDRSSVSRSIQKQAMEPGGMSVGLAEDGQLHPKRITLNTIVTEYLTNQHSL
 CNNPVTTCPQFDLYEPHKCPDPKPSRLLSSNYNLTSRHARTQAGFNTSRFDRRYV
 HTHFSPWRSIRSADYEDLEFTCCDLAGKYIIVGTQQGDGRVFNMNDGVEQFFSNC
 HNFSVDAIKANRAGDLVITSSFWRTPTSILWSIADDEFKLLRLPDVITYCEFSQTV
 QDRLLGTQNEVY
 10 >/tmp/aaaaafrla|GENSCAN_predicted_CDS_2|1371_bp
 atgcagaccaaaggaccattacggatgcggactgtatcgtggaatggcctgtagggccttggcgggacttgctcgtccgatc
 gggtcaggcagatcgtcagcaagcttccactcttggcagcggacaactccagacgctgatgcgggatcccatactccaggaga
 agcgcgcggaacatgtaatctccaaaagtacgcattggagttgctagaacgagtgctcgggtaagacgaaaccgctaaataatcc
 15 ttggatccatcgtgtccaacatgcacaaggccaatgtaatgccagacacgcacatccagtataacaagcagcagctgtatcagc
 ttatcttcgagcacctggaagcaacgggtctctcccagacagcccaaatgctgcaacgggaggtgggtcttccgctacagactcc
 cactacgcgcagtttcatcaatcaccttccgactacaaaagtctccagtggttagctcgtgtctagaatcgtctcgaagc
 cgcatgcaagatgtgaacgcagcagataatgggcaatggagacttaaacagaagtttggtaggactcctcgcggcaggagcc
 ggtgtagcaatgcgggagatggagtcagcataccaaatttagctcccttaacacaacgcagacgcccataaaaaataaggagg
 20 acggatagaagttcagttagccgctctatccagaagcaggcaatggagcctggtggcatgctcagttggtcttgcgaagatgtca
 actgcatcccaagaggatcaccciaataaccatcgtaacggaatacctcaccaaccagcactcgtgtgcaataatccggtgaca
 acctgcccgcagtttgattgtacgagccgcacaagtgtccagatccgaagcccagccgattgctaagctcgaactacaacctga
 ctatcgcatcgtcgaacccaagccggatttaataaccagtcgttggacgtcgtatgtgcacacgcactttaccatggcgta
 gcattcgatcggcgactacgaggacataggttcacctgttgcgatttggcggtaatacatcattgtgggcacgcagcaggg
 25 cgacggacgagtggtcaacatgaacgatggcgtggagcagttcttccaactgtcacaactttagcgttgatgtatgaaggcta
 agagccggagacttggtcatcacatctagcttctgcgccacacccaccagcattctatggtctattgcggacgatgagttcaagcta
 aagttgcgacttccgatgtcacgtactgtgagttcagtcacaacggtgcaggatcgttgttgggcacccagaatgaggtatactaa
 corresponds to CG10082
 30 *Drosophila* EST several including SD04293 (AI532704)
 Annotated *Drosophila* genome genomic segment AE003454
 Annotated *Drosophila* genome Complete gene candidate CG10082 – novel protein with
 35 homology to enhancer Pi uptake
 Human homologue of Complete gene candidate 1665793 dbj|BAA13393|
 (D87452) Similar to
 40 *S.cerevisiae* YD9335.03c
 protein (S54640) [Homo
 sapiens] (2e-43)
 45 **Putative function** Putative phosphatase or enhancer of Pi uptake protein
Confirmation by RNAi Reduced G1 and G2/M peaks indicating fewer cycling cells

Example 57 (Category 5)

Line ID 131/8
Category 2nd chromosome, small imaginal discs
Reversion R
Map Position 60A

Rescue ID BamH1

Rescue Sequence 1

10 CACGATTGCNNGGCCCATCGAAGTGTGGGTCTATCGATACTCGTGGGTAAATAA
 ACAAGTTCTGAACTGCGATTTTCGGGGGTTTGAGGGGTCAATTGTCCCCTGTGT
 TGGAAATGTGTTCCCTAAATCTACACAAACACTCCCTAAGCTTATCCTAAACTTAT
 AAATATTGGTTGCTATTTAAACCCCATTTACGGTTATCCAGCACGCCCCCTGA
 ACTGTGACCCACATCCCCGATTTTAGTGACTAGTTTATACTTATCGTGGTTGG
 15 CATTTGGTACACTACACTTTCTTATTCACCTAGATCGCCGACTCCGCGCACGGT
 CGCGCTCCCGTTCCCGCTCCCGATCTCGGCTGCGACTGCGGTGCGGATCCCGTT
 CCCGGTCGCGGCGACCGGCGCCTCCANATCCGGATCCCTAANCGGCANCNCT
 CNTGGTGGCAATCNNGGAATGTTCCGGGGNCCNCTACCNAGTGNAATCAC
 TGGTACGTCCCACCGCNAAACTCCGCCANTGCGGTTGCCGGAACGGGTGGC
 20 ANTGCCAATGGGTGCTGCAGAAGGTACCATCACAGCAATCGCTCACGGANC
 CCGAAGACTGCCTCTGCCGCCCGGTGGGCCACTCATACACGCTACACGGTCG
 GAAATACTACATTGATCACAATGCGCATACCACGCACTGGAATCATCCGTTGG
 GAACGC

25 **Rescue ID** EcoR1

Rescue Sequence 2

AATTGATTTCCGGACATATAAACAGAATCCAGAACTCATCCGGCAGCAGGCTC
 AGTCAGGCCAGTAAATCCGAAAAGAGAGTAACCAGCAGGAAAAGAGAATCC
 ACGTAAATACAGAGAAAATGGCTCTACGCGTCCAATTCGAGAACAAACGACGA
 30 CATCGGCGTATTCACTAAACTAACCAACACATACTGCCTGGTGGCCATCGGTG
 GATCCGAGACCTTCTACAGCGCCTTCGAGGCGGAGCTGGGCGACACCATCCCG
 GTGGTGATGCGAATGTGGGCGGCTGCCGGATCATCGGCCGCCTCACCGTGGG
 CAACCGCAACGGCCTGCTGGTGCCCAACTCCACCACCGACGAAGAGCTGCAA
 CACCTGCGTTACANCCTGCCANAACCCCGGAAANATTTATCGTGTGGAAGAAC
 35 GCCTGTCCGCGCTGGGCAACGTTATCGCCTGCAATGATTATGTGGCCCTGGTG
 CACCCGGATCTGGACAAGGAGACCGAGGAGATCATCGCGGACGTGCTCAAAG
 TANANGTCTTCCGCCAGACCATTGCCGACAACACTACTGGTGGGCTCTTACGCC
 GTGCTGAGCAACCAGGGGGGCATGGTGATCCCAAGACNAGCATTACAGGAAC
 AGGACAACCTGTCGTCCCTGCTGCAGGTTCC

40

Genomic hit, Accession No. CSC:AC020517

Associated ORF

Genscan ORF1 predicted sequences >22:13:05|GENSCAN_predicted_peptide_4|357_aa
 45 MALRVQFENDDIGVFTKLNTYCLVAIGGSETFYSAFEAELGDTIPVVHANVGG
 CRIIGRLTVGNRNGLLVPNSTTDEELQHLRNSLPDAVKIYRVEERLSALGNVIACN

173

DYVALVHPDLDKETEEIADVLKVEVFRQTIADNSLVGSYAVLSNQGGMVHPKTS
IQDQDELSSLLQVPLVAGTVNRGSEVLAAGMVVNDWLSFVGMNTTATEISVIESV
FKLNQAQPATVTTKLRAALIEDISRSRVAGGGGGGGGGSSGNSSSGPSTSRRTT
5 RNNAAATAADRPKINEADLEGKSPREEVMLKTMGFCTFDTTKNRKVEGNDVGEV
HVILKRKYRQYMNRRKGGFNRPLDFVA

>22:13:05|GENSCAN_predicted_CDS_4|1074_bp

atggctctacggtccaattcgagaacaacgacgacatcggcgctctcactaaactaaccaacacatactgcctggtggccatcgg
tggatccgagaccttctacagcgcttcgaggcggagctggcgacacatcccgggtgcatgcgaatgtggcggtgccc
10 ggatcatcgccgctcaccgtgggcaaccgcaacggcctgctgggtcccaactccaccaccgacgaggagctgcaacacct
gcgtaacagcctgccagacgcgtgaagattatcgtgtggaggagcgctgtccgcgctgggcaacgttatcgctgcaatgat
tatgtggccctggtgcacccggatctggacaaggagacggagagatcatcgcgacgtgctcaagtagaggttccgccag
accattgccacaactcactggtgggctcttacgcctgctgagcaaccaggcgccatggtgcattccaagacgagcattcag
gaccaggacgaactgtcgtccctgctgcaggttccctcgtggcggaacagtgaaccggggcagcgaagtactcggcgccg
15 gcatggtgctcaacgactggctctcctcgtgggcatgaacaccacggccacagagatcctcgtgatcgagagcgtcttcaagctt
aaccaggcacagcccggccacagtgcagcaacgtgctgcccctcatcgaggacatcgcggtcgagggtcgccgga
ggaggaggaggaggagcgccggcggaagcagcgccggcaacagcagctccggaccatcgacgtcggaaggacgacg
aggaacaatgcggcgccacagctgcggaccggcccaagatcaacgaggcgacgtgagggttaaatcgccgggaagaggt
cgagatgctgaagacaatgggattctgcacgttcgacaccaccaagaacaggaaggtcgaggggcaacgatgtcgagagaagtg
20 atgtaatcctcaagcgaagtaccgccagtacatgaatcgcaagggtggcttcaaccggcgctcgattctgtggcatag

Drosophila Gene Hit rescue sequence and TBLASTN with ORF1: b(2)gcn
(EUKARYOTIC TRANSLATION INITIATION FACTOR 6
)((X97641)

25 **Human Homologue** BLASTX with X97641: integrin beta 4 binding protein (HUMAN
EUKARYOTIC TRANSLATION INITIATION FACTOR 6)
(NP_002203.1)

Drosophila EST GH08760 (AI109537 similar by BLASTN to X97641
"D.melanogaster b(2)gcn gene.")

30

Annotated Drosophila genome genomic segment AE003462

Annotated Drosophila genome Complete gene candidate CG17611 – bcgn benign
gonadal neoplasia homology
to Eif6 translation factor

35

Human homologue of Complete gene candidate 6016331 EUKARYOTIC
TRANSLATION
INITIATION FACTOR 6

40

(EIF-6)(aa) and 4504771
|ref|NP_002203.1|pITGB4BP|
integrin beta 4 binding
protein(aa)

45 **Putative function** eukaryotic translation initiation factor 6 (eif-6)(aa)

Confirmation by RNAi Slightly reduced G1 and increased G2/M indicating block in
G2/M

Example 58 (Category 5)

	Line ID	135/25
	Category	2nd chromosome, small imaginal discs
5	Reversion	NR
	Map Position	24A
	Rescue ID	EcoR1
	Rescue Sequence	
10	ATAACATGGGCNCTGGTTTTTAAGTNAAGCTCTANTNATTGGCCCCCATTCTTA	
	NNCTCTCTCGCTCTCTTCTCGCTCTTTCGCCTGCTCTCTCGCCTGATTATTCTGC	
	TTGGTCGGCTGATGGTTTTTNGTTTINATCTGGTGTATTTCTGCGTAGTTTATG	
	ACAAACCGGCTGGTTCCTGTTGTTATTGCCGTATTCTAATATATTTCCCCTATTG	
	TTCTTATTTTTGTTGCAGCCTGCACACCTCGGAGGTTCTAGATGATAAGGGGTG	
15	TAGCGATGGTGGGGGGCTGTCTTGANGGGCTTCTCGCCTTGAGCTCTTGTATTAT	
	CTTTGGTCATTTGTTATTGTTTAATGCACGGCAATATTATTGGTAAACAAGTTA	
	GCCAACAGCACTAAACGCCAATCGCATTCTTTCTAAAAACCAAGTCTATTGT	
	CGATCTTGCTAGGGAATGATGATGACTCAGGTGCAATTGGGATCTTATCTAT	
	GGCTGTCTGGGAATCAAGAAGTGTTCGCCGAGAATTCGTGAANTACTGCCGCT	
20	CTCTCCATGGGGCCATTATTTGCACTCGTTTTNCGCGAAATACCATNAATTAGC	
	ATAAAGACACGTCGCCGGCAATCGTGACGTAGGCTATNAATGCCTTCTATGCA	
	TGTGCNAACTCGCGGAAGCATAGCAATTTGAAGGAACAATTTTCANTGCAG	
	GTTTTAATGGGCTAAAAAA	
25	Genomic hit, Accession No. CSC:AC014199	
	Associated ORF	
	Genscan ORF1 predicted sequences >20:54:54 GENSCAN_predicted_peptide_3 117_aa	
	MSASPTARQAITQVMPMITRKVVISDPIQMPEVYSSTPGGTLYSTTPGGTKLIYER	
30	AFMKNLRGSPLSQTPPSNVPSCLLRGTPRTPFRKCVVPTELIKQTKSLKIEDQEQQF	
	QLDL	
	>20:54:54 GENSCAN_predicted_CDS_3 354_bp	
	atgtccgcttcacccaccgcccgtcaagccatcaccaggttatgccatgatcaccaggaaggtgtcatctcggatccgatcca	
35	gatgcccaggtgtactcctcgacgcccggcgaacccttactccaccactcctggaggcaccaacttatctacgagcgggc	
	tttcatgaagaatctcgtggctccccattgagccaaactccgctccaacgtgccagttgcttctgaggggaactccgcgta	
	ctcccttcgcaagtgcgtgcccgtcccacggaactgatcaagcagaccaagtcgtgaagattgaggaccaggaacagttcc	
	aactggatctgtag	
40	Drosophila Gene Hit TBLASTN with ORF1: BcDNA.HL08053 mRNA (AF132557)	
	Human Homologue TBLASTN with ORF1 and BLASTX with AF132557: eukaryotic	
	translation initiation factor 4E binding protein 2 (EIF4EBP2)	
	(L36056)	
45	Annotated Drosophila genome genomic segment	
	AE003579	

175

	Annotated <i>Drosophila</i> genome Complete gene candidate	CG8846 - phas1 translation initiation factor 4E binding protein 2
5	Human homologue of Complete gene candidate	CG8846 - 4758260 ref NP_004087.1 pEIF4EBP2 eukaryotic translation initiation factor 4E binding protein 2 (4e-16)
10	Putative function	EIF4E translation factor binding protein
15	Confirmation by RNAi	Slight reduction in G1 and G2/M indicating fewer cycling cells

Example 59 (Category 5)

Line ID 141/12
Category 2nd chromosome, small imaginal discs
Reversion R
Map Position 21A/B

Rescue ID BamH1

Rescue Sequence

10 GGCTCTTTTCCAAANAGGCAGTTTCTTGNCCCATTCTTGGATTGCTTTGTAGT
 GAACTNAATCGTTTTGTGTTGTTTCTCTGTCGTCCAGTCTTGTGAAAATTCGTG
 ATAATAATGCCTGGATAAATANTTAAGCATTGGAACCGGGGAAAAAGGG
 CTAAGTTGTGTGAAGGAAACAATTGAAGTGACCCTTTGTNTATAAACATTCCA
 CGACGTGTTTCGAAAAACAAACAAAGATATGCGGAAACAAAGTGTTAATAAAA
 15 GAGCNAAAAATAGAGAGAGAGTGTGCGGATAAGCGGTTGAGCGAGATAGAG
 AAAATTGTTGATTAAAAATGTGTGTCNAAATAAACATCAAGCCGCTTGAACGA
 ACAGTCAGTTAGTTGCTTCTGATAATAACCATGGGAAGCGGCNCGTGTGCTTC
 GCTCCTCGTTACTTATAAAATATTTAAACGTTTGCACTTCTCNTATTTCCGAAT
 TTTTGCNCCCCTGAANCAACTTNGTTAAACTGCAAATAGCAATGCAAACAAAC
 20 GAATAGAAACTGAAATCGACAACNACATGTGAAATTCACAAATCAAATCGCA
 ATTGTCATCCCAAAGATATAGAACAAGCTATAGGGAAGATANAGAATGTAAG
 TGCCAAACTAAAATAAACAAACAAGAATAACATTTCCACAGGTGTTTTGCATT
 TCAAATGCATATTTCCGTGGCGGNTACAAATCTTTTCAAACCG

25 **Genomic hit, Accession No.** CSC:AC017815

Associated ORF

Genscan ORF1 Predicted sequences >17:48:30|GENSCAN_predicted_peptide_2|554_aa
 MSNKKMFNRRTTSVSPGQLHYHYHTDFYYSMPDLHKTRKMHGVKRVLVFCLMIVIL
 30 PAILIIMPLHLRKTVFADVYPMAESDIEIRAGISSIFCSKHTLRMNSNFNAFQLRNL
 PEIATNRKHIRLKKSMITLPDDTLEYWGFFLLKGAKVRVKFCSRYDGSRLIIHGHR
 ELNLCLGLTDHNKNKLGANYAKGHEQVQVFFEDNVEITEEKGNQDVLMEHENHG
 GEDLTEDIPQPQVNIPVKQNNSIQPKLIRKKLKGTHHGEHDMHAITDLQGSHTT
 EHILNHHDHSSNSPAHHHNSAHHREHSSNITNEETSRNHIRNEDEDPDQNSSKTH
 35 YSAESPPIRERLKRHNRAHRNQKRQDLYDTLYKRSKRENVYDRKTIHGGNAIN
 FTETDESNSVSSFETGLFQCFNGMILLQEFFRPKNECSNPHIMDTSPNKSSMVVHN
 VIEDGYYYYYIFYSDNDHVQNEIHAFDIYKPTYQYSNMSESQSCLNTTNCTFNISFL
 SDEIVVVEVPTRDIEHEEDDITNLISTCHPRSEIYAIFPITVLVLILCCSFL

40 >17:48:30|GENSCAN_predicted_CDS_2|1665_bp

atgtccaacaaaaagatgtcaacaggactacgtcagtaagtcctggacagttgcattattatcacaggatttctattactcaatgcc
 ggatttgcataaaacccgcaaaatgcacggcgtgaaaagggtgctggtttctgctgatgattgtgatactgccggccattctatc
 attatgccgctgcatttgcgaaagacggtgttgcgacgcatcatccatggcggagtcgatacattgagattcgggcagga
 atctcgtcgatcttttctcgaacacacactgcgtatgaactcaattcaacgcttttcaactacgtaataagccggaattgcgac
 45 gaatcgcaagcaccattaggtgaagaagtcgatgacattgccgatgatacgttgaatactggggcttctcttgcgaaagggtgc
 caagggtcgagtgaaattctgctccgctacgatggatcccgatcctgatcatccatggtcacagggagcttaattcttgcggtct

gaccgatcacataagaataagtggcgccaattatgccaaagggtcacgaacaggtgcaggtgttcttcgaagacaatgtggag
 atcacggaagagaagggaaccaggatgtgctaattggagcacgagaaccacggcggagaggatttgactgaggatattccaca
 gccgcaggtgaacatacctgtcaagcaaaacaattctatacagcctaagtaattaggaaaaaactgaaaaagggcacaattcatc
 atggcgaacatgatatgcatgctataacagattgcaaggatcacaccatacggacacatattgaatcacatgacagctcta
 5 attctccagcacatcacacaatagtactgccatcatcgggagcacagttcgaatatcacaacgaagaaactagtctaatcaca
 tacgaaatgaagatgaagatccagatcagaattcaagtaagaccattatagtgcggaaagtccgcctcaccgggaacgtctcaa
 aagacacaatagggtagcccataggaatcagaagagacaggatctttacgatacgtttataaaagatcaaaggaggagaatgtc
 tacgatagaaagacgatccatggaggaaatgctataaattttacggaaacggacgagtcgaattcgggtgtccagctttgagacagg
 actatttcagtgtttcaatggaatgatcctgctgcaggaggttcttcaggccaaaaaatgaatgctcaaatccgcacataatggacactt
 10 cgcccaacaagagttccatgggtgtgcacaacgtcatcgaggatgggtactactattatattctacagcgacaatgatcacgttc
 aaaacgagatccacgccataatcgaatttacaagccgacgtatcagtactcaaacatgagcgagtcacaaagctgtctgaatacc
 acaaattgcacattcaacatcagtttcctttcggtatgagattgtgtgtgtggaggtccaacacgggatggtatcgagcacgagga
 ggacgatataaccaatctgatctccacctgtcatccgcgcagcgagatatagccatctttccattacgggtgctgtgtgctgaccttt
 gctgctccttctgtag

15 corresponds to CG9524

20 **Annotated *Drosophila* genome genomic segment** AE003623
Annotated *Drosophila* genome Complete gene candidate CG9524 - novel His-rich
protein

Human homologue of Complete gene candidate none

25 **Putative function** No homologies which indicate function

Confirmation by RNAi Reduced G1 and G2/M peaks indicating fewer cycling cells

Example 60 (Category 5)

	Line ID	146/2
	Category	2nd chromosome, small imaginal discs
5	Reversion	NR
	Map Position	26B
	Rescue ID	EcoRI
	Rescue Sequence	
10	TTTNATCCAAACTGAGANACTNTTGGCCCCAAAACTGAAAACTCGGACTCGGG CGCGTAAGGGAGTCGGTCNTCGGGAGTCGGTCGTCTTTTGTGATCTTGAGAC TGAAATTCCAATTGTTGATTTATCTCTCGGCTGCTGCGCCGCGGCTGCGCTGCT GCAGCGCAGTCCCACTCCGATTTGACCAGCGACCAAGTTTATAAACTTTGAG CCAAAATGCAGCGGCGCACAGTTGTTACCAAAACGTTGCACGCGTCGTGGCCC	
15	TCATCAAAACAAAAAAAAAAAAATATAAGCGAAAATGAAAACGAAATTCGGTTA ACGTCAACAGAAGCTGACAAAAGGCAGAAAAGACCGAAACAAGTTGCAGGG CCAGAGTAAGCCAAGTTAAATGCGAAAGAGAAGCAAGAGNCAAGAAGAAAN AATGGGCACTACATACATATATTATAGCCAGCTAATCTGTTGTGCAGTGC GTT TTATCAGCCNCGAAAAGAAAACGAAAACGAAAAGTCGGTCCAAGTTTCGGAC	
20	TCAAAATCCAAACAGAAGAGACTCCATNCCATCAGAGACACGCGGATCTCAT CTCGGTAATGTCTCAATAAAAAGTAATCTTAACTGCCGCCGGAATGTTGAAA AAGTGAAAATTGAAGCGCTTAACGTGTTTCGAAATACGATACATGAGAAGTCC CAAAAAAAAAA	
25	Genomic hit, Accession No. CSC:AC019865 <i>Drosophila</i> EST GH19286 (AI388389)	
	Annotated <i>Drosophila</i> genome genomic segment	AE003481
30	Annotated <i>Drosophila</i> genome Complete gene candidate	
		CG11353 - novel with weak homology to sugar acetylase? CG7525 - tie receptor protein tyrosine kinase.
	Human homologue of Complete gene candidate	
35		CG7525- 4e-23 4557869 ref NP_000450.1 pTEK TEK tyrosine kinase, endothelial >gi 464868 sp Q02763 TIE2_
	Putative function	Sugar acetylase and receptor tyrosine kinase
40	Confirmation by RNAi	
		Both gave a reduction in G1 and increase in G2/M peaks indicating arrest in G2/M

Example 61 (Category 5)

Line ID 155/13
Category 2nd chromosome, small imaginal discs
5 **Reversion** R
Map Position 21B

Rescue ID BamH1

Rescue Sequence 1

10 GNTTTAGTCCNCTTTTGANAGGGNCTTGGNGNCTTAAANAANNAAAAAAGGG
GNCCCGGCNCCCAGCAAANAGNNTAAAACTTGAATGGTTTAATTCGAAAATC
TTTATAGAAATGTCGCCTAATACCTTATCGGTATAGAGTTCACCTCGTCTCCTAA
TCCATATTTTAAGATATCAATATCTATTAACAATTTTATCGTATGATTAGAAA
TTCGCATTGTTTTATTATTTTCGACCTTTGGGCTTTACATCGACAGCTACTCTCTA
15 TCCAGACAGGAGACTGGGAGAGAGAGCACGATGCTGTCTGAAAGCATGAATG
ATGGATGCTGTGCCTATGTGCGATATGCACGTTGCCTGAGCTAAAACGAAACG
AGATTATTAATCTATCCGCAAGATTGAGATGCTGATTCCACATGAGTGAGCGA
GTCCGTGAGTGGATATTGCTCTCTCCGAAATGCATGCATGAGTGAGCAGGGGG
GCTTCAATCGCNCTNTCGATNTGCGACAGNGACATNTTTTATCTTCGACNAT
20 GCNCTCNCTCCCTCCCACAGAAATCTTGCGCTNGNTCTCCGANNTNGGGNTNG
ANGGCNCTCTTCTCTNTCCTTAAATTGGGANTTNNCTTTTTTCNAANAAGGGN
NAGA

Rescue ID EcoR1

Rescue Sequence 2

25 AATCNTTTTNTCCATTNGGCGNCTTNTCAAAACATATTCACATTTGGNCCCAA
CGGCGTANGACTTNATCTCACGATTGTTTGGTTTCCTACTCTCCCGCGCTCCCT
CTCTTCTGAGTCTCTTTCTGGCTGATTTCGATTTCGATTTTAGCCGCTGCCATCG
CCGTTGTTTGCCTACCTATGTGTGTGTGTGAGGAGTGTGTCTTGATTTTCAGT
30 CCGCAATGCGCTCCGCTCATTATTTGTTTGANCGCCGCGGTGTAAAGTTGTAA
AAAGTCCAAGTGCTCGTGGAAGCTCGATGCAAGACGGGGAAAACGAAACGCG
ATAAATCGTGAGAAAAAGAGAGTGCCTAAAGGAAGAGGGAGTGATAATCAN
ACGAAATGGAATAATGTNTTGCAGAGGCNACAACAACATGCAAATAGTTG
TCATTGAGGCGCAATGAATGATAATTAGTGCTTANTTGAAATCATAATCNTGA
35 AGAAAGCGTAAAGCTCGATTNTGGCAATNTATTCTTGATTACCANTGAGTCTG
TGATATTGCCGTGTGTNCCGAAAATGGANGTTATNAAACCCATGGACTTCAGC
ACCTTCTCCGCGTTCTGCGAACATCTTAACAAATCTCCACAAAATTGCAGCAA
CAACTGCANCGACGGTACCGCCAATAAANCAATGGAAAANGCATTATTTG
GAGGTAANAGCNAAAAATACCAATNTTCCAATGCGAAATTGCNAGCNTGG
40

Genomic hit, Accession No. AC004274

Annotated *Drosophila* genome genomic segment AE003590

Annotated *Drosophila* genome Complete gene candidate CG13693 - novel

45

Human homologue of Complete gene candidate 6e-05 4507659 translocated

promoter region (to activated
MET oncogene)
>gi|1730009|sp|P12270|TPR_
HUMAN POOR MATCH

5

Putative function No homologies to indicate function

Confirmation by RNAi Only wild type profiles observed

Example 62 (Category 5)

Line ID 162/24
Category 2nd chromosome, small imaginal discs
5 **Reversion** R
Map Position 55C

Rescue ID EcoR1

Rescue Sequence 1

10 TTTTNTTTCANGGNTCTTTGCNCATAAAAANACACGNGCCCTCNTGTCCATTTCAC
ATTTTACTTGGAGTCGGTAACGTTGAGTTCCGCGTCCGTGCGTTCTGCCTTCCA
ATACAAAGTCTGGTGTGAATCTACCAAGCATTCCAGTGNGAAAATCAACTCAC
ATTGCTCGGTGATCCNTGCGGCGGTATNATCGCACCCGGAATTGCATAAGTTG
CGGNGAGCGGAAAGAGAGTGCACGGATTTCNGTTATCNAAGGGCCGGCANC
15 NGTGGGGCGGCGACGGNAGAGCACGCAGAANAANAATANANTGNNGTGGCG
AATTNAAAAATANNATNAAAGAAAATTCGGGCGCGCTAATTTTTCTTCAAATTT
GTGTGCGGTGCGCGAAAAACAACGTGTTTTCNATGGTTGATAATACACACGG
ACGGNNCACTCGCGCTCACCCACATAGTCACNAAAGTCGGCGACGTCGACGA
CCNCNACNCTCACATANGGACNTTTAATCCCGTNCATNCGTGTAGCGTNCNTA
20 TTTAACCNTNTCTGTCCATCGGAACGCNCGCNTTCTCGCCTTCNTTCTNCTTTA
CTTTAATTTCTATTTNNAAGGGGNAGNCCNATCTTTTNCCTNTCNNTGCCNT
TTAANNTCATCCACANCTCNCTTTNTCNTTCCCTCCNCCTTNTNTCTTTTCNTC
TTNCTTNTGNCCTTGCCCTCGTTCCTTCTCTCNTCTCCTTNCCCTTCTCCTCCTTT
TTTCTCCTTCCCCC

25

Rescue ID BamH1

Rescue Sequence 2

AAGNCNCCTTGGCCGNNTTNAACGGNAANTAANCCGGGNCNCNCGGNCNCGA
TAATCAGGTCNANCCTTGTGCCTACCACCACCAAATTGAAAAAGAGCNAAGA
30 TTCTCTAAGGCAAAAACTCCCCAATCTGTGGAATTTCCGGAAGCGAGAGCAC
ATTCAAAGCTACCAGTTATCAGCGAGCAGCATGTCTAAGCTCAGGAACCTGTT
GCCCACAATCTTTGGCGGGAAGGAGGCACAGAATCCGACACCCGTCGAGGGA
CGCTGGAATAGGACGCGAGCTCCCGTGGACGACAACGAACCNGATTACTACT
ACTGCGGAGCCATGGCGCTGCCCTCCACCGCTGGCACGCCCACAGCCTCCTCG
35 GATCTGACCGAATCCGTGCTGCGCGAGCTCAGCGACCCAAACTACAATTCAAT
GGATGTGGTGCTTTTCNNCCTNTTTTCCGGGCACTCTCAGTAACGTCCAGACAA
ACAACACCATGAACGTTACNCGCGCCAGCAACAGGTGGTCATGAACCTTCTCG
AATGCCAATAATCTGCACTTCGGCTCCGTCTTCAACTTCAACCAAACTTGAG
CGCTGCGNGCTCNCGAANGGGTTTACCNGTTCGCANAAGAATCGGTGCGCTC
40 TCCANACNGT

Genomic hit, Accession No. CSC:AC012981

Associated ORF

45 Genscan ORFs: ORF2 predicted sequences
>18:26:17|GENSCAN_predicted_peptide_7|1320_aa

MEETNNATTIEQQPIALINGQEQVANEQQPSSPTSVATPTSTTSGGTGNATPAFSY
 DDLFPALPANTS AQSGASGSLARVTSSQKTHIVHVPCKERKSTESKFGEGES
 KRICQKITKETGAQIEIASRQVTPREHFRVILGKGGQRLREIERVTATRINIPSQSD
 ESEFITIAGTKEGIAQAEQEIRQLSAEQYKKSSDRITVPKVYHPFIVGPYSENLNKLQ
 5 EETGARINVPQQVQKDEIVISGEKDAVAAAKAKVEAIYKDMKKCSTVSVEVAK
 PKHRYVIGPKGSTIAEILQLTGVSVEMPPNDSPSETITLRGPQVALGNALTVVYQK
 SNSVKSVEINAAHWHKYVFGRKGANMKQLEEDCPNVNVCLEDKIKLEGDPEN
 VDRAVAYLSEIKNYEENFTFEVMTVNPSSYKHIIGKAGANVNRLKDELKVNINIE
 EREGQNNRIEGPKEGVRQAQLELQEKIDKLENEKSKDVIIDRRLHRSIIGAKGEKI
 10 REVKDRYRQVTTITPTQENTDIVKLRGPKEDVDKCHKDLLKLVKIEQESSHIEVPI
 FKQFHKFVIGKGGANIKKIRDETQTKIDLPAEGDTNEVIVITGKKENVLEAKERIQK
 IQNELSDIVTEEVQIPKYYNSIIGTGKGLISSIMEECGGVSIKFPNSDSKSDKVTIRG
 PKDDVEKAKVQLELANERQLASFTA EVRAKQQHHKFLIGKNGASIRKIRDATGA
 RIIFPSNEDTDKEVITIIGKEESVKKAREQLEAIKECDEVTEGEVSVDPKHHKHFA
 15 KRGFILHRISEECGGVMISFPRVGINSDKVTIKGAKDCIEAARQRIEIVADLEAQT
 IEVVIPQRHHRITIMGARGFKVQQVTFEFDVQIKFPDRDATEPVEGLTNGGSGENG
 GENEGQGEQEVEKEAEQEPVRQCDVIRITGRIEKCEAAKQALLDLPIEELSVPF
 DLHRTIIGPRGANVRQFMSKHDVHVELPPSELKSDVIKVCCTPARVAEAREALVK
 MIEDYEADRADRELRSFVLQVDVTEFHSKLIGRHGAVINKLRADHDVVISLPKRD
 20 EPNDRIISITGYQANAEAAARDAILEIVGDPETLHREVIEIDKRIHPLIGQRRRTIRKII
 EDNKVNIKFSADDDNPNISIFISGKIEDVENVKELLFGMAEDYERDYLNDVAIAPPTI
 GAFLTGFWIRCRRCQRRERIRHQRRTVGEAKAGQKPDCAQHSVAGGLPALRCWRG
 SGGHLAYHLRVGPQKLSASGRVSRSPA VAAILQVGVRRGSELEMDQELEQKLELE
 LEDYRAMSGRAAAVVRTSL

25

>18:26:17|GENSCAN_predicted_CDS_7|3963_bp

atggaggaaactaacgcaactaccatcagcagcagccatcgctctcattaatggccaagagcaggtggccaacgagca
 gaaccatcctcgccaactcagtgccacgcccactagtagcggcggaactggcaatgccacaccgccttagctac
 gacgacctgttccggccctgcccggccaactcggctcaatcgcaatccggagcttccggttcgactctagctcgtgacgag
 30 ttcccaaaaaactcatattgtgcatgttccctgcaaggagcgcaagtcacggagtcggagaagtttggcgaaggcgagtcgaag
 cgtatttgcagcagatcaccaaggagaccggagccagatcgagattgccagtcggcaggtgaccgttccctgggagcacttc
 cgcgtcatcctcggaagggtggccaacggctgcgcgaaatcgagcgtgttactgcgacgcgcacatcaacatcccagccagag
 cgatgagagcgagttatcacgattgccgaaccaaggaggtattgccagggcggcagggagatccgtcagctgtcagccg
 agcagtacaagaagtcacggaccgcatcacgggtgcccaagttaccatccctcatcgtgggcccctacagcgagaacataaa
 35 taagctgcaggaggagaccggcgttaggatcaacgtcccgcgcagcaggttcagaaggacgagatcgtcatctcgggcgag
 aaggacgcggctgcagcggcaaggccaagggtggaggccattacaaggatatgaaaagaagtgtctaccgtcagtgaggga
 gtagctaaagccaagcaccgatattgtcattgtccgaagggtccaccatcgccgagattctgagttgaccggtgtgtctgtag
 agatgcctcccaatgactccccctcgagacgatcactttgctgggccgaagtggctttggaaatgccctaaccgttgtctac
 caaaagtccaactcggcgaagtgttgagatcaatcgccgacattggatccacaagtatgtgttcggtcgcaagggggccaaca
 40 tgaagcagctggaggaggactgccccaacgtgaacgtgaattgctggaagacaagatcaagctggaggggagatcccgagaa
 cgttgacagggtgtgacctactgtccgaaatcatcaaaaactacgaggagaacttcacattcgaggtgatgacggttaaccttc
 gtactacaagcacatcatcgtaaggctggagccaacgtaaatgcctgaaggatgaactgaaggttaacattaacatcgaagag
 cgcgagggccagaacaacatccgatcaggggtcccaaggaggagtagggcagggcgagctgaattacaagaaaaatcg
 acaactggaaaacgaaaaatcgaagatgtgatcatcgaccgctcctcatcgttctattatcgagctaaggggcgagaagatt
 45 cgcgaggtgaaggaccgctaccgccaggttacaatcacgatactacgccccaggagaataccgatattgtgaagctgcgcgg
 acccaaggaggatgtggacaaggtgcacaaggtatgcttaagctggtaaggagattcaggaatcgtcgacattatcgaggtg
 cccatcttaagcagttccacaagttgattggcaaggggcgccgtaacatcaaaaagatccgcatgagaccagactaaaat
 tgatctgcctccgagggtgacaccaacgaagtatcgaatcaccggcaagaaggagaacgtgctcagggcgaaggaaacgta

tccaaaagattcaaaacgagctttccgacattgtcaccgaggagggtgcaaatcccgcccaagtactacaactcaatcatcggcact
 ggcgggcaaacatctcctcgatcatggaggaatgcgggtgtgttctatcaagttcccaacagcgactccaagagcgataaggt
 cactattcgcgggtccaagacgatgtggagaaggctaagggttcagctattggagctggccaacgaacggcagctggtctctt
 accgccgagggtgcgcgccaagcagcaacaccacaagttcctgatcggcaagaatggcgcttctatccgtaagattcgcgatgcc
 5 actggtgcccgcattatcttcccttcaaacgaggacactgacaaggaaagtatcaccatattggcaaggaaagcgtaaaga
 agggccgtgagcagctggaggcgatcatcaaggagtgcgacgaagtaaccgaagggtgaggtttctgtcgtatcccaagcaccac
 aagcacttcgtggccaagcgtggcttcatcctgcaccgcatctcgaggagtgcgccggcggtgatgatctccttccccgtgctgg
 catcaactccgataaggtgacgatcaagggtgccaaggactgattgaagcgcccgccagcgcatcgaggagatcgctcgccg
 atctggaagcgagaccaccatcgagggtggtgattccacagcgatcatcgcaccatcatggcgccacgtggttgaaggttca
 10 acaagtcacctttgagttcgtatgtgcagatcaagttccctgatcgtgatgccaccgaacccgtcgagggtctgaccaacggaggc
 agcggagagaatggaggcgagaatgaaggccaggaggagagcaggaagtagagaagggaagccgaacaggagccgggttc
 gtcagtgcgatgttatccgaatcacgggcagaattgagaagtgcgaggccgccaacaggctctgcttgatcttccccatcgag
 gaggagttgtcgggtgcctttcgacciccatcgtaccatcatcgaccgcgcgggtgccaatgtgcgtcagttatccaagcacgat
 gtgcacgtagagctgccacctagtgaagcttaagtcggatgtgatcaaggtctcggtacgcccgtcgcgtcgccgaggcccg
 15 gaagcgtggtgaaaatgattgaggattacgaggtgataggccgatcgtgagctgcgcctcttgttctccaggtggacgtaga
 tacggaattccattcgaagctcattggtcgtcatggcgctgtgatlaacaagctgcgtgcccgatcacgacgtcatcttctgcct
 aagcgggatgaaccaatgacgcacatctctatcacggctaccaggccaatgcggaggcagcccgcatgccaatcctaga
 gattgttggcgaccccgagacacttcacgcgagggtatcgagatcgataaacgcatccacccccacctcattggccaacgccga
 cgacacattcgaagatcatcgaggataataagggtgaacatcaagttctcagctgatgatgacaaccccaattcgtatctcatagt
 20 ggcaagatagaggacgttgagaacgtcaaggagttgctcttcggcatggtgaggactacgagcgtgactacttgataacgtg
 gcgatagcggccgccaacgattggtgccttctaactgggttcttgatccgatccgcaggtgccagcgagaacggattcgtcatc
 aaagacgcaccggtgggagaagcaaaagcaggccaaaaactgactgcgccaacactcagtcgagggagacttcccgcact
 tcgctgctggcggggctccgggtggcctccacgcctatcacctccgtgtggggcccaaaaactaagtgcacggccgagtgctc
 ccgatcgccagcagtagcagcaatactacaagtcggggtgcgccgggagtcggagctggagatggaccaggagctggagca
 25 gaagctggaactggaactgaattggattatcgggcaatgagcggcagagcagcggcagtcgtgcggacatctcttag

- Drosophila Gene Hit** BLASTN with rescue sequence 1: dodeca-satellite protein 1 (DDP-1) (AJ238847). TBLASTN with ORF2:dodeca-satellite protein 1 (DDP-1) (AJ238847).
 30 **Drosophila EST** GH20785 (AI389573), LP07358 (AI294065)

Annotated Drosophila genome genomic segment AE003799

- 35 **Annotated Drosophila genome Complete gene candidate** CG5170 - Dpi dodecasatellite DNA binding protein
 CG5576 - Bg5 involved in cytoskeleton organization and biogenesis which is putatively a component of the plasma membrane
 40

- Human homologue of Complete gene candidate** CG5170- 4885409
 ref|NP_005327.1|pHDLBP| high density lipoprotein binding protein
 45 >gi|2498434|sp|Q00341|HB

5

CG5576- 2e-07 4506539
ref[NP_003795.1|pRIP|
UNKNOWN >gi|3426027
(U50062) RIP protein kinase
[Homo sapiens]

10

Putative function CG5170: DNA binding protein (homology with Scp160p, a new yeast protein associated with the nuclear membrane and the endoplasmic reticulum, is necessary for maintenance of exact ploidy)
CG5576: death domain containing protein, possibly involved in signal transduction

15

20

Confirmation by RNAi CG5170: Reduced G1 and G2/M peaks indicating fewer cycling cells and more polyploidy
CG5576: Loss of G1 peak

Example 63 (Category 5)

Line ID 40/2
Category 2nd chromosome, small imaginal discs
Reversion NR
5 **Map Position** 39B

Rescue ID BamH1

Rescue Sequence 1

TTTTGCCTCCGCTTTTAAATTAAAAAAATGTNTGTTTNGCCCTGGAGCTCTCG
10 GTCTGTTAGCGAGCGTTGCCACCTTTCTGCGAGCTGTTGCTGCACACTGCCACT
TTACGAACACAGCTCTGATAGCGGGACAAAATACGTCAAGGCAGCGACGGTG
GGTTACTAGTGAATTTGGAACGGTGGTCTTAAGACGTACTGGTCTTTTATATTT
TCATTATTTTTTAAATTGTCGCTCATTTACCAATAAACCTTTTACTTTTTCTTG
ATAGTCCGAAGTCAGATCAAATAGGAAGTTTCACAAAAAATTTTCATCCAGAG
15 AAAATACGCCGACGCTATTCGAGTTTTTGTATTTCGTTAACCGGGAAAGAATA
GTTTCGAATTCGTTTCGCACTTTATCGATAGTAGATTGCTATTATGGAGCCCACTA
GTAAATTAATTAATTCAGACTGATAAAAGCGATCAACTTTTGTTAATGGGT
TTAANTCTATAATAATNCTTAGTCCAAATTGTNTCAAAGTAGTCGATAATTTAT
AATAACAGTTTTAGATGACCTCTAGGAAATAACTAATTACCCACATNCTTCAA
20 GAAAGTGTNNCAATTTGTNCTATAATTAATAACAGTTGTATTAAATTATGTTG
TNATTGTNACTCATAATACAAATTAACAATATAAACACACATAAATAAGAG
AATTGGAATATTTGTCTCAGATTAGATTTNCCAC

Rescue ID EcoR1

25 **Rescue Sequence 2**

AACGGGGGGGCTTCCGCGNCNCCAAAACGCAATNTACCGTTCATGCTGTGAAG
CGAAAAAGAGTGGTAGCGCCTACCNTGGCATATGTAGTTAAATCCGTGAAAT
AAGTGAATAAGAATATATGTATGTACTTAATTCGAAAACCTTTTCGCCGTCAG
CACAACGGGTGAACGAGAGAGCGGAAGTGGAGTTTTTTGTGGCGGGTCGTCT
30 CGCTCGCACC GCAAANGTCGTCCGTGGCTGCGTGTATGGGTGTGTGGA
GCGTCGAGGTGAATGTGGATTTCTAACCACACCAGCATTGCAAAGACATTGAT
TGATATTTAAAGCTGCAGCAGCGAACAAGCAAATCCTAATTTTCGGCAAAGTT
TAAGAATAACGAGTGAAGTGGGCGCGCGCAATAAGATAAAATTGAAGGTTAT
CTGTGTGCGTGTGAGTGACCGTNTACCAGTGTGTGTGTGCGANCGTCCATTGT
35 AAACAAAAACAAGTGTGTGAGCGGAGAGAAGAAAGGGAAAGAGAGAAAG
AGCGAACAGACTGGCGAGAGAAAAAAGAGATGCCACAAANAAAGCAGCGCA
CAAAGGAAAGCTGAAAATTTTCAATAAATCTGCAAAAGTGAAGAAAACCA
GAACCCGCAGTCNTGTAAATAAAACCCAGANTCCAAGAAACNTTAAAGAA
GCAGTGCAAAACAACTGGTGCTNTGAATGCGGTTTATTTGAAAAAAATGCA
40 ATTCGGTCCGATGGAA

Genomic hit, Accession No. CSC:AC014744

45 **Drosophila EST** several including LD46342 (AI544109 BLASTN similar to mRNA L07550)

186

Annotated *Drosophila* genome genomic segment AE003669
Annotated *Drosophila* genome Complete gene candidate CG8678 - novel with ankyrin
homology

5

Human homologue of Complete gene candidate CG8678 -gi7661580
B69CEC399B56F35C
[ref]NP_056425.1|DKFZP434J
154 protein [Homo sapiens]
(2.20E-85)

10

Putative function Novel protein with ankyrin domains, unknown function

Confirmation by RNAi Reduced G1 and G2/M indicating fewer cycling cells

15

Example 64 (Category 5)

Line ID 55/12
Category 2nd chromosome, small imaginal discs
Reversion NR
Map Position 49C

Rescue ID BamH1

Rescue Sequence

10 TCTCATGNTCAGGGGGCCTTTACNATGTCAAAGAGCAAATTGTCCACAGGGCA
 GCAACCGCAAGTGAGAGACGGGTGGAAAACCTGGGCGGCATGACCATGAATGA
 AAGCCGCGACCGGCAAACGTGGCCCGCCACAAAGCGAGCATTTTCACATTTT
 AACTGTCTGGACATTTTGTAAAGTTACACCAAGGCAATGATACCAGTAAAAAAG
 AAGAAACAATCATTTTTGAATAGATTAATCACCTGATTAATGTTGGTTGTATGT
 15 TGATTGTAGGTGTTTTAATATACAATGTCTCTATTACTGCTTTCCTTTATTCAA
 AGCCATGTGTAAGTGTAAAGTTCTCGATTTCGGCTAGATTTTGAAGTTCTGCCAT
 TATCAATTAAAAGTCCAGTTCCTCTATAAAATTGGTAATAAAATAGCTCTTTACA
 GCCAAGTATATGTGCAATTTTGTAAAGATTAAANGTCCAAATGTTGTGAACCTT
 TCCTGGCCCTGAATTTTAAAAAACCATTAAATTGGTCCCATTGACATTAAATG
 20 TTCTATGTACATTAATATGACTTTTGTGGATGGTTTTATAACAAGCATTACT
 ATATTCTAAAAATCAAGGATAAAGGACNAGCTTTACAGGAGGTAAACATTCCTA
 TTGTACGGCTTTATTTTCTTATACCCATAAGAGCATACCACTAGGATCCGTCGA
 CCTGCAGATCTCTAAAAACTTGCCTTTGCTGGCGTTTTCCATAA

25 **Genomic hit, Accession No.** AC007085

Associated ORF

Genscan ORF1 predicted sequences >21:54:11|GENSCAN_predicted_peptide_3|108_aa
 MGLVTAAFKLKRKDIQDRYQHDINRICHTRSTAHTAYAHFAEHLRRSPRQRFVN
 30 GKGAALVLILLVSAARQFSGSTGAYKLGNRVGVKVEGEQQEYKLQDRTTTHFCGN

>21:54:11|GENSCAN_predicted_CDS_3|327_bp

atggggctggttaaccgccgcttcaagctgaagcgcaaggatatccaggacagatatcagcatgatattaaccgcatctgccaca
 cacgtagcacggcacacacggcgatgctcatttgcggagcatctgtgacgaagtccacgtcaacggttgtcaacggcaa
 35 aggtgctgcgcttgctcatcctcctcgttctgcggtcgacaatttctggctcgacaggtgcctacaaactgggtaataagagttg
 gaaaagtagaaggggaacagcaggaatacaaaactacaagacagaacaacacattttgtggcaattaa

Corresponds to CG8732

40 **Annotated *Drosophila* genome genomic segment** AE003836
Annotated *Drosophila* genome Complete gene candidate CG8732 - l(2)44Dea
 homology to fatty-acid-
 Coenzyme A ligase, long-
 chain previously described
 45 spindle/chromosome

abnormalities in neuroblast
squashes

- 5 **Human homologue of Complete gene candidate** 1e-171 4758330
ref|NP_004448.1|pFACL3|
fatty-acid-Coenzyme A ligase,
long-chain 3
10 >gi|4165018|dbj|BAA371 and
LCFD_HUMAN LONG-
CHAIN-FATTY-ACID--COA
LIGASE 4 1e-157
- Putative function** Fatty acid CoA ligase
- 15 **Confirmation by RNAi** Only wild type profiles observed

Example 65 (Category 5)

Line ID 6/7
Category 2nd chromosome, small imaginal discs
5 **Reversion** NR
Map Position 28E

Rescue ID BamH1

Rescue Sequence 1

10 TATNAATAATCATAGGGCTCTTGCTCTTACGTGTAAGGCCTGCCCCCTCTNCCA
GTCTATATACAAAGAAAAACACACACACACTGGCACACTGGTGTTCGCATATG
CCAAAGCCGAGTTAATTTCACTTTGTTAATCTATCGTTTGGTGTTCGATTT
TTTAACCGCGCAAACGGTATTTGCGCGTTTTCGCGCTCTTACTTTGCGATTTAT
TGCACCGCTTGGCTGTGTTTTCGCAATTTCTATCTTGATTTTCATTGGTATTCACG
15 CGTAATGTAATTTCTTAGCAGCGTGACCGCGCCGATAACGATAAAAAATACCAC
GGGACCAAAAATAAATACCATATGATACCACTTCAGGGAAAAGAAATCCTAT
TTAATACCACTCACTTTAAAAATAAGTTTTTAAAAATATATATNTTTATTTAAA
AAAAGGTGTATTTATAATCAAATACTCGGTACTTNTAATTACTCCAAGAANA
ATTAATTTGAAAAAAGGGGTTCATTATAAAATATATATTAACCGCTTACAC
20 ATAATCCCCAAACAAAACAGCGATTGGGATTTAAAAGGTTCTAAGTCCATCAT
TATAAAAGATCATTTCCGAAAAACAAAAGAAATAGATTCAAAATTAGGCGAC
ATCAGCCCGCTGATAANGATCATAAAAATACAGAAAGCTNATTCAGCGAATCA
GAAANTCCTACTCGCCACTATCCGAAAACNTNGAAAAAAAATGG

25 **Rescue ID** EcoR1

Rescue Sequence 2

TGAAAGGTAGCAACAACGTTTCCTTGAAAAAGCTGTAAATAGTAAACAAAA
TTGTCAAGTTAACGAGCCAAAGTTATTAATAAGGTTTCGAGTACGTTGGCATC
GGCTGCCCAGGCAGCAAANAAAAACAAAGACGCAGTTCAAGATCAGCTGGAC
30 ACTTAGAAGANTTTAAGAATTGAAGCACATTNNAAGAAGANAAACAAGAAC
CCCACCAAAAACCCGCGTGCCTTTGTATGTGTGTGTGCCATCAAATTTCCCGC
ACTGGGTGAATGTGCNTGCGTGTGTNTGTGTGCATTTAATTTTCCTACCAATAA
TCGCCTTCCAAGAAGTGAATACCAGCCGATCCACCGCTAAATCGAAAAAAGTT
TNACTCTGGGTTAANTCACTGTTTACGGCTTTTGTGCTATAATTACCTTTCCCG
35 TAAGCNGTGGGAANCTAAAANCCAAAACNTNAGAATCCGAATTCCG

Genomic hit, Accession No. CSC:AC017934

Associated ORF

40 Genscan partial ORF1 predicted sequences
>22:35:21|GENSCAN_predicted_peptide_4|128_aa
MGTNSGATAGINNKPVGATGAGVLVGGGVGGANSSIGGVLSNSLGGGGSGGLS
ISGLNAGGQNAVGGMGNVGGDDGGNGMVGGGVNNQQATTPQYTIPGILHFIQ
HEWSRFELERSQWDVDRAELQ
45 >22:35:21|GENSCAN_predicted_CDS_4|384_bp

190

atgggcaccaattcgggagccaccgctggcataaacaagaacgggtggcggtgcaacaggagccggcgctctgttaggcg
 gcggtgtggcggtgccaattcctcgatcgcggtgtcctgtcgaacagcctggcggtggcgagcgcggtctgagcatc
 agcggcctcaacgctgggtggacagaaacgccaatgtggcggaatgggcaacgttggcgcgacgacggcgaaacgggatg
 gtggcgcggtgtaaataaccagcaggccacaacgcccatacacaataccgggcattctgcacttcatccagcacgagtg
 5 tcgcgttcgagctggagcgatcacagtgggacgtggacagggccgaattgcag

Human Homologue TBLASTN with ORF1: very weak homology with striatin,
 calmodulin-binding protein (STRN) (NM_003162.1)

Drosophila EST several including LD42534 (AI516610), LD03224

Annotated Drosophila genome genomic segment AE003619

Annotated Drosophila genome Complete gene candidate CG7392 – novel WD40 family
 member

15 **Human homologue of Complete gene candidate** CG7392- SG2N_HUMAN
 CELL-CYCLE NUCLEAR
 AUTOANTIGEN SG2NA
 (S/G2 ... 622 e-178 A cell-
 cycle nuclear autoantigen
 20 containing WD-40 motifs
 expressed mainly in S
 and G2 phase cells

25 **Putative function** WD40 protein a novel nuclear protein mainly expressed in S and
 G2 phase cells that was characterized using autoantibodies from a
 cancer patient

Confirmation by RNAi Reduction of G1 peak , more polyploidy

30

Line ID 103/1

Category 2nd chromosome, small imaginal discs

35 **Reversion** R

Map Position 57B

Rescue ID BamH1

Rescue Sequence 1

40 GATTTCAAAATTAGGCGACATCAGCCCGCTGATAAAGAATCATAAAAAATACT
 GAGGCTTATTTTAGCGAGTCAGAGACTCCTACTCGCCAACATATCGAAAACATA
 GNGAAGATATAGTCGCCAACCGATCTGCCTTCTATAGTGTTGCTTATTGTTGTC
 CCCTAATCAAATTAATAAAAAATCTGCATTAGGCTGCTTCGCCGGCCAGCAACA
 AATGTTTACACCTACTGTACTTTTCGCAGAACAGAGATCCAAATGCAGGATC
 45 GTTTCATGACTGTACATTTATTTCGGATTAGACATTAAATTACACCCTACAGCT
 ATACATACTAACAGTGAACACGGCAAATGCTTAGCTAGCATTGGGCCACTTTC
 GTTGACTGCGAATAAAAAATGATTGGCCGATGCCTTTAGCAGATTCTTTTGAT
 CGAATTACTCGGATGGCTTGTGTGTCCACCTCTTACAAGAACTCCTCGCACCA

ATCGTTGAGACAGTTGTAGCAATCGGATGCTTGGTTGGAGCTGGCGTGGCACA
 CCTTCTTCATCCAGTCCCTTGGACAGNTTCTTGGNCCTTTTCAGNANCAGGATCT
 GGTCCCAAACGGNGGAAGGCCTAAAACGAATGGNAATTGATCGGTAGCCCTT
 GACTGGCATTGGTAATTTGCGCACATGGGNGTCATCGGATTTACACACGCACC
 5 ATATCGAATCAGCGTCCTTAAGCGTCAACCGAGGGTTTCCCCAATTCCGGCCA
 GTTCCGTCACCGACTTGGTTGCCATTGG

Rescue ID EcoR1

Rescue Sequence 2

10 ATCAAAGCGNCTGGGCCCCGTGCATCGCCNCAGCGTTCGTCTTAATTAATTAGT
 GATTGCAAGCGGGTGCAATTATGCACAAAATTACGGACTAATACAACGCCC
 GCTTCGCGCTCTCTCCATCTCCCTTCCAAATAGTCGTTTGCTCTTCGCACACAA
 AAGTGTAACCCTGTGAAAGGTAGCAACAACGTTTCCTTGGAAAAAGCTGTA
 AATAGTAAACAAAATTGTCAAGTTAACGAGCCAAAGTTATTAATAAGGTTTCG
 15 AGTACGTTGGCATCGGCTGCCCAGGCAGCAAAGAAAAACAAAGACGCAGTTC
 AAGATTCAGCTGGACACTTAGAAGAGTTTAAGAATTGAAGCACATAAAAAAG
 AAGAGAAACAAGAACCCACCAAAAACCCGCCGTGCGTTTGTATGTGTGTG
 TGCCATTCAAATTTCCCTGCACTGGGTGAGTGTGCGTGCGTGTGTGTGTGTGTC
 AGTTTAATTTTCTACCAATAATCGCCTTTCCAAGACGTGATTACCAGCCGATC
 20 CACCGCTTAAAATTGATAAACGTTTTAACTCTTGCGTTACATCAGCTGTTTTAC
 GGCTTTTGTGCTATAAGTTACGCTTTTCCCGTAAGCCGTTGGCAACACTAGAA
 CGCAAAAGAGCATAAAGAATCGCGAGTACCGTANAGAGGAAGAGAGGAAGA
 GAGAGAGATAGAGAGTGTGAGCGTGTGAGTGAGCGGGGAATGTGGGGGCGGT
 TCCGGTGCGAAAAAACGTAGTAGTAGTACATNNAGAGAGTGCGAACGAGAGG
 25 GAGGCAGCCAGCGAGTGTCTGCGACTGCTCCCCCTTTACCCTCGTCGCTTTT
 CTATTCGGAAAAATTCAATGACCTCATTTGTTTCATGTGCCGAACTTTGCTTTTC
 TTTCCCAACCTAAAAACGCAAAAAAAAAAAAAACNCCAAACAGGATATACGTNG
 GAACANTGANCAAAACNANTTCGANAAAACCAACAACNANGGACCGTGCCCTG
 GGGCNCCTGAAAGGCAAACAGCTGGCNCNCAATCCGGAAGAGGATCNGGAA
 30 NAACAGGATCNGCGGGCNCAAGGATCNCCGGAACAGGCAAAGGAAACNCCC
 GGCNCACNGCACAAGCCNCTGAAAAGCAACNTGAACCAATGGGCACCANTTC
 CGGGANCCACCGCTGGCATTAAA

Genomic hit, Accession No. CSC:AC017934

35

rest of results as for line 6/7

Example 66 (Category 5)

Line ID 65/24
Category 2nd chromosome, small imaginal discs
5 **Reversion** NR
Map Position 48A

Rescue ID BamH1

Rescue Sequence

10 TACGATTTTTGCANTGCNCCATTTTCGTGGCACCCGATTTGTATATATATTTTTT
ATATAACCCACGGATTGCCAACTTTCATTGCCCTTTCACACTCTTATTCGCCAT
TTATGAACTCTTCTTTGACGATTGGAACGGTTCCTTTTCGCTATTTTCGACTGC
ACCCGCGCTCTTTTCGCTTCGCTCTCCTCCCTCTCTACACACCGCTCTTTATCCT
TAATTGCTTTTTCTATTTAGCGGAATTGATCGTTCTCAACTTGGTCGCCATTGC
15 AGCTCCACAGGCGAAAAAATCGGTGGAAATGCCAATACAGGTGCACGGCGAG
TGCCGATAAGCTGGAAAATCGGGAAAACGCACGCCTACACATTTCATTGCCAG
CATCGGCTTTGCCTTTTCGCTGTGCGAGATTAGCATATTTCCACTTTTGGTTCGC
GCACAACACTANCTAAATTATTGNTATTTTTTTCCCAACTGTGAGGTGAAAC
TGTGAAACAAAACCACTGTGGGCGGGTCAGTGTGACCCTCTCGCGGTGGGTG
20 AAAATCCTAGTGAGCTTCGTTGTTAGGGCTGTATGACACGAAAGCAAGTTGAA
AAGAACTTTTTTAAAATTATTTGGTTAATTGAGCAGAACTAAACTATATN
AAAATATTTAAGAATNCAGATTAGTGATGTATTTAATATAATAATAGTAAGAT
GTTT

25 **Rescue ID** EcoR1

Rescue Sequence 2

CTTNTTTGATAGANATAGGCTTCTTTTAAAAAANAAGCAGCANCAGGGG
CCCNAAAGTGCGTGNNTGTGAACGCTGATTGCTTGCAAGTGTGTTTCGTGTGTG
TGTGATTGTGTGCTCCGANCAAGTGAAATCAATAATATTTGCAGCCACAAGCA
30 ATTAATAAAAACTGCAATAATGTCAAAAAATCTAATTGAGGCAACAAATTAN
CAAAGCCATNAAAGCAGGCTGCACTGCGAGAAAATTGTGCCTTTCCACAGAT
CTTCTGCTGCAAAAGCNAAAGAANGTAAGCAAGTCGGCCANTTTATTNCATTCT
TCTCATCTCTCTTCTTCGCGAATTGGCGCNTANCACTTACAATAATTNATATNA
CTTCTTAAATTTCAAANTCCCTTTCNTGAACGGANCTTTTAACGGAAAACAAA
35 GCGGGTAAACTAACTTAACTAACTAATTANAANTGTANGTATAAATGAACC
GAACTCGCTTTAGATATNATGCGTTTCACTAACANATTANAACAACTTTGAA
GCTGTANTGTCAGGTTGTTATTNCGTTCACCANATGTAGACTGNCCGNNAATT
TNACCTTTCCCATANTCTGTTCTTAANTGINTTGTTTTTTCCCAATNNTTTGATC
ATNCNTTGGTNAATNANCTNAACGGCCCAAAGTNAATGAATTCCANTCACGTC
40 CACTGGCTCTGGTTCNATANTTAATNGGCTGTTTCTTACTTCCCTTAACCCTAA
CATCTNTTAATCACCTGTGCCATNTGTTTGTGTGTGTGTGAACGAATGAGAAA
AAAAA

Annotated *Drosophila* genome genomic segment

AE003825

Annotated *Drosophila* genome Complete gene candidate CG9005 - novel putative cell adhesion

5 **Human homologue of Complete gene candidate** CG9005- Ensembl predicted gene
ENSP00000006008
Gene:ENSG00000005238
Clone:AC004472
Contig:AC004472.00001 6.00E-38
10 (KIAA1539 protein AB040972) and
AK022837 Homo sapiens cDNA
FLJ12775 4e-33

15 **Putative function** Putative cell adhesion protein

Confirmation by RNAi Reduced G2/M peak

Example 67 (Category 5)

	Line ID	74/3
	Category	2nd chromosome, small imaginal discs
5	Reversion	NR
	Map Position	47A
	Rescue ID	EcoR1
	Rescue Sequence	
10	GCACAGAATGGCNCCTTCACGACAAAAGATCTNCNAATTAGGATGATGCAGA	
	AGGAGGACACGCTTTTCATTATCTGGTTGCCACCTAATTTAAGTTCCACATCAA	
	GGGAAGAAGGAAATACGTTCCAACGGACGTCAAATTTACTAACTACACTACTT	
	GAAAAGCCTGTCTATAAAAACACGATAACGTTTTTGCTAATCTCAAGACAATG	
	TTAAATATAATTGGAGAAAGTATTGAATATGAATATCACAAAAATTGTTTAGG	
15	GTCTCTACGTGGTAAATAGTATTTGGCATAGACAGTGAGATGTGAGTCGTACG	
	TACTAATTAATAAAAGTTGTTCAARAGAACCTCATATACTGTAAGTGACAACGA	
	ACGAAGCTGACAACCTCTGCTTGACATATTTGGCGGAGTTTCGAAAATATCATC	
	GCATTGGTATTGTTTTGTNTCCACCNTGGGGCGAGATTTTGTGTGTGCTTTAC	
	TTTGCTTGTTTTTCNCCACAAANCGAACCATAATGTTTCGAAATGGTAAAATTA	
20	CCGTGCCAACAAAGCTCTCTCTCTCCCACTCCGAACTCTCTCATCTCTCCTTG	
	CAATTGTTTAAGGTGTGCAAGGAAATGAAAAATGTCCCGGCTGTGTINCCATG	
	CATTCCCCTTCAAAGCCAATTATNTTTGTGCCTCTCCAACNTTTTTGATCGGNN	
	TGATTTTTTTGGCTCCCNANTCCCCCCCCCTTCNCCCATTCCGGGTTANAT	
	TATTNTNCCAATTTTCCTATTTTACGGTCCCNGTCCCTGGAAATANTTCCTNC	
25	AATCNCCGCTCCATNTCNCCATNTTTGACAGATTTTC	
	Annotated <i>Drosophila</i> genome genomic segment	AE003829
	Annotated <i>Drosophila</i> genome Complete gene candidate	CG12052 lola -a specific RNA polymerase II transcription factor involved in axon guidance
30		
	Human homologue of Complete gene candidate	1e-09 3789797 (AF059569) actin binding protein MAYVEN [Homo sapiens]
35		
	Putative function	lola-like specific RNA polymerase II transcription factor
	Confirmation by RNAi	Almost no G1 peak and increase in G2/M peak indicating arrest in G2/M
40		

Example 68 (Category 5)

Line ID 79/7
Category 2nd chromosome, small imaginal discs
5 **Reversion** R
Map Position 55B

Rescue ID BamHI

Rescue Sequence 1

10 GTCTCATGCACCCTGGCCCTNAGCTGCATAAGTGTAAGTGTGTGNCTGTGTGC
GAGTGTGGGTAGGCGGCGGCAACTATCTCGCTTGCTCTTGCGTCCGGGGTTAT
CGGTAGCTTCTTCTAGGCTGAGTGCATTTTCGTTGAATCGTGGATGTTGAAAGTT
GTCTAATTTCCGAACATTGATTTTTCCCTTCCCGTCAAGAACTGCATTGT
TGCTTCTTGAAGACCAGTTTTGGTAACATCAGGAGAATGGAAAGGAGCGAGT
15 GAGTCGGTGAGTAAGTGAGTGAGCGATGCGAGCGACAAAATCAACAACAACA
ACAACAACGGTCAAAACGAGTTCCAACGAAAGTTGCAACACTCTCAACAATT
TGAGCAGCTCCGTTTGTGTTATTGCATTACTCAATCGGGAAGACTCTACACTC
GACGGAATAGTGTGCTCGTCTGAAATTTATCNATTTCCATTCTTCCTTTGTTT
TTGGGCCAAACAATGGCNTCGGCAANCGTTCGTGGAAAACCGCAGGAACCAC
20 CAAAATGCCTGGCGTCACATTAACCGAGCCGCCTTTGTTTATGCAAATATTATT
GTAATATTTGGTNAAAATTAAGTCGCGCTTCNCGTTACTTTTTATTTTCATATAC
ACGCAGCAGCAGCACGCATACAGTCACGTCACGCACACATACAATCGCCGTN
CACATACACTTGTCTTTTTNCCACACACTTTCCTAATCAT

25 **Rescue ID** EcoRI

Rescue Sequence 2

NGNGTCTCATGCACCCTGGCCCTNAGCTGCATAAGTGTAAGTGTGTGNCTGT
GTGCGAGTGTGGGTAGGCGGCGGCAACTATCTCGCTTGCTCTTGCGTCCGGGG
TTATCGGTAGCTTCTTCTAGGCTGAGTGCATTTTCGTTGAATCGTGGATGTTGAA
30 AGTTGTCTAATTTCCGAACATTGATTTTTCCCTTCCCGTCAAGAACTGCA
TTGTTGCTTCTTGAAGACCAGTTTTGGTAACATCAGGAGAATGGAAAGGAGCG
AGTGAGTCGGTGAGTAAGTGAGTGAGCGATGCGAGCGACAAAATCAACAACA
ACAACAACAACGGTTCAAAACGAGTTCCAACGAAAGTTGCAACACTCTCAAC
AATTTGAGCAGCTCCGTTTGTGTTATTGCATTACTCAATCGGGAAGAACTCTA
35 CACTCGACGGAATAGTGTGCTCGTCTGAAATTTATCNATTTCCATTCTTCCTT
TGTTTTTGGGCCAAACAATGGCNTCGGCAANCGTTCGTGGAAAACCGCAGGA
ACCACCAAAATGCCTGGCGTCACATTAACCGAGCCGCCTTTGTTTATGCAAAT
ATTATTGTAATATTTGGTNAAAATTAAGTCGCGCTTCNCGTTACTTTTTATTTTC
ATATACACGCAGCAGCACGCATACAGTCACGTCACGCACACATACAATCGCC
40 GTNCACATACACTTGTCTTTTTNCCACACACTTTCCTAATCATNNTA

Genomic hit, Accession No. AC004296

Associated ORF

45 Genscan: ORF2 predicted sequences >15:31:31|GENSCAN_predicted_peptide_3|109_aa
MVTSEFRHLRDEKSFTDVTLACEGQTCKAHKMLVLSACSPYFKALLEENPSKHPHIL

KDVSYIHLQAILEFMYAGEVNVSQEQLPAFLKTADRLKVKGLAETPSSIKREG

>15:31:31|GENSCAN_predicted_CDS_3|330_bp

atggtgacctcggtccgtcacctgcgcgacgagaagagcttcacagatgaacactcgctgcgaggccaaacctgcaaagcc
 5 cacaaaatgggtgcttccgcttcagtcctactttaagcgctactggaggagaacccatcgaagcatccgatcattatcctgaaa
 gatgtctctacattcacctacaggctatactggagttcatgtacgccggtaggtgaacgtgtccaggaacaattgccagcatt
 cttagaccgccgatcgctcaaagtgaaggcctcgagagacaccagttcgataaagcgggaaggttga

10 **Drosophila Gene Hit** TBLASTN with ORF2: several zinc finger proteins including
 Broad-Complex mRNA for BRcore-Z2 protein (X54665)
Human Homologue TBLASTN with ORF2: kelch (*Drosophila*)-like 2 (Mayven actin
 binding protein) (KLHL2) (AF059569)

15 **Annotated Drosophila genome genomic segment** AE003800
Annotated Drosophila genome Complete gene candidate CG5738- lola, lola-like
 putative kelch-like putative
 specific RNA polymerase II
 transcription factor known to
 affect disc morphology

20 or could be CG10914 - novel
 unknown

25 **Human homologue of Complete gene candidate** CG5738- 9e-09 3789797
 (AF059569) actin binding
 protein MAYVEN [Homo
 sapiens]

30 CG10914- predicted gene
 ENSP00000051207
 Gene:ENSG00000047313
 Clone:AC068261
 Contig:AC068261.00019
 4.00E-49 (potential cell
 35 division GTP binding protein
 1: ENST00000051207

Putative function CG5738: lola like specific RNA polymersae II transcription factor,
 CG10914: Possible GTP binding protein

40 **Confirmation by RNAi** Both show marked reduction in G1 to G2/M ratio

Example 69 (Category 5)

Line ID 80/2, 81/8
Category 2nd chromosome, small imaginal discs
Reversion R
Map Position 57D/E

Rescue ID BamH1

Rescue Sequence 1

10 CANTTTCAGAGGCCATAGNCCTTCACAAAATTCNCCATCTCTGCCCCGGCATCC
 GTGCTTGAAAATGGTGCCAATGCGTCGTGGAGAATCTGCTGCACTCGATGGTC
 TGCAAAATTGCACATTTATTAGATTTAATAAAATTTTCAACTGTCCGCGANCAC
 GTTTGCTCGTGTTGAATTTTCGAGTACAAAATTAGTGCGACTGTTGGATTGCATT
 GAAATGCCAAAAATCGGTGTGACCATTTCGAAGTCCCCACAGGCTCATGACTT
 15 TCGCGGTTACCAAATCCAAATAACGCAAGCTGGTCACGCTGTCAAACATCGG
 TGACGGAATGGTGACGACACAAACAATTTGCTTAAAAACTTTCTTGCGGCCGT
 AAAAATGCGCAAGCAGCCTGGCAGCGCAACGCACGTACACGTAATTGGAACA
 AATGTTTGCTGAACCACAACCGCCCACTAAATGTTANCCGCCAAGTCTTTTCC
 CCCGCCGCCCGTCNTCNCNTCNCNCCGATTATTTGGTTTACAATTTGCTTAC
 20 ACAAGTGCAATCGTCGATAGCGCTTCATTTGGAGTAACAAGTAATATTTTGC
 GCCGTAAGTGTTCGCCGTATCAGACAGAAGGTTGGTATCAGTTCGACGCAG
 CTTGTGACGGTATTGCATACGCGGCGAAACGCCACGTGAAAACGGATCGCA
 GTTCTCGAAAACCTCNGGATAAAAA

25 **Rescue ID** EcoR1

Rescue Sequence 2

TGGGGTCTCANGCCCCGACGGCCATATTTTAAACACAAGATTCNNCANCTCTGC
 AGGGCATCCGTGCTTGAAAATGGTGCCAATGCGTCGTGGAGAATCTGCTGCAC
 TCGATGGTCTGCAAAATTGCACATTTATTAGATTTAATAAAATTTTCAACTGTC
 30 CGCGAGCACGTTTGCTCGGTGTTGAATTTTCGAGTACAAAATTAGTGCGACTGT
 TGGATTGCATTGAAATGCCAAAAATCGGTGTGACCATTTCGAAGTCCCCACAG
 GCTCATGACTTTTCGCGGTTACCAAATCCAAATAACGCAAGCTGGTCACGCTG
 TCAAACATCGGTGACGGAATGGTGACGACACAAACAATTTGCTTAAAAACTTT
 CTTGCGGCCGTAAAAATGCGCAAGCAGCCTGGCAGCGCAACGCACGTACACG
 35 TAATTGGAACAAATGTTTGCTGAACCACAACCGCCCACTAAATGTTAGCGCCA
 ACTNCTTTTCCCCGCCGCCCGCGGTCGTCNTCNCNCCGATTATTTTGTGTTACA
 ATTTGCTTACACAAGTGCAATCGTCGATAGCGCTTCATTTTGGAGTAACAAGT
 AGTATTTTGCGCCGTACTGCTGTTCCCGTATCANACAGAAGGTTGGTATCAG
 TTCGACGCAGCTTGTGACGGTATGCATACGCGGGGAAACGCCACGTGAAAAC
 40 GGATCGCAGTNCTCGAAACTCNGGATAAAAAGAAAAAGTAGGCTGAATG

Genomic hit, Accession No. AC007175

Associated ORF

45 Genscan: ORF2 predicted sequences >16:09:09|GENSCAN_predicted_peptide_3|2497_aa
 MNEGNSAGGGHEGLSPAPPAVPDRVTPHSTEISVAPANSTSTTVRAAGSVGAALP

ATRRHHQHIAATQVKGIASSSSKQQKQLASAPVPLSPLPQQQQQTAEATAAAAAP
AHSNVSVSSSTIEASVLPQAKRQRLDDNEDRTSAASIVGPAESSNIVSSLLPASVA
SSSEVGGLSSTALQDLNALKKRILQQKLQILRNLKERHLENVSEYFYLQNGGSM
DYPAWRKKTPTPQFISYSNANRIDQLIHEDKPSTSAAAAAQNQKYTTQQTDSVE
5 SSLVSGIGTGATKAPLDGNISNSTVKTNTQSQVPSKIGSFTESTPAATESNSSTTV
GTATSGAATSTSATSASGNVLAVEAEIKIPAVGATPVAISTKLPAAVVQLTQQG
GTPLPCNTSAGSTALRRPQGQNNASSGSAASGGGSLTPTPLYTGNPAALGG
SGGLTPGTPTSGSLLSPALGGSGTPNSAAQEFSEFKAKQEVYVMQRISLQREGL
WTERRLPKLQEPSRPAHWDYLLEEMVWLAADFAQERKWKKNAAKKCAKMOV
10 QKYFQDKATAAQRAEKAQELQKRVASFIAREVKSFWWSNVEKLVEYKHQTKIEE
KRKQALDQHLSFIVDQTEKFSQQLVEGMNKSVAADTPSLNSSRLTSPKRESDDDFR
PESGSEDDEETIAKAEEDAADVKEEVTALAKESEMDFDLNDLPPGYLENRDKL
MKEEQSSAIKTETPDDSDDEFEAKEASDDDENTISKQEEAEQIDHKKEIDELEA
DNDLSVEQLLAKYKSEQPPSPKRRKLAPRDPELDSDDDSTAVDSTEESDAATED
15 EEDLSTVKTDTDMEEQDEQEDGLKSLMADADATSGAAGSGSTAGASGNKDDML
NDAAALAESLQPKGNTLSSTNVVTPVPFLLKHSRLREYQHIGLDWLVTMNERKLN
GILADEMGLGKTIQTIALLAHLACAKGNWGPFLIVVPSSVMLNWEMEFKKWCPG
FKILTYYGSKERKLKRVGWTKPNAFHVCITSYKLVVQDQQSFRRKKWKYLILD
EAQNIKNFKSQRWQLLNFSTERRLLLTGTPLQNDLMELWSLMHFLMPYVFSSHR
20 EFKEWFSNPMTGMIEGNMEYNETLITRLHKVIRPFLRLKKEVEKQMPKKYEHV
ITCRLSNRQRYLYEDFMSRAKTRETLQTGNLLSVINVLMLQRLKVCNHPNMFARP
TISPFQMDGITFHTPRLVCDIMEYDPFTQINLETNLNLLHLEQTMTAYVSHKSRL
APRKLIEDIDTAPLPAPRPCNGKYRFHVRSAELAQRILNNAVKVGASPAMRLE
GSKIMPMRNLPSGRVLKRVASINPVNMAKPVVINSVVTSSSTTASSPTGAL
25 SVLSNSKLLGARSQINAPTPAKVAKTMQDGKPFYLTPTNSGAAGARLTLSKT
TASASTTTSRTTVTASTTSGQQLIRDPIVKDLATHVKSTVQKQSIANGKTEPEEETE
AEDPYKVQELIQMRKEQRLAALKRMAMINRRRTDATPIYGEDCREAIQRCMQAT
RSLKRSTWQTRGYANCCTAMAHNRGWSLNLHLLKSFEERCADLKPVFANFVIYVP
SVCAPRIRRYVQNLSSSTHWQHEQRIENIVDQALRPKLALLHPIISEMTTKFPDRLI
30 QYDCGKLQTMDRLLRQLKVNGHRVLIFTQMTKMLDVLEAFLNYHGHYLRDGS
TRVEQRQILMERFNGDKRIFCFILSTRSGGVGINLTGADTVIFYDSWDNPTMDAQA
QDRCHRIGQTRDVHIYRLVSERTIEVNILKKANQKRMLSDMAIEGGNFTTTYFKSS
TIKDLFTMEQSEQDESSQEKENKDRIVATTTLSSTPSTVVETEKQSLRAFEHALA
AAEDEQDVQATKTAKAEVAADLAEFDENIPIATEDPNAEGGPQVELSKADLEMQ
35 NLVKQLSPIERYAMRFVEETGAAWTAEQLRAAEAELEAQKREWEANRLAAMHK
EEELLKQETEAEEMLTYSRKDSSNQVNTKTDSSNSNKRRLVRENRRNSAQKLSRSV
SSHSTGSNNKNSKSATTRGNSQNSLNQTVPGSGISRVNRTGAGVSSSSRGKSNS
KSTGKGTDAAAPQVRRQTRLHSLGAVNMASARTPPTRKTTRTALAASAAASTLED
ASLIVEERPKRQSANIAMSKMMKTPFKQNVPSNISIKTTPPKRGRRDSVAAAATRS
40 KLLERRATIAAPLKHMDDESQDEEEQEEQSEEDTEGEEANATVDDDEEGEEEL
ASLDEETIQTGSQTNDEEDDDDEEVGEEGMVDIDTEDSEADVKSSTYGTADGK
PEEAESLDGWDADHDQVQDTTMTSSTYYNVSEESDTDEHHDSKAEAKEPPQNSDK
SDESEAVGHTPRTRSRGTVKINLWTLDVSPVANALNKSSANRSLKKAPRTESTPK
ESQSEPRRKITQPKLPKKEETNNKSNSNIGTLHRWISKSPRVMLRSTPVTAAASSS
45 AAVSGVSGGNASSSGTAR

>16:09:09|GENSCAN_predicted_CDS_3|7494_bp

atgaatgaaggaattcagcaggagggggcatgaaggctcagccggccctcctgctgtgccagacgcgtaactccaca

gttcacaaattaatgctccaacgcccgtataagtagcgaaaacgatgcaagacggaaaaccattttctacctcacaccggcgac
 gaattcaggagcagcaggagcgctctaccctgacaagcaaaaccacagcctcggcgccacgacgacctccagaacaaca
 gttacagcatcaactacttctggtcagcaactaataagggatccattgtcaaaagatttggccactcatgtaaaaagcacagtacaa
 aagcaaaagcattgccaatgggaagacggagcccagggaagaaactgaagcagaggatccctacaaagtacaggagctgattc
 5 agatgcgcaaggagcagcgattggcagcgcttaaacgtatggcaatgataaatgctgcggaacggatgccactcccatatacg
 gcgaagattgtcgcgaggtatatacagcgctgcatgcaggcgacccgatccctaaagcgtacaaactggcagacgctgggatac
 gccaaactgctgcactgccatggcgcatcggaacggttgggtccctaaaccacttgcgaagagcttcgaggaaaggtgcgctgac
 taaagccagtgttggcaacttggatctacgttctctgttggcgccccggatccgctgtatgtacaaaatctctacgcacgc
 actggcagcacgaacaaaggattgaaaacattgtgcatcaggccctgcggcctaagctggcgttgctgcatccaatcatttcgga
 10 aatgaccactaagttcccagatccgcgtctcatccaatacactgttgcaagttgcagaccatggatcgttggctacgccagctaaa
 gggttaacgggcatcgtgtactgatactcactcagatgaccaagatgttggatgtttggaaagcttttctcaactaccacggctcatattat
 ctgctgttagatggctctactcgggtggaacacggcagatccgtatggagcggttaattggagataaacgaatcttctgtctc
 ctctccacgcggctggttgagtggtggcatcaatttgacgggtgccgatactgtgacttttactcactccgactggaacccacaatg
 gatgcgcaggcccaagatcgttgcctcgtatgtgaacgcgagatgtacatactaccgtctgtctccgaaagaacataga
 15 gggttaacattcttaagaaggcaaaccaaaagcgaatgtgagcgacatggccatcgagggtggcaactttacaactacgtacttta
 agagttccaccataaaggatctctttacaatggagcagagcgagcaggacagtcgagccaagagaagtcggaaaacaaggat
 agaattgtgtctacaacaacgcttcagatacgcttcgacggttggagacggagaagcagtcactgcgtcatttgagcacgc
 gttggctgccgccgaggacgagcaggatgtgcaggccacgaaaacggctaaagccgaagtggcagctgatctgcccaggttc
 gacgagaacattcctattgcaacagaagatccaatgcggaaggaggtcctcaagtggaaactcagcaaggccgatctggagatg
 20 cagaacttggttaaacagctctcaccgatagagcgatagccatgcgcttgggaagaaactggagcagcatggacggcgga
 caattgcgagccgagggaagcggagctggaggcccagaaacgcgagtgaggaggccaatcgcttggcgccatgcacaaggga
 ggaggagctgttgaaagcaagaaacggaagcggaggagatgcttacctacagtcgcaaggattcgagtaacagggttaatac
 aacagattccaatccaataagcgacgactggtgagggaataatcgcaaaactcagctcagaagctgagcaggagtgtagcag
 ccatagcaccggtagcaacaacaagaacagtaaatcggaacacccgtggaataagccagaacagcctcaatcagactgtac
 25 cagtggggtcagggaataagcagggtgaatagaacggggggggggtcagtagcagtagccgagggcaaaagtaacagcacga
 agtcaacggggaagggaacagacgccgcaccgcaagtccggcgagacccgtctccactctctggcgagctcaatatggc
 cagcgcggcgaacaccgcccactagaagacaacacgtacagctctggctgcatctgcagctgcatctacttttagaggatgcctct
 tgatcgtcgaggagcgtccaaaagacagtcggccaacatagctatgagcaagatgatgaagacgcccttcaaacagaatgttc
 catccaacatcagfataaagacaactcctctaaaggggcggaagagacagtggtgcagctgccgccacacgcagtaaaactgc
 30 tggaagaagagctacaattgtctcctttaaacaatagtgatgatgaaagtaccaggatgaagaggagcagggaagagcagg
 agtctgaagaagataccgaggggcagggaagcaaatgccacagtagacgacgacgaggagggggaggaggaggttggcgtca
 ctgacgaagagaccatacaaacggatcgaaacaaatgatgaagaagacgatgacgaggaagaagtggtgaagaggga
 tgggtgataatgatactgaagattcagaggcagatgtcaaatccagctccacctatggtacagcggcagatggttaagcccgaagaa
 gccgaagcttggtggtggctgggtgacacgaccaggtgcaggacaccacaatgactagctccactactacaatgtcagcga
 35 ggaatcagacacggatgagcatcacgatagcaaggcggaggctaaagagccgccgcaaaattccgataagagcgacgagag
 cgaggctgttgacacacaccacgtacaaggtcgcgcggcacagtaaaagatcaatctgtggaccctggacgtgagtcctcgtagc
 aaacgcattgaataaaagcagcgccaataggagcctcaaaaaagcaccgaaggactgagtcacgccaagaggagctcagagc
 gagccaaggcgaaagattactcagccaaagctgccgaagaagaagaactaacaacaaatctaacagaatattggcacctta
 caccgctggatagcgaagtcccccgagtaatgctcgtatccacacctgttacggcagcgagcgtagctcatcagcagcagtc
 40 gtggtgtttcgggagggaatgcctcctcagcggaacagccaggtga

Drosophila Gene Hit TBLASTN with ORF2: brahma protein (M85049) and imitation-SWI protein (ISWI) (L27127) and chromodomain-helicase-DNA-binding (CHD-1)

45 **Human Homologue** BLASTX with EST TBLASTN with ORF2: Snf2-related CBP activator protein (SRCAP) (AF143946) and SWI/SNF related, matrix associated, actin dependent regulator of chromatin, subfamily a, member 4 (SMARCA4) (NM_003072.1)

	<i>Drosophila</i> EST	several including SD07794 (AI534784), LD34465 (AA990657)
	Annotated <i>Drosophila</i> genome genomic segment	AE003453
5	Annotated <i>Drosophila</i> genome Complete gene candidate	CG9696 – domino an enzyme involved in DNA repair homology to snf2 family helicases
10	Human homologue of Complete gene candidate	CG9696- gi4557447 416409C913D6A935 [ref]NP_001261.1 chromodomain helicase DNA binding protein 1 [Homo sapiens] (1.90E-85
15	Putative function	snf2 helicase family member protein that contains a chromodomain, which occurs in proteins that are implicated in chromatin compaction, and an SNF2/SWI2-like helicase domain, which occurs in proteins 20 that are believed to activate transcription by counteracting the repressive effects of chromatin structure
25	Confirmation by RNAi	Loss of G1, peak, increase in G2M indicating arrest in G2/M

Example 70 (Category 5)

Line ID 99/31
Category 2nd chromosome, small imaginal discs
5 **Reversion** NR
Map Position 53E

Rescue ID EcoR1

Rescue Sequence 1

10 AAGGCCCGACCAGAAACGAAATTTTCGGCGCGTNTTTTAAAATGCGCGGTAA
ATTCACTTTGATTTTTGTTTTCTCTCTCGTTCTTCACACACACAGTTAGTTAGA
TTGTGTGTTTCGCCTGGCTTTGCCTTTTAATTTTTTATTACCTGCATCCGATTTCG
GTATTTGAAACAGCCGTTGAGTCTCCTTTGGCTTTTTTATCAGCGACGTCATCA
GTGGCGGCAGAAGCAGAAGCGTCGACAGCGGCGGGGGATTTCGGCTGCATCTT
15 TGGAGCCCCTTTCCGGCTGTGCCCCACGGCTTTCGCCACCCCCGCAGTAACC
GATGCATTTTCCACATCGCTTACCTTATCGGCGGCATTTTCTTTGGCTGCCGTT
TCTGCCGCTTTGTTAGCATCCTTTTCGTGCGGCGANGGCATGGAAAGATACAA
ATCAGAATTGGATTACACTTGCTAATTTTTTGGCGGNCAATACAATGGTTCGG
TGCGCCTATTCTTTTTTAATCGAATCGCAATTGAGTGTNAATTAAGTCTCCGCA
20 ATGCAATTTGTGTATCTGTCTCCTCCCCGANC GAACAACGATNGAAAAAGGAA
CCAGAAATAAAANAGGNAATGAAAAACACATTGCAATCTATAAGGCCACAC
ACACACATATCATCCCGTCTACCANTCCATCGGATTTCGANCCACANAANCCAT
NTTTATACCNCAACGAACGNGGAAAAAACNATATCNGNAATTACCCCCCGAA
AATTGTTGCCNCTTTTACCCAAATATTTACAACCNCCGTTCAATCACTCCTGGA
25 ACATTCCNGGCTTTCCCAATTTTCNCCTTTACTACAATTTCAATGGTTTCTTTTT
CCTCAC

Rescue ID BamH1

Rescue Sequence 2

30 CCTNAAATGTNGCGCTGGGNCCTAAANCGTCNCTCCTTGTGTCTCTCTTGTTTA
CCGCGCTATGCTGATGTTGGCATGTGGTTCGATCCCCCTCCGTGTCGATGTTTA
CAAAACCATNATTAGAGTTTGATGATTGAGTTCTCTTAACTTTCCTTCCTCCTT
CCTTCCTTGGCTTTTGTTCATGCTAAATCCTTTAAATGGGGTTCTGCGTAGTTT
AATGCCGAGGTACAGCAAACTTCAATATTCATGTTCCCTTGCCTGCCAAAC
35 GAAATTAGCATTGGACGTCCCAAGGTTGAAGACATTTNATTATTTTAACATCT
TTTNATTTTATTACATTTGAACTCTTACAAGTAATAATAATTACAATTAATAT
TATAGCTGCAGCGACAAAAAGGAGAAATCCCCCTCGCCGGTAATAAAGAAT
CCAACAATAAGGATGCTNAAAANGAAGAAAACCCNAAAAAGGAGAAGAAAA
ATCGGAANAAGGNGATGAGCCNGAAGATGAGGNNGATGAGAAAGCTAGCGA
40 TGAAGAGAGCGAGAAGAAGAAANCGANATGAGATGCAGAGGACAGATAAAG
GATGCCACNGATGAATCCAAGCCAAAATCGGGAGCCGATAAGCCCAAGAAAC
TGAGCCCAAGGCCAAGAATGGCAAGGTGGNT

Genomic hit, Accession No. CSC:AC020063

45

Associated ORF

Genscan ORF1 predicted sequences >16:48:25|GENSCAN_predicted_peptide_1|722_aa
 MPSPHEKDANKAAETAAKENAADKVSDVENASVTAGVAKAVGAQPERGSKDA
 AESPAAVDASASAATDDVADKKAKGDSTAVSNTESDAAAADKKEKSPSPVIKKS
 NNKDAKKEDNSEKDEENSEDGDEPEDEADEKASDEESEKKKPKLDAEDKIKDAT
 5 DESKPKSGADKPKKPEPKAKNGKVAKEEDDDEEDEDDEDAEDDDGDENDGLDK
 NNEVAEDDENVVALAIEDRINENINKTRVDGLQTLHAICFGAQGNVVKKNLRS
 FAGFEFAKDSAEYNKKLEAIKKVDNKGRLSICEILTLDRKGSKNETVLRVLKFLM
 EPDESLCLEQGDEEEEDAEDEDLDEDEEDPPSEEDKKRKSGKSSGGAGRGSARN
 STGRPRRATAGKKMSAYVDFSSDDSEQKVA VPKRRRNDDES GSDYNPSANS
 10 SDGGRGGGAGAAGRKVP SRGGRGRPARKSRRRNSDSEEEEESEVSDADSDVPKR
 KRGSVGKRGRPAAPASAGRRGRGRGAASRK RKDS DSEDEEVSEDEEEEDVSDFA
 SDQSEVCKFNLISSIWCFIKYMPIFQEERPKKSKKPITPAKNSKANNKSKPAGKADS
 RSKKSKKESSEEDDDVDDKDESEDEPLTKKGKQAFPTDEQIRGYVKEILDKANL
 EEITMTKTVCKQVYAKYPDFDLTDKKDFIKATVKADGVQDLDGSPELIPRGRTTVT
 15 IWLICCCNNQIFGET

>16:48:25|GENSCAN_predicted_CDS_1|2169_bp
 atgccatgccgcacgaaaaggatgctaacaaagcggcagaaacggcagccaaagaaaatgccgccgataaggaagcgcgatg
 tggaaaatgcatcggttactcgggggtggcgaaagccgtgggggcacagccgaaaggggtccaaagatgcagccgaatc
 20 ccccgccgtgtgcagcctctgcctctgcgccactgatgacgtcgtgataaaaaagccaaaggagactcaacggctgtttca
 aataccgaatcgatgcagctgcagcggacaaaaaggagaatccccctgccggtataaagaagtccaacaataaggatgc
 taaaaaggaggacaactccgaaaaggacgaggagaactcggaagacggcgatgagccagaagatgaggctgatgaaagc
 tagcgtatgaagagagcgaagaagaaccgaaattagatgcagaggacaagataaaggatgccactgatgagccaagcca
 aaatcgggagccgataagcccaagaacctgagccaaaggccaagaatggcaaggtggctaaggaggaggacgacgacga
 25 agaggacgaggatgatgaggatgccgaagatgacgatggagacgagaacgatggcctggacaagaacaacagggtggccg
 aggatgatgagaatgtcgtcgtctgccgagattgatgcattaatgagaatatcaacaagactcgtgtagatggtctgcaacat
 tgcatgcaatctgcttggcgcccaaggcaagaacaatgtgtgcaagaagaacttgcgatcctttgccggttcgagtttgccaagg
 attcagcggaggtacaacaaaagctggaggccatcaaaaagggtgataataaggcctgcgcagcatctgcgagatccttacc
 tcgatcgcaaggccagcaagaacgagactgtcttcgagtgctcaattcctaattggaaccggacgagtcgcttcttgaggca
 30 gggatgatgaggaggaggaggaatgccgaggacgaggatctggaatgaggaggacccgccagtgaaaggagaca
 agaagcgcaagagcggaaagtctagcggcggtggcagaggctctgcacgaattccaccggacgtccaaggcgcgca
 cggcaggaaagaaaatgtccgctatgtagatttccagctctgacgatagcagcagaaagttgagttcccaaaaggagacg
 aatgatgactccgagtcgggctcagattacaatccttctgccaattccgactctgacgggtgctggtggtggtgctggtgcagc
 aggtcgcaaaagtcgaagccgcgggtggacgcggtcgtcctgcgcgcaaaagtcgcagaagaaactctgattccgaggagaa
 35 gaggaaatcggaagtttcgatgccgatagtgatgtccaaacgtaaacgtggttccgtgggtaaacgtggacgacccggcagct
 cctgcgtcagctggacgaaggggtagaggacgagggtgcagcttcccgaagcgtaaagattcagatagcgaagatgaggagg
 tatccgaggatgaagaggaggaggtgtctccgatttgcagcgatcaaaagcgaagtgtaaaatttaattatcgagcatttg
 gtgtttatcaagtatatgccaattttcaggagggaacgtcccaaaagagcaagaagccattacgcctgcgaaaaatagcaaa
 ctaacaacaagtcaaaaccagctggaaaggccgatagtcgatcgaaaaatcaagaagggaatctccgaggaagatgatgat
 40 gtcgatgacaaagatgaatccgacgaggatgagccactaaccaaaagggaacaggcattccaacggatgaacaaatacc
 cggatgtcaaaagagattctggataaagccaatctgaggagattacgatgaaaccgtgtgcaacaagtttatgcaaaatacc
 agactttgacctaacagacaagaagacttcatcaagcgacagtgaaagcggacggaggtcaggatttgatggtagtcgccga
 actgatccgcgtggccgaacaacggttacaatatggttgatctgctgttgcaacaatcagatatttggggagacgtaa

45 **Human Homologue** TBLASTN with ORF1: poor homology with DEK gene
 (D6S231E) (NM_003472.1)
Drosophila EST several including LD33301 (AA979048)

	Annotated <i>Drosophila</i> genome genomic segment	AE003805
	Annotated <i>Drosophila</i> genome Complete gene candidate	CG5935 - EG:EG0003.6 - novel with weak homology to DEK oncogene
5		CG8648 - EG:EG0003.3 - novel XPG/ flap endonuclease-like, DNA repair?
10	Human homologue of Complete gene candidate	CG5935- 1e-17 4503249 ref NP_003463.1 pD6S231E DEK gene >gi 544150 sp P35659 DEK_H UMAN DEK PROTEIN >gi 284375
15		CG8648- 4758356 ref NP_004102.1 pFEN1 flap structure-specific endonuclease 1;
20		MATURATION FACTOR 1 (MF1); DNase IV; RAD2_HUMAN(aa)
25	Putative function	CG5935: function unknown but putative DNA-binding protein predicted to be involved in chromosomal organisation. The translocation (6;9), associated with a specific subtype of acute myeloid leukemia, results in the fusion of two genes, dek and can, and the expression of a chimeric, leukemia-specific dek-can mRNA
30		CG8648: Novel XPG/ flap endonuclease-like, DNA repair protein
	Confirmation by RNAi	Both show slight reduction of G1 peak

REFERENCES

- Deak, P., Omar, M.M., Saunders, R.D.C., Pal, M., Komonyi, O., Szidonya, J., Maroy, P., Zhang, Y., Ashburner, M., Benos, P., Savakis, C., Siden-Kiamos, I., Louis, C., Bolshakov, V.N., Kafatos, F.C., Madueno, E., Modolell, J., Glover, D.M. (1997)
- 5 Correlating physical and cytogenetic maps in chromosomal region 86E-87F of *Drosophila melanogaster*. Genetics 147:1697-1722.
- Torok, T., Tick, G., Alvarado, M., Kiss, I. (1993) P-lacW insertional mutagenesis on the second chromosome of *Drosophila melanogaster*: isolation of lethals with different overgrowth phenotypes. Genetics 135(1):71-80
- 10 Saunders, R.D.C., Glover, D.M., Ashburner, M., Siden-Kiamos, I., Louis, C., Monastirioti, M., Savakis, C., Kafatos, F.C.(1989) PCR amplification of DNA microdissected from a single polytene chromosome band: a comparison with conventional microcloning. Nucleic Acids Res. 17:9027-9037
- Lefevre, G. (1976) A photographic representation and interpretation of the
- 15 polytene chromosomes of *Drosophila melanogaster* salivary glands. In: The Genetics and Biology of *Drosophila*, Eds Ashburner, M. and Novitski, E. Academic Press.
- Jowett, T. (1986) Preparation of nucleic acids. In "*Drosophila*: A Practical Approach." Ed Roberts, D.B. IRL Press Oxford.
- Pirrotta, V. (1986) Cloning *Drosophila* genes. In: . In "*Drosophila*: A Practical
- 20 Approach." Ed Roberts, D.B. IRL Press Oxford.
- Altschul, S.F. and Lipman, D. J. (1990) Protein database searches for multiple alignments. Proc. Natl. Acad. Sci. USA 87: 5509-5513

Burge, C. and Karlin, S. (1997) Prediction of complete gene structures in human genomic DNA. J. Mol. Biol. 268, 78-94.

Each of the applications and patents mentioned above, and each document cited or referenced in each of the foregoing applications and patents, including during the prosecution of each of the foregoing applications and patents ("application cited documents") and any manufacturer's instructions or catalogues for any products cited or mentioned in each of the foregoing applications and patents and in any of the application cited documents, are hereby incorporated herein by reference. Furthermore, all documents cited in this text, and all documents cited or referenced in documents cited in this text, and any manufacturer's instructions or catalogues for any products cited or mentioned in this text, are hereby incorporated herein by reference.

Various modifications and variations of the described methods and system of the invention will be apparent to those skilled in the art without departing from the scope and spirit of the invention. Although the invention has been described in connection with specific preferred embodiments, it should be understood that the invention as claimed should not be unduly limited to such specific embodiments. Indeed, various modifications of the described modes for carrying out the invention which are obvious to those skilled in molecular biology or related fields are intended to be within the scope of the following claims.

CLAIMS

1. A polynucleotide selected from:
 - (a) polynucleotides comprising any one of the nucleotide sequences set out in Examples 1 to 70 or the complement thereof.
 - 5 (b) polynucleotides comprising a nucleotide sequence capable of hybridising to the nucleotide sequences set out in Examples 1 to 70, or a fragment thereof.
 - (c) polynucleotides comprising a nucleotide sequence capable of hybridising to the complement of the nucleotide sequences set out in Examples 1 to 70 or a fragment thereof.
 - 10 (d) polynucleotides comprising a polynucleotide sequence which is degenerate as a result of the genetic code to the polynucleotides defined in (a), (b) or (c).
2. A polynucleotide selected from:
 - (a) polynucleotides comprising any one of the nucleotide sequences set out in Examples 1 to 14 or the complement thereof.
 - 15 (b) polynucleotides comprising a nucleotide sequence capable of hybridising to the nucleotide sequences set out in Examples 1 to 14, or a fragment thereof.
 - (c) polynucleotides comprising a nucleotide sequence capable of hybridising to the complement of the nucleotide sequences set out in Examples 1 to 14 or a fragment thereof.
 - 20 (d) polynucleotides comprising a polynucleotide sequence which is degenerate as a result of the genetic code to the polynucleotides defined in (a), (b) or (c).
3. A polynucleotide selected from:

- (a) polynucleotides comprising any one of the nucleotide sequences set out in Examples 15 to 19 or the complement thereof.
- (b) polynucleotides comprising a nucleotide sequence capable of hybridising to the nucleotide sequences set out in Examples 15 to 19, or a fragment thereof.
- 5 (c) polynucleotides comprising a nucleotide sequence capable of hybridising to the complement of the nucleotide sequences set out in Examples 15 to 19 or a fragment thereof.
- (d) polynucleotides comprising a polynucleotide sequence which is degenerate as a result of the genetic code to the polynucleotides defined in (a), (b) or (c).
- 10 4. A polynucleotide selected from:
 - (a) polynucleotides comprising any one of the nucleotide sequences set out in Examples 20 to 30 or the complement thereof.
 - (b) polynucleotides comprising a nucleotide sequence capable of hybridising to the nucleotide sequences set out in Examples 20 to 30, or a fragment thereof.
 - 15 (c) polynucleotides comprising a nucleotide sequence capable of hybridising to the complement of the nucleotide sequences set out in Examples 20 to 30 or a fragment thereof.
 - (d) polynucleotides comprising a polynucleotide sequence which is degenerate as a result of the genetic code to the polynucleotides defined in (a), (b) or (c).
- 20 5. A polynucleotide selected from:
 - (a) polynucleotides comprising any one of the nucleotide sequences set out in Examples 31 to 53 or the complement thereof.
 - (b) polynucleotides comprising a nucleotide sequence capable of hybridising to the nucleotide sequences set out in Examples 31 to 53, or a fragment thereof.

- (c) polynucleotides comprising a nucleotide sequence capable of hybridising to the complement of the nucleotide sequences set out in 31 to 53 or a fragment thereof.
- (d) polynucleotides comprising a polynucleotide sequence which is degenerate as a result of the genetic code to the polynucleotides defined in (a), (b) or (c).
- 5
6. A polynucleotide selected from:
- (a) polynucleotides comprising any one of the nucleotide sequences set out in 54 to 70 or the complement thereof.
- (b) polynucleotides comprising a nucleotide sequence capable of hybridising to the nucleotide sequences set out in 54 to 70, or a fragment thereof.
- 10
- (c) polynucleotides comprising a nucleotide sequence capable of hybridising to the complement of the nucleotide sequences set out in 54 to 70 or a fragment thereof.
- (d) polynucleotides comprising a polynucleotide sequence which is degenerate as a result of the genetic code to the polynucleotides defined in (a), (b) or (c).
- 15
7. A polynucleotide probe which comprises a fragment of at least 15 nucleotides of a polynucleotide according to any of Claims 1 to 6.
8. A polypeptide which comprises any one of the amino acid sequences set out in Examples 1 to 70 or in any of Examples 1 to 14, Examples 15 to 19, Examples 20 to 30, Examples 31 to 53 and Examples 54 to 70, or a homologue, variant, derivative or fragment thereof.
- 20
9. A polynucleotide encoding a polypeptide according to Claim 8.
10. A vector comprising a polynucleotide according to any of Claims 1 to 7 and 9.

11. An expression vector comprising a polynucleotide according to any of Claims 1 to 7 and 9 operably linked to a regulatory sequence capable of directing expression of said polynucleotide in a host cell.
12. An antibody capable of binding a polypeptide according to Claim 8.
- 5 13. A method for detecting the presence or absence of a polynucleotide according to any of Claims 1 to 7 and 9 in a biological sample which comprises:
- (a) bringing the biological sample containing DNA or RNA into contact with a probe according to Claim 9 under hybridising conditions; and
 - (b) detecting any duplex formed between the probe and nucleic acid in the
10 sample.
14. A method for detecting a polypeptide according to Claim 8 present in a biological sample which comprises:
- (a) providing an antibody according to Claim 12;
 - (b) incubating a biological sample with said antibody under conditions which
15 allow for the formation of an antibody-antigen complex; and
 - (c) determining whether antibody-antigen complex comprising said antibody is formed.
15. A polynucleotide according to according to any of Claims 1 to 7 and 9 for use in therapy.
- 20 16. A polypeptide according to Claim 8 for use in therapy.
17. An antibody according to Claim 12 for use in therapy.

18. A method of treating a tumour or a patient suffering from a proliferative disease comprising administering to a patient in need of treatment an effective amount of a polynucleotide according to any of Claims 1 to 7 and 9.
19. A method of treating a tumour or a patient suffering from a proliferative disease,
5 comprising administering to a patient in need of treatment an effective amount of a polypeptide according to Claim 8.
20. A method of treating a tumour or a patient suffering from a proliferative disease, comprising administering to a patient in need of treatment an effective amount of an antibody according to Claim 12 to a patient.
- 10 21. Use of a polypeptide according to Claim 8 in a method of identifying a substance capable of affecting the function of the corresponding gene.
22. Use of a polypeptide according to Claim 8 in an assay for identifying a substance capable of inhibiting the cell division cycle.
- 15 23. Use as claimed in Claim 22, in which the substance is capable of inhibiting mitosis and/or meiosis.
24. A method for identifying a substance capable of binding to a polypeptide according to Claim 8, which method comprises incubating the polypeptide with a candidate substance under suitable conditions and determining whether the substance binds to the polypeptide.
- 20 25. A method for identifying a substance capable of modulating the function of a polypeptide according to Claim 8 or a polypeptide encoded by a polynucleotide according to any of Claims 1 to 7 and 9, the method comprising the steps of: incubating the polypeptide with a candidate substance and determining whether activity of the polypeptide is thereby modulated.

26. A substance identified by a method or assay according to any of Claims 21 to 25.
27. Use of a substance according to Claim 26 in a method of inhibiting the function of a polypeptide.
28. Use of a substance according to Claim 26 in a method of regulating a cell division
5 cycle function.